

Gulf St Vincent Prawn *Penaeus (Melicertus) latisulcatus* Fishery 2017/18



L.J. McLeay, C. L. Beckmann and G. E. Hooper

**SARDI Publication No. F2007/000782-8
SARDI Research Report Series No. 999**

**SARDI Aquatics Sciences
PO Box 120 Henley Beach SA 5022**

October 2018

Fishery Assessment Report to PIRSA Fisheries and Aquaculture

**Gulf St Vincent Prawn
Penaeus (Melicertus) latisulcatus
Fishery 2017/18**

Fishery Assessment Report to PIRSA Fisheries and Aquaculture

L.J. McLeay, C. L. Beckmann and G. E. Hooper

**SARDI Publication No. F2007/000782-8
SARDI Research Report Series No. 999**

October 2018

This publication may be cited as:

McLeay, L.J., Beckmann, C. L. and Hooper, G. E. (2018). Gulf St Vincent Prawn *Penaeus (Melicertus) latisulcatus* Fishery 2017/18. Fishery Assessment Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/000782-8. SARDI Research Report Series No. 999. 44pp.

South Australian Research and Development Institute

SARDI Aquatic Sciences
2 Hamra Avenue
West Beach SA 5024
Telephone: (08) 8207 5400
Facsimile: (08) 8207 5415
<http://www.pir.sa.gov.au/research>

DISCLAIMER

The authors warrant that they have taken all reasonable care in producing this report. The report has been through the SARDI internal review process, and has been formally approved for release by the Research Director, Aquatic Sciences. Although all reasonable efforts have been made to ensure quality, SARDI does not warrant that the information in this report is free from errors or omissions. SARDI and its employees do not warrant or make any representation regarding the use, or results of the use, of the information contained herein as regards to its correctness, accuracy, reliability and currency or otherwise. SARDI and its employees expressly disclaim all liability or responsibility to any person using the information or advice. Use of the information and data contained in this report is at the user's sole risk. If users rely on the information they are responsible for ensuring by independent verification its accuracy, currency or completeness. The SARDI Report Series is an Administrative Report Series which has not been reviewed outside the department and is not considered peer-reviewed literature. Material presented in these Administrative Reports may later be published in formal peer-reviewed scientific literature.

© 2018 SARDI

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968* (Cth), no part may be reproduced by any process, electronic or otherwise, without the specific written permission of the copyright owner. Neither may information be stored electronically in any form whatsoever without such permission.

SARDI Publication No. F2007/000782-8
SARDI Research Report Series No. 999

Author(s): L.J. McLeay, C. L. Beckmann and G. E. Hooper

Reviewer(s): S. Mayfield, C. Noell (SARDI) and S. Shanks (PIRSA)

Approved by: S. Mayfield
Science Leader - Fisheries

Signed:

Date: 25 October 2018

Distribution: PIRSA Fisheries and Aquaculture, SAASC Library, Parliamentary Library, State Library and National Library

Circulation: Public Domain

TABLE OF CONTENTS

LIST OF FIGURES.....	IV
LIST OF TABLES	V
ACKNOWLEDGMENTS	VI
EXECUTIVE SUMMARY	1
1. INTRODUCTION.....	3
1.1. Overview	3
1.2. Description of the fishery.....	3
1.2.1. Access	3
1.2.2. Management arrangements	6
1.3. Biology of the Western King Prawn	9
1.3.1. Distribution and habitat preferences	9
1.3.2. Reproductive biology.....	10
1.3.3. Early life history	10
1.3.4. Stock structure	11
1.3.5. Growth	11
1.3.6. Natural mortality	11
1.3.7. Biosecurity and prawn health	12
1.4. Research program.....	12
1.5. Information sources used for assessment	13
1.6. Stock status classification.....	13
2. METHODS	16
2.1. Fishery-independent surveys	16
2.2. Fishery statistics	19
2.2.1. Catch and effort.....	19
2.2.2. Prawn size	19
2.3. Catch rate standardisation.....	20
3. RESULTS	21
3.1. Fishery-independent surveys	21
3.1.1. CPUE.....	21
3.1.2. Recruitment.....	23
3.1.3. Size.....	23
3.1.4. Spatial patterns in CPUE and size	24
3.2. Fishery statistics	26
3.2.1. Catch and effort.....	26
3.2.2. CPUE.....	29
3.2.3. Spatial patterns in catch and effort	30
3.2.4. Prawn size	33
4. DISCUSSION.....	35

4.1. Information sources used for assessment	35
4.1.1. Fishery-independent data.....	35
4.1.2. Fishery-dependent data	35
4.2. Status	36
4.3. Future directions	37
5. REFERENCES.....	39

LIST OF FIGURES

Figure 1. Double-rig trawl gear and location of hopper sorting and prawn grading systems used in the Gulf St Vincent Prawn Fishery.	4
Figure 2. Trawl net configuration showing trawl boards, head rope, ground chain and codend with crab bag as used in the Gulf St Vincent Prawn Fishery.....	4
Figure 3. The research management blocks for the Gulf St Vincent Prawn Fishery.....	5
Figure 4. The 109 survey stations specified in the 2017 Management Plan and regions of the Gulf St Vincent Prawn Fishery.	18
Figure 5. Key outputs from May fishery-independent surveys. (a) Nominal catch per unit effort (CPUE, kg.trawl-shot ⁻¹) from diamond v T90-mesh; (b) Standardised versus nominal catch per unit effort (CPUE, kg.trawl shot ⁻¹); (c) the Fishery Recruitment Index (FRI) (recruits.h ⁻¹), and (d) prawn size (prawns.kg ⁻¹)	22
Figure 6. Catch rate and mean size of prawns sampled from the diamond-mesh codend during May FIS' in the GSVPF between 2011/12 and 2017/18.	25
Figure 7. Fishery-dependent catch and effort data outputs for the Gulf St Vincent Prawn Fishery; (a) Annual catch (t) and (b) effort (vessel-nights) separated by early spawning (November–December), late spawning (January–March) and non-spawning (April–October) and (c) annual commercial effort (hours * 1000).....	27
Figure 8. Comparison of standardised annual commercial CPUE and nominal annual commercial CPUE (kg.block ⁻¹ .vessel-night ⁻¹) in the Gulf St Vincent Prawn Fishery from 1990/91–2017/2018.	30
Figure 9. The number of blocks fished pre–Christmas (November–December) and all other months in the Gulf St Vincent Prawn Fishery from 1989/90–2017/18.....	31
Figure 10. The annual catch per fishing block in the Gulf St Vincent Prawn Fishery from 2007/08–2017/18.....	32
Figure 11. Size-grade composition (%) of monthly harvests in the Gulf St Vincent Prawn Fishery from 2005/06–2017/18.	33

LIST OF TABLES

Table 1. Key Gulf St Vincent Prawn Fishery statistics between 2015/16 and 2017/18.	2
Table 2. Major management milestones for the Gulf St Vincent Prawn Fishery.	8
Table 3. Current management arrangements for the Gulf St Vincent Prawn Fishery.....	9
Table 4. Stock status terminology (Flood <i>et al.</i> 2014).....	14
Table 5. Decision rules for classifying the status of Western King Prawn stock in GSV	15
Table 6. Categories assigned to reported prawn grades from commercial logbook data for the Gulf St Vincent Prawn Fishery.	20
Table 7. Analysis of deviance (Type II test) for the GLM used to standardise survey CPUE for the GSVPF.....	23
Table 8. Monthly distribution of commercial (and survey) catch in the Gulf St Vincent Prawn Fishery from 1989/99–2017/18.	28
Table 9. Analysis of deviance (Type II test) for the GLM used to standardise annual commercial catch in the Gulf St Vincent Prawn Fishery.	29
Table 10. Size-grade composition (t) of monthly harvests in the Gulf St Vincent Prawn Fishery from 2005/06–2017/18.	34

ACKNOWLEDGMENTS

Funds for this research were provided by PIRSA Fisheries and Aquaculture, obtained through licence fees. SARDI Aquatic Sciences provided substantial in-kind support. Thanks go to the Gulf St Vincent prawn fishers for their substantial contributions to surveys (including vessel time and personnel). Observer work during the May survey was undertaken by Peter Hawthorne, Andrew Hogg, Doug Graske, Sarah-Lena Reinhold, Troy Rogers, Nat Navong and Stuart Sexton. The catch and effort data from the SARDI Information Management System were provided by Ms Melleessa Boyle of the Information Systems and Database Support Unit at SARDI Aquatic Sciences, and we thank Dr Craig Noell for his assistance with statistical procedures. The report was formally reviewed by Dr Craig Noell and Dr Stephen Mayfield of SARDI Aquatic Sciences, and Steve Shanks of PIRSA Fisheries and Aquaculture and approved for release by Dr Stephen Mayfield, Science Leader, Fisheries (SARDI Aquatic Sciences).

EXECUTIVE SUMMARY

This report determines the current status of the Western King Prawn (*Penaeus (Melicertus) latisulcatus*) stock in Gulf St Vincent (GSV) through analysis of data collected by several long-term monitoring programs.

In 2017/18, the total commercial catch of Western King Prawn in the Gulf St Vincent Prawn Fishery (GSVPF) was 235.3 t, with an additional 3.5 t taken in the May 2018 fishery-independent survey (FIS). Fishing was conducted over 291 vessel-nights, comprising 97% of the Total Allowable Commercial Effort (TACE) of 300 vessel-nights. All 50 allocated pre-Christmas vessel-nights were fished (Table 1).

Estimates of standardised annual commercial catch per unit effort (CPUE) have been relatively high in recent years and were $\geq 750 \text{ kg.block}^{-1}.\text{vessel-night}^{-1}$ in 2014/15, 2015/16, 2016/17 and 2017/18.

Estimates of standardised FIS CPUE have also remained at relatively high levels since 2013/14, exceeding $30 \text{ kg.trawl-shot}^{-1}$ in the last five May surveys. Similarly, the FIS recruitment index (FRI) has also been relatively high ($\geq 600 \text{ recruits.h}^{-1}$) for four of the last five annual surveys (2013/14, 2015/16, 2016/17 and 2017/18).

Primary Industries and Regions South Australia (PIRSA) has adopted the National Fishery Status Reporting Framework (NFSRF) to determine the status of all South Australian fish stocks. The current management plan for the GSVPF provides the decision rules for classifying stock status relative to limit, trigger and target reference points defined for three performance indicators relating to stock abundance and recruitment (PIRSA 2017). The performance indicators are: 1) standardised annual CPUE; 2) standardised FIS CPUE; and 3) the FRI.

In 2017/18:

- Standardised annual commercial CPUE was $895 \text{ kg.block}^{-1}.\text{vessel-night}^{-1}$, which was 0.7% higher than in 2016/17 ($889 \text{ kg.block}^{-1}.\text{vessel-night}^{-1}$), and within the target range defined for this performance indicator;
- Standardised FIS CPUE was $38.6 \text{ kg.trawl-shot}^{-1}$, which was 1.1% higher than in 2016/17 ($38.2 \text{ kg.trawl-shot}^{-1}$), and within the high range defined for this performance indicator; and
- The FRI was $670.3 \text{ recruits.h}^{-1}$, which was 14.6% lower than in 2016/17 ($784.4 \text{ recruits.h}^{-1}$), but within the high range defined for this performance indicator.

When the 2017/18 estimates for each performance indicator are applied in the decision matrix used to classify the stock status of Western King Prawn in GSV, the fishery is classified as '**sustainable**'.

Table 1. Key Gulf St Vincent Prawn Fishery statistics between 2015/16 and 2017/18. Note, estimates of standardised annual commercial CPUE and standardised FIS CPUE will differ each year as data are updated within the Generalised Linear Model.

Statistic	2015/16	2016/17	2017/18
Total allowable commercial effort (TACE)	300 nights (30 pre-Christmas)	300 nights (50 pre-Christmas)	300 nights (50 pre-Christmas)
Total commercial catch	217.8 t (+4.1 t from survey)	224.6 t (+2.9 t from survey)	235.3 t (+3.5 t from survey)
Total effort	296 vessel nights (30 pre-Christmas) 2,799 hours	287 vessel nights (49 pre-Christmas) 2,836 hours	291 vessel nights (50 pre-Christmas) 2,858 hours
Standardised annual commercial CPUE	929 kg.vessel.night ⁻¹	889 kg.vessel.night ⁻¹	895 kg.vessel.night ⁻¹
Standardised FIS CPUE	54.3 kg.trawl-shot ⁻¹	38.2 kg.trawl-shot ⁻¹	38.6 kg.trawl-shot ⁻¹
FRI	837.1 ± 115.6 recruits.h ⁻¹	784.4 ± 116.3 recruits.h ⁻¹	670.3 ± 82.8 recruits.h ⁻¹
Status	Sustainable	Sustainable	Sustainable

Keywords: Western King Prawn, *Penaeus (Melicertus) latisulcatus*, trawl fishery, Gulf St Vincent, South Australia.

1. INTRODUCTION

1.1. Overview

Stock assessments for the Gulf St Vincent Prawn Fishery (GSVPF) are part of the South Australian Research and Development Institute's (SARDI) (Aquatic Sciences) ongoing assessment program. This report assesses the current status of the Western King Prawn (*Penaeus (Melicertus) latisulcatus*) (Kishinouye 1996) stock in Gulf St Vincent (GSV) and includes new data from the 2017/18 fishing season.

The report has three objectives: 1) present information relating to the fishery and biology of Western King Prawn; 2) assess the 2017/18 status of the Western King Prawn resource in Gulf St Vincent; and 3) identify future directions for the research program.

1.2. Description of the fishery

1.2.1. Access

Three commercial prawn fisheries occur within South Australia: the Spencer Gulf Prawn Fishery (SGPF); the West Coast Prawn Fishery (WCPF); and the GSVPF. The SGPF is the largest prawn fishery in South Australia in terms of total catch and numbers of licences (39). The WCPF is the smallest of the prawn fisheries with three licences. There are currently ten commercial fishing licences issued for the GSVPF.

All three prawn fisheries use demersal otter trawls to target Western King Prawn at night between sunset and sunrise (Figure 1). Trawls tow funnel-shaped nets along the sea floor and retain the catch in bags located at the end of the net (most commonly referred to as codends) (Figure 2). Commercial licence holders in the GSVPF are permitted to retain and sell two species of by-product harvested incidentally during prawn trawling: Balmain Bug (*Ibacus* spp.) and Southern Calamari (*Sepioteuthis australis*). A smaller penaeid (*Metapenaeopsis crassima*) is also permitted to be retained in South Australian waters but is of low commercial value.

The fishing season occurs from 1 November to 31 July of the following year with a closure in January and February. Fishing during the season generally occurs between the last quarter and first quarter of the moon and is permitted in depths greater than 10 m north of the geodesic joining GSV, Investigator Strait and Backstairs Passage. The GSVPF is divided into 121 prawn fishing blocks for research and management purposes (Figure 3). The major home ports for the GSVPF are Port Adelaide and North Haven.




Figure 1. Double-rig trawl gear and location of hopper sorting and prawn grading systems used in the Gulf St Vincent Prawn Fishery. Figure from Carrick (2003).




Figure 2. Trawl net configuration showing trawl boards, head rope, ground chain and codend with crab bag as used in the Gulf St Vincent Prawn Fishery. Figure from Carrick (2003).




Figure 3. The research management blocks for the Gulf St Vincent Prawn Fishery. South Australian Marine Park Zoning also shown.

1.2.2. Management arrangements

The *Fisheries Management Act 2007* provides the legislative framework for the management of fisheries resources in South Australia. General regulations for South Australia's prawn fisheries are described in the *Fisheries Management (General) Regulations 2007*, with specific regulations located in the *Fisheries Management (Prawn Fisheries) Regulations 2006*. These three documents provide the statutory framework for management of the GSVPF.

Management arrangements for the GSVPF have evolved since the fishery's inception in 1967 (Table 2). Between 1967 and 1979, the number of licences increased to a total of 22 before being reduced to 10 by 1990. Following licence rationalisation in 1990, the Western King Prawn stock was considered overfished and the fishery was closed during the 1991/92 and 1992/93 fishing seasons to allow recovery. The fishery reopened in 1993/94, and in 1998 the first management plan for the fishery was introduced. From 2004/05, four surveys per season were undertaken. A second management plan was implemented in 2007 (Dixon and Sloan 2007).

A review of the GSVPF in 2011 (Knuckey et al. 2011) recommended that the fleet improve its operating efficiencies. Two key management changes were subsequently implemented for the 2011/12 season: 1) the number of surveys used to manage the fishery during the fishing season was reduced to two (i.e. April and May); and 2) in March 2012, all trawl nets used for commercial fishing were modified to T90-mesh codends and fitted with Nordmøre-grids to improve catch selectivity, reduce levels of bycatch and facilitate the escapement of small prawns (Dixon et al. 2013).

Due to poor economic performance, the fishery was closed again in 2012 at the request of licence holders. Following the fishery's closure in 2012/13–2013/14, a new management framework was developed with stakeholder input and implemented in November 2014. The management framework included an individual transferrable effort (ITE) system, with transferrable fishing nights as the effort unit. From 2013/14, the number of surveys was further reduced to one undertaken in May.

From 2014/15, under the new management framework, control rules for spatial management, mean prawn size and mean nightly catches per vessel were transferred and adopted into the St Vincent Gulf Prawn Boat Owner's Association (SVPBOA) Industry Code of Practice. The Industry Code of Practice is a non-legislated document that describes the expected activities for licence holders in the GSVPF. Transferrable fishing nights could be used anytime during the season, however, the number of fishing nights that could be fished pre-Christmas (November and December) was restricted. A

revised daily logbook and nightly fishing reports were also implemented from 2014/15. The daily reports require licence holders to provide more detailed spatial information (start and end coordinates of each trawl-shot), and the nightly fishing reports provide a summary of the total catch unloaded per grade within 48 hours of unloading. Also, in November 2015, restrictions on the construction of the T90 codend were modified so that no more than 33 meshes were made up of standard mesh. Previously, between March 2012 and November 2015, no more than 10 meshes were allowed.

The latest management plan for the fishery was implemented in April 2017 and provides the overarching policy and harvest strategy for management of the fishery (PIRSA 2017). The GSVPF is currently a limited-entry (10 licence holders) fishery, which is managed by gear restrictions and a restriction on the number of fishing nights. Fishing takes place between sunset and sunrise. No trawling is permitted in waters shallower than 10 m. Current management arrangements are summarised in Table 3.

The harvest strategy listed within the management plan for the fishery uses decision rules for setting the total number of fishing nights (TACE) and number of pre-Christmas fishing nights each season (PIRSA 2017). The decision rules are based on limit, trigger and target reference points defined for three performance indicators relating to stock abundance and recruitment: 1) standardised annual catch per unit effort (CPUE); 2) standardised fishery-independent survey (FIS) CPUE; and 3) the FIS recruitment index (FRI). These indicators are also used to determine the status of the Western King Prawn stock in GSV (see Section 1.6).

Table 2. Major management milestones for the Gulf St Vincent Prawn Fishery.

Date	Management Change
1967	Commercial prawn fishing commences in GSV
1968	All SA waters closed to trawling except for specific managed zones for which permits are offered and all waters less than ten metres are closed to trawling
1969	The <i>Preservation of Prawn Resources Regulations 1969</i> is introduced and vessels licensed to fish for prawns
1975	The fishery is split into two zones when five permits are issued to specifically fish in Investigator Strait
1982	Number of Investigator Strait zone fishers reduced to two
1982	Triple rig trawl nets introduced
1986	A review of management was completed by Prof Parzival Copes
1986	A licence rationalisation strategy was implemented as an outcome of the review
1987	The <i>Fisheries (Gulf St Vincent Prawn Fishery Rationalisation) Act 1987</i> is introduced
1987	The two Investigator Strait entitlements removed and four GSVPF licences removed over the following four years and the two zones are once again amalgamated
1990	Prof Parzival Copes was requested to complete his second review of the fishery
1990	Licences reduced to 10 in GSVPF
1991	Fishery closed in June
1991	A Select Committee of the House of Assembly of South Australia reviewed the fishery's management options
1994	The fishery re-opened in February
1995	A review of the fishery was conducted by Dr Gary Morgan
1998	First management plan for the fishery was introduced (Zacharin 1997)
2000	<i>Fisheries (General) Regulations 2000</i> enabled "large" vessels to enter the fleet
2007	The second management plan was implemented (Dixon and Sloan 2007)
2011	A review of the fishery was undertaken by (Knuckey et al. 2011).
2012	The fishery was closed in November by unanimous agreement of industry. Introduction of the T90-mesh codend
2013	Morgan and Cartwright (2013) completed a review of the fishery management framework
2014	Dichmont (2014) review of the stock assessment methods, processes and outputs. The fishery reopened in November 2014. Individual transferable units were introduced. <i>A revised framework for longer-term management of the Gulf St Vincent Prawn Fishery</i> was developed
2017	The third management Plan for the South Australian Commercial Gulf St Vincent Prawn Fishery was approved by the Minister for Agriculture, Food and Fisheries (PIRSA 2017)

Table 3. Current management arrangements for the Gulf St Vincent Prawn Fishery.

Management tool	Current restriction
Permitted species	Western King Prawn (<i>Penaeus (Melicertus) latisulcatus</i>), Balmain Bug (<i>Ibacus</i> spp.), Southern Calamari (<i>Sepioteuthis australis</i>).
Limited entry	10 licences
Licence transferability	Permitted
Corporate ownership	Permitted
Spatial and temporal closures	Yes
Method of capture	Demersal otter trawl
Trawl rig	Single, double or triple
Trawling times	Not during daylight hours
Maximum combined headline length	27.43 or 29.26 m (non-amalgamated gear), 43.89 m (amalgamated gear)
Minimum codend mesh size	58 mm
Maximum vessel length	22 m
Maximum vessel power	336 kW
Catch and effort data	Daily logbook and catch disposal logbook submitted after each trip
Landing locations	Landings permitted anywhere in the State
Landing times	Landings permitted at any time during the season

1.3. Biology of the Western King Prawn

1.3.1. Distribution and habitat preferences

The Western King Prawn is a benthic crustacean that inhabits sand and seagrass habitats within estuarine and gulf waters of the Indo-west Pacific (Grey *et al.* 1983; Tanner and Deakin 2001). In South Australia, its distribution is primarily restricted to the relatively warm and reproductively favourable waters of Spencer Gulf, GSV and along the West Coast of South Australia (Ceduna, Venus Bay and Coffin Bay) (Penn 1980; Courtney and Dredge 1988).

Both juvenile and adult prawns exhibit strong diel patterns in behaviour characterised by daytime burial and nocturnal activity (Rasheed and Bull 1992; Primavera and Lebata 2000). Levels of activity are strongly linked to the lunar cycle and water temperature. Consequently, prawn catchability in fisheries is higher in the dark phases of the lunar cycle and during warmer months (Penn 1976; Penn *et al.* 1988).

Tagging experiments showed that prawn movements in GSV were predominately north to south (Kangas and Jackson 1997), with a small number of tag recaptures in eastern

Investigator Strait also indicating a general west and north-west movement in GSV (Kangas and Jackson 1997).

1.3.2. Reproductive biology

Adult prawns aggregate, mature, mate and spawn in deep water (>10 m) between October and March (King 1977). Ovary development followed by the spawning of fertile eggs occurs during a single intermoult period (30–40 days) where fertilisation likely occurs immediately prior to or on release of the eggs by the female (Penn 1980). Multiple spawning events may also occur as spawning frequency is related to moulting frequency (Penn 1980; Courtney and Dredge 1988). During the peak spawning period, the sex ratio of harvested Western King Prawns is typically female-biased, likely due to greater foraging activity of females (Penn 1976; 1980; Svane and Roberts 2005). The ideal temperature range for spawning is between 17°C (Penn 1980) and 25°C (Courtney and Dredge 1988), which generally coincides with spring and summer in South Australia (~October to March). In both gulf fisheries in South Australia, the majority of spawning occurs in November and December. Research by Roberts *et al.* (2012) indicated that reproductive success may be maximised at this time of the year under relatively high gulf temperatures that cause shorter larval durations and higher rates of larval survival (Roberts *et al.* 2012).

In all three South Australian prawn fisheries, fecundity increases exponentially with carapace length (CL), however, the increase is more pronounced in the cooler waters of GSV (Carrick 2003). Consequently, larger prawns make a greater contribution to total egg production due to their higher fecundity (Penn 1980; Courtney and Dredge 1988; Carrick 1996).

1.3.3. Early life history

Western King Prawns have an offshore adult life history phase and an inshore juvenile phase. Prawn larvae undergo metamorphosis through four main stages: nauplii, zoea, mysis and post-larvae. Key parameters that affect larval development and survival are generally considered to be temperature, salinity and food availability (Preston 1985; Jackson and Burford 2003; Bryars and Havenhand 2006; Lober and Zeng 2009). Water temperature is accepted to be one of the most important factors affecting larval growth and survival, with higher water temperatures resulting in faster rates of development and higher rates of survival (Hudinaga 1942; Roberts *et al.* 2012).

Prawn larvae are dispersed by wind-driven and tidal currents. Kangas (1999) modelled larval dispersion in GSV and demonstrated that spawning events in northern GSV were more likely to result in settlement in northern nursery areas than spawning that occurred

in southern GSV. However, rates of larval settlement from different spawning areas within the fishery were not quantified.

Kangas (1999) also indicated that post-larvae settled in inshore nursery areas at 1 mm CL, with juveniles 2–7 mm CL comprising 90% of the surveyed population. Post-larvae produced from early spawning events settle in nursery areas during December or January, when growth rates are high, before migrating to deeper water in May or June (at ~20 mm CL). Post-larvae produced from late spawning events (January to March) settle in nurseries from March, where they ‘over-winter’ before recruiting to the trawl grounds in the following summer (Kangas 1999).

1.3.4. Stock structure

Genetic analyses using ribosomal-DNA have shown significant genetic differences in the distribution of Western King Prawn haplotypes between South Australia and Western Australia (Carrick 2003). However, an analysis of the genetic structure of Western King Prawns using electrophoresis suggests a homogenous stock within South Australia (Carrick 2003).

1.3.5. Growth

As with other crustaceans, prawns undergo a series of moults to increase their size incrementally. The inability to directly age prawns has increased the reliance on tag-recapture and cohort analysis to determine growth rates. Tag-recapture studies of prawns in GSV were undertaken by King (1977) and Carrick (1982). Growth parameters were determined from 464 recaptures using a modified von Bertalanffy growth model by Kangas and Jackson (1997). Differences in growth were apparent between sexes and within years. Maximum growth occurred in April for males and in early March for females. No relationship was identified between growth and water temperature between October and December, likely due to prawns allocating more energy to reproduction during this time than growth.

1.3.6. Natural mortality

The instantaneous rate of natural mortality for GSV prawns (both sexes combined) was estimated by Kangas and Jackson (1997) as 0.003 per day. Morgan (1995) provided similar estimates of natural mortality from a range of empirical methods and models. These values are within the range of those estimated for Western King Prawns in Spencer Gulf (0.003 to 0.005; King 1977) and on the West Coast of South Australia (0.001 to 0.014; Wallner 1985) and in Western Australia (0.002 to 0.005; Penn 1976).

Sex-specific estimates of natural mortality from Xiao and McShane (2000) provided equivalent results (0.003 per day for females and 0.003 per day for males).

1.3.7. Biosecurity and prawn health

The health of Western King Prawn populations in South Australia and the potential effects of coastal pollutants, parasites and disease on growth, survival and reproduction is poorly understood. Roberts *et al.* (2010) assessed the disease status of prawns (focusing on viruses) collected from key nursery sites in both Spencer Gulf and Gulf St Vincent. A naturally occurring (endemic), and likely harmless, monodon-type baculovirus (MBV) was observed in ~60% of prawns. The MBV is a common virus known to occur throughout Australia.

White spot disease (WSD) is a highly contagious and lethal viral disease first detected in farmed prawns in south east Queensland in December 2016. The virus can cause high rates of mortality in affected prawns and is currently the subject of a national surveillance program. To date, all samples that have been collected and tested within jurisdictions outside of Queensland have returned negative results ([National Disease and Pest Outbreaks 2018](#)).

1.4. Research program

There have been numerous fisheries' research projects relevant to the Western King Prawn stock in GSV. Research has been undertaken to investigate population dynamics and biology (King 1977; Kangas 1999; Xiao and McShane 2000; Tanner and Deakin 2001; Roberts *et al.* 2012), stock structure (Carrick 2003), biosecurity and disease (Roberts *et al.* 2009), fishing gear technology (McShane 1996; Broadhurst *et al.* 1999; Dixon *et al.* 2013; Gorman and Dixon 2015), trawling impacts (Tanner 2003) and to develop fisheries models (Xiao 2004). Previous stock assessment and stock status reports detail the biological information and the history of commercial catch used to assess the status of the Western King Prawn stock in GSV (Kangas and Jackson 1997; Xiao and McShane 1998; Boxshall *et al.* 1999; Boxshall and Williams 2000; Boxshall and Johnson 2001; Svane 2003; Svane and Johnson 2003; Roberts *et al.* 2007a, 2007b; 2008; 2009; Hooper *et al.* 2009; Dixon *et al.* 2011; 2012; Beckmann *et al.* 2015; 2016; McLeay *et al.* 2017).

From 2011 to 2014, three separate independent reviews of the stock assessment and harvest strategy for the GSVPF were conducted (Knuckey *et al.* 2011; Morgan and Cartwright 2013; Dichmont 2014). As a result of these reviews, there has been a rationalisation of the research program. The principal change was a reduction from four

fishery-independent surveys (FIS') per year to one (conducted in May). The focus of the 2017/18 research program is to assess the status of the Western King Prawn stock in GSV using fishery-independent and -dependent information collected from the GSVPF during the 2017/18 fishing season.

1.5. Information sources used for assessment

Fishery-independent data have been collected for the GSVPF using surveys coordinated by SARDI since 1984. SARDI also maintains a comprehensive fishery-dependent catch and effort database for the GSVPF using data obtained from the South Australian Fishing Industry Council (SAFIC) annual reports between 1968 and 1987. Data from July 1987 were obtained from daily commercial logbooks provided to SARDI by licence holders. No fishing was conducted from July 1991 to February 1994.

1.6. Stock status classification

A National Fishery Status Reporting Framework (NFSRF) was developed to enable the consistent assessment of the status of Australian fish stocks (Flood *et al.* 2012, 2014; Stewardson *et al.* 2016). The NFSRF of Flood *et al.* (2014), which the management plan for the GSVPF refers to, considers whether the current level of fishing pressure is adequately controlled to ensure that stock abundance is not reduced to a point where recruitment is impaired. The system combines information on both the current stock size and level of catch into a single classification for each stock against defined biological reference points. The stock is then classified as 'sustainable', 'transitional-recovering', 'transitional-depleting', 'overfished', 'environmentally limited', or 'undefined' (Table 4).

Table 4. Stock status terminology (Flood et al. 2014).

	Stock status	Description	Potential implications for management of the stock
	Sustainable	Stock for which biomass (or biomass proxy) is at a level sufficient to ensure that, on average, future levels of recruitment are adequate (i.e. not recruitment overfished) and for which fishing pressure is adequately controlled to avoid the stock becoming recruitment overfished	Appropriate management is in place
	Transitional-recovering	Recovering stock—biomass is recruitment overfished, but management measures are in place to promote stock recovery, and recovery is occurring	Appropriate management is in place, and the stock biomass is recovering
	Transitional-depleting	Deteriorating stock—biomass is not yet recruitment overfished, but fishing pressure is too high and moving the stock in the direction of becoming recruitment overfished	Management is needed to reduce fishing pressure and ensure that the biomass does not deplete to an overfished state
	Overfished	Spawning stock biomass has been reduced through catch, so that average recruitment levels are significantly reduced (i.e. recruitment overfished). Current management is not adequate to recover the stock, or adequate management measures have been put in place but have not yet resulted in measurable improvements	Management is needed to recover this stock; if adequate management measures are already in place, more time may be required for them to take effect
	Environmentally limited	Spawning stock biomass has been reduced to the point where average recruitment levels are significantly reduced, primarily as a result of substantial environmental changes/impacts, or disease outbreaks (i.e. the stock is not recruitment overfished). Fisheries management has responded appropriately to the environmental change in productivity	Appropriate management is in place
	Undefined	Not enough information exists to determine stock status	Data required to assess stock status are needed

PIRSA has adopted this classification system to determine the status of all South Australian fish stocks. Status is categorised for the Western King Prawn stock in GSV from a combination of three performance indicators as defined in the harvest strategy for the fishery:

1. Standardised annual commercial CPUE ($\text{kg.block}^{-1}.\text{vessel-night}^{-1}$) that is estimated from daily commercial logbook data provided to SARDI.
2. Standardised FIS CPUE ($\text{kg.trawl-shot}^{-1}$) that is estimated from data collected in the May FIS.
3. FRI (recruits.h^{-1}) that is estimated from data collected in the May FIS.

To categorise stock status, estimates for each of the three performance indicators are assessed in relation to how they align against limit, trigger and target reference points for each performance indicator as defined in the harvest strategy for the fishery (Table 5) (PIRSA 2017).

Table 5. Decision rules for classifying the status of Western King Prawn stock in GSV (PIRSA 2017). *Transitional may refer to transitional depleting, or transitional recovering. Transitional depleting is based on the definition in Flood *et al.* (2014) as “biomass is not yet recruitment overfished, but fishing pressure is too high and moving the stock in the direction of becoming recruitment overfished”. Transitional Recovering is based on the definition in Flood *et al.* (2014) as “biomass is recruitment overfished but management measures are in place to promote stock recovery and recovery is occurring”.

		Performance Indicator – Standardised Annual Commercial CPUE (kg.block ⁻¹ .vessel-night ⁻¹)			
		Limit <600	Trigger ≥600 - <750	Target ≥750 - <900	High ≥900
Performance Indicator - Standardised FIS CPUE (kg.trawl-shot ⁻¹)	Limit <20	Overfished	Transitional* OR Overfished when FRI <450 recruits.h ⁻¹	Transitional* OR Sustainable when FRI ≥450 recruits.h ⁻¹	
	Trigger ≥20-<25	Transitional* OR Overfished when FRI <450 recruits.h ⁻¹	Transitional* OR Sustainable when FRI ≥450 recruits.h ⁻¹		
	Target ≥25-<30	Transitional* OR Sustainable when FRI ≥450 recruits.h ⁻¹		Sustainable	
	High ≥30				

2. METHODS

2.1. Fishery-independent surveys

Surveys using commercial vessels with fishery-independent observers on-board, were conducted in GSV prior to the December, March, April and May harvest periods from 2004/05 to 2010/11, in April and May in 2011/12, and in May from 2013/14. This report uses data from May surveys only to enable temporal comparison of the FIS data collected since 2004/05.

Surveys were typically done using three to six vessels over two or three consecutive nights commencing on the second and third nights following the last quarter of the moon. There were two exceptions; 1) 2005 to 2008, when surveys were conducted during one night on the dark of the moon with approximately ten vessels and, 2) the 2014 survey, which was extended over the dark phase of the moon with only one vessel participating. From 2005 to 2016, the number of sites sampled ranged from 101–112, except for 2014 where a ‘reduced’ survey of 48 sites was completed. The survey in May 2018 includes data from 101 of the 109 survey locations specified in the current Management Plan (PIRSA 2017) (Figure 4).

Survey shots were done at semi-fixed sites in the ten regions of the GSVPF (Figure 4). Each shot began close to a known location (recorded by Global Positioning System, GPS) and then continued in a set direction for a specified period of time (usually 30 minutes). The total distance covered was dependent on trawl speed, which was influenced by vessel power, tide and weather conditions. Data collected for each shot location include the total catch weight, catch weight of each prawn size grade, number of nets used, trawl duration, tide direction, and the number of prawns in a 7 kg bucket (‘bucket count’ - a rapid measure used to estimate prawn size).

T90-mesh (conventional diamond net turned 90 degrees) codends with bycatch reduction grids were introduced into commercial fishing operations of the GSVPF in 2012 to reduce discards. However, the reference levels relating to standardised FIS CPUE ($\text{kg.trawl-shot}^{-1}$) and FRI (recruits. h^{-1}) described within the current harvest strategy were derived from historical data collected from diamond-mesh codends only since 2005. Consequently, only data collected from diamond-mesh codends since 2005 were used to estimate standardised FIS CPUE and FRI in the 2017/18 season because these data directly inform stock status determination. Data from T90 mesh codends are still collected during the FIS to enable potential calibration with data collected from diamond-mesh codends in the future.

Estimates of mean prawn size (N prawns. kg^{-1}) are calculated from the 7 kg subsample, with higher counts representing smaller prawns and lower counts representing larger prawns. To present the spatial distribution of catch rate and prawn size, prawns were defined as ‘small’ (>33 prawns. kg^{-1}), ‘medium’ (30-33 prawns. kg^{-1}), ‘large’ (28-30 prawns. kg^{-1}) and ‘extra-large’ (<28 prawns. kg^{-1}), and CPUE was defined as ‘very low’ (<50 kg.trawl-shot $^{-1}$), ‘low’ (50-100 kg.trawl-shot $^{-1}$), ‘moderate’ (100-150 kg.trawl- shot $^{-1}$) and ‘high’ (>150 kg.trawl-shot $^{-1}$). The GPS locations of each trawl station were used to indicate prawn size and CPUE.

A random sample of 100 prawns is also taken from each shot to obtain information on sex ratio and length-frequency. The catch rate of recruits per trawl-hour was determined for each survey. ‘Recruits’ are defined by prawn size: ≤ 32 mm CL for males and ≤ 34 mm CL for females.




Figure 4. The 109 survey stations specified in the 2017 Management Plan and regions of the Gulf St Vincent Prawn Fishery. The northern gulf (pink) includes Region 1 (RG1) and Region 6 (RG6), the central gulf (green) includes Region 2 (RG2), Region 5 (RG5) and Little Hole (LIH), and the southern gulf (blue) includes Deep Hole (DPH), Southern Deep Hole (SDH) and Investigator Strait (INV).

2.2. Fishery statistics

2.2.1. Catch and effort

Fishery catch (t) and effort (vessel-nights or trawl hours) data are presented for available data since the 1968/69 fishing season. In this report, a ‘fishing season’ is defined as the period from 1 November to 31 July of the following year. Estimated prawn catch for each shot was adjusted using validated post-harvest catches reported in monthly logbooks. Catch and effort data includes fishery-dependent ‘searching’ and spot survey catches, as well as FIS catches that are retained for sale by the SVPBGA.

The main spawning period for Western King Prawns in GSV extends from November to March, so catch and effort data are also presented for early (November–December), late (January–March) and non-spawning (April–October) periods. The spatial distribution of the annual harvest per fishing block is presented from 2006/07–2017/18, however, confidential data (<5 licence holders) are not shown.

All data are entered and maintained by SARDI Aquatic Sciences. Data are checked for errors during data entry, and further validated via electronic lookup tables and code-driven software that is activated during data entry and reporting operations. SARDI staff contact licence holders to correct any errors or missing information identified in the quality assurance process.

2.2.2. Prawn size

Mandatory reporting of commercial prawn-grade data in daily logbooks was introduced in 2005/06. The grade is determined from the number of prawns to the pound (e.g. ‘U10’ = under 10 prawns per pound). To facilitate interpretation of the prawn-grade data, grades were assigned to four size categories (Table 6). Soft and broken prawns that were not graded, were assigned to a fifth category ('S&B'). Data presented are from commercial fishing nights only and reported as the proportion of the total catch occurring in each of the size classes (see Dixon *et al.* 2012).

Uncertainty associated with the calculation of prawn size arises from: 1) data not being available from the entire fleet for all seasons; 2) differences in the grading among the vessels; and 3) uncertainty associated with the unvalidated grade data provided in logbooks (Dixon *et al.* 2012).

Table 6. Categories assigned to reported prawn grades from commercial logbook data for the Gulf St Vincent Prawn Fishery.

Prawn grade	Categories in logbook
Extra Large	U6 U8, XL U10, L
Large	9/12 U12 LM 10/15 13/15 10/20 (50%), 12/18 (50%)
Medium	10/20 (50%), 12/18 (50%) 16/20, M
Small	SM, 19/25 21/25 S, 20+, 21/30 26+ 30+, 31/40
Soft & Broken	S/B, B&D, MIX, REJ, SMS, blank, ERR

2.3. Catch rate standardisation

CPUE is commonly used as an indicator of relative biomass (stock abundance) in crustacean fisheries worldwide. However, to improve the relationship between CPUE and relative biomass, it is important to standardise CPUE to account for the influence of variables that are not related to population size. Generalised linear models (GLMs) were used to standardise FIS CPUE ($\text{kg.trawl-shot}^{-1}$) since 2005, and annual commercial CPUE ($\text{kg.block}^{-1}.\text{vessel-night}^{-1}$) since 1990/91, as per the methods in Noell *et al.* (2015). Candidate variables for standardising FIS CPUE were fishing year-survey, region and vessel, while those for standardising annual commercial CPUE were fishing year, month, region, lunar phase, effort, licence number and cloud cover. Standardised estimates of CPUE are presented with 95% confidence intervals.

3. RESULTS

3.1. Fishery-independent surveys

3.1.1. CPUE

Nominal FIS CPUE (diamond mesh) increased from $22.8 \pm 1.6 \text{ kg.trawl-shot}^{-1}$ in 2004/05 to $42.3 \pm 3.5 \text{ kg.trawl-shot}^{-1}$ in 2007/08, but then declined to a near historical low of $22.3 \pm 2.1 \text{ kg.trawl-shot}^{-1}$ in 2011/12 before the fishery was closed in 2012/13 and 2013/14 (Figure 5a). Nominal FIS CPUE from a ‘reduced’ survey in 2013/14 was $56.5 \pm 5.0 \text{ kg.trawl-shot}^{-1}$ but declined to $31.5 \pm 2.1 \text{ kg.trawl-shot}^{-1}$ in 2014/15. In the 2015/16 fishing season, nominal FIS CPUE increased 45.2% to $45.7 \pm 3.1 \text{ kg.trawl-shot}^{-1}$, but in 2016/17 declined to $33.1 \pm 2.4 \text{ kg.trawl-shot}^{-1}$. In 2017/18, the nominal FIS CPUE estimate of $36.2 \pm 2.7 \text{ kg.trawl-shot}^{-1}$ was similar to that recorded in 2016/17. Nominal seasonal estimates of FIS CPUE measured from the T90-mesh since 2011/12 were 12.3–46.6% lower than those estimated from diamond mesh but followed a similar trend (Figure 5a).

Fishing year-survey, region and vessel were all highly significant variables in the GLM used to standardise nominal estimates of FIS CPUE. However, a low overall model goodness-of-fit (adjusted R^2 value 0.13) indicates other sources of variability are unaccounted for. A total of 14.3% of the deviance in survey catches was explained by the model (fishing year-survey being the most important at 7.4%), which indicates that 85.7% of the deviance was caused by unknown factors (Table 7).

Trends in standardised FIS CPUE and nominal FIS CPUE measured from diamond mesh between 2004/05 and 2017/18 were similar, differing by less than 18.9% in each year surveyed over this period (Figure 5b). Estimates of standardised FIS CPUE have remained in the high range for this performance indicator ($\geq 30 \text{ kg.trawl-shot}^{-1}$) in surveys undertaken since 2013/14, ranging from $35.6 \text{ kg.trawl-shot}^{-1}$, 95% CI [23.4, 51.9] in 2014/15 to $54.3 \text{ kg.trawl-shot}^{-1}$, 95% CI [37.6, 75.6] in 2015/16 (Figure 5b). In 2017/18, standardised FIS CPUE was estimated from a total of 101 trawl shots and was $38.6 \text{ kg.trawl-shot}^{-1}$, 95% CI [28.2, 51.5]. This estimate was slightly higher (by 1.1%) than the 2016/17 estimate of $38.2 \text{ kg.trawl-shot}^{-1}$, 95% CI [25.2, 55.4].




Figure 5. Key outputs from May fishery-independent surveys: (a) nominal catch per unit effort (CPUE, kg.trawl-shot⁻¹) from diamond v T90-mesh; (b) standardised versus nominal catch per unit effort (CPUE, kg.trawl shot⁻¹) (diamond net); (c) the fishery recruitment index (FRI) (recruits.h⁻¹), and (d) prawn size (prawns.kg⁻¹). Indices were calculated from 101 survey shots in the 2017 Management Plan. Note: T90 and diamond gear were sampled side by side from 2011/12 and a reduced survey was conducted in 2013/14. Error bars = standard error for all metrics except standardised CPUE where error bars are upper and lower (95%) confidence intervals. Grey area indicates the period when no survey occurred during the fishery closure between 2012/13 & 2013/14.

Table 7. Analysis of deviance (Type II test) for the GLM used to standardise survey CPUE for the GSVPF. Abbreviations: SS, sum of squares; df, degrees of freedom; F , F -statistic.

Effect	SS	df	F
Fishing year-survey	242.1	33.0	9.2***
Region	154.6	7.0	27.7***
Vessel	73.3	17	5.4***
Residuals	2817.8	3531	NA

Significance: *** $p < 0.001$

3.1.2. Recruitment

In 2013/14, the FRI estimate recorded in the ‘reduced’ survey was the highest on record at $1,069 \pm 158$ recruits.h $^{-1}$ (Figure 5c). Following re-opening of the fishery in November 2014, the FRI estimate in 2014/15 was 327.6 ± 62.2 recruits.h $^{-1}$, which was the lowest estimate since 2004/05 (239.2 ± 31.5 recruits.h $^{-1}$). In 2015/16, the FRI was 837.1 ± 115.6 recruits.h $^{-1}$, more than double the FRI estimated in 2014/15 and the second highest value on record.

In 2017/18, the FRI was estimated from a total of 101 completed trawl shots and was 670.3 ± 82.8 recruits.h $^{-1}$ (Figure 5c). This estimate is 14.6% lower than the FRI estimate recorded in 2016/17 (784.4 ± 116.3 recruits.h $^{-1}$).

Similar to the patterns in CPUE observed between diamond and T90-mesh since 2011/12 (Section 3.1.1), estimates of FRI from T90-mesh were lower (24.1–62.9%) than those estimated from diamond mesh but followed the same trend (Figure 5c).

3.1.3. Size

Mean prawn size, measured from the 7-kg subsample taken per shot, has an inverse relationship with the number of prawns per kg (i.e. larger mean prawn size = less prawns per kg). Mean prawn size has fluctuated since 2004/05 (Figure 5d). Prior to the fishery closure, mean prawn size in 2011/12 was one of the smallest estimates on record at 37.1 ± 1.2 prawns.kg $^{-1}$. Following the fishery closure, mean prawn size in 2013/14 increased 11.2% to 33.0 ± 1.1 prawns.kg $^{-1}$, and in 2014/15 mean prawn size was the largest on record at 29.3 ± 0.8 prawns.kg $^{-1}$ (Figure 5d). Since 2014/15, mean prawn size has steadily decreased, and in 2017/18 was the smallest on record at 37.2 ± 1.2 prawns.kg $^{-1}$ (Figure 5d).

Trends in mean prawn size estimated from the T90-mesh since May 2012 are similar to those estimated from diamond mesh. However, annual estimates of mean prawn size from T90-mesh were slightly larger (0.1–7.3%) than diamond mesh (Figure 5d).

3.1.4. Spatial patterns in CPUE and size

Between 2011/12 and 2017/18, there was a general north to south gradient in the size of prawns, with a relatively higher CPUE of small and medium prawns (≥ 30 prawns. kg^{-1}) in northern GSV and extra-large prawns (< 28 prawns. kg^{-1}) in southern GSV.

In 2017/18, high CPUE (> 100 kg.trawl.shot $^{-1}$) of extra-large prawns occurred in the southern part of the gulf (DPH) while moderate CPUE (50–100 kg.trawl.shot $^{-1}$) of small and medium prawns occurred in the northern and central parts of the gulf (RG1, RG2) (Figure 6).





Figure 6. Catch rate and mean size of prawns sampled from the diamond-mesh codend during May FIS' in the GSVPF between 2011/12 and 2017/18. A reduced survey was undertaken in 2013/14 following the fishery closure between 2012/13 & 2013/14. Map on bottom right shows the regions used to describe the spatial distribution of fishing.



3.2. Fishery statistics

3.2.1. Catch and effort

The total commercial catch of Western King Prawn in the GSVPF in 2017/18 was 235.3 t, not including 3.5 t taken in the May 2018 survey (Figure 7a, Table 8). The 2017/18 catch was greater than taken in 2015/16 (217.8 t), 2016/17 (224.6 t) and the average since 2000/01 (228.7 ± 15.8 t; Figure 7a, Table 8).

In 2017/18, 58.0 t of the catch was taken during the early spawning period prior to Christmas (November–December), which was 24.7% of the total annual catch, and 26.6% greater than that taken in the same period in 2016/17 (45.8 t). The amount of catch taken during the late spawning period (January–March) in 2017/18 (25.2 t) was 15.4% less than that taken in 2016/17 (29.8 t) (Table 8). Similar to 2015/16 and 2016/17, the highest proportion of catch (152.1 t, 64.7%) taken in 2017/18 was harvested during the non-spawning period (April–July).

Commercial fishing was conducted over 291 vessel-nights in 2017/18, comprising 97% of the TACE of 300 vessel-nights (Figure 7b). All 50 allocated pre-Christmas vessel-nights were fished. Total effort was within the range recorded since the fishery reopened in 2014/15 (287–296 vessel-nights), but low compared to the amount of nights fished prior to 2003/04 (Figure 7b). The total commercial effort recorded in 2017/18 was 2,858 trawl-hours, which was similar to that recorded in 2015/16 (2,799 trawl-hours) and 2016/17 (2,836 trawl-hours), and among the lowest levels of effort recorded in the history of the fishery (Figure 7c).




Figure 7. Fishery-dependent catch and effort data outputs for the Gulf St Vincent Prawn Fishery; (a) Annual catch (t) and (b) effort (vessel-nights) separated by early spawning (November–December), late spawning (January–March) and non-spawning (April–October) and (c) annual commercial effort (hours * 1000). Catch data from surveys not included.

Table 8. Monthly distribution of commercial (and survey) catch in the Gulf St Vincent Prawn Fishery from 1989/99–2017/18.

Season	Nov	Dec	Feb	Mar	Apr	May	Jun	Jul	Total
1989/99	35.8	9.4	-	47.8	33.6	42.3	-	-	168.8
1990/91	-	5.3	-	48.0	36.0	31.9	13.1	-	134.3
1991/92	-	-	-	-	-	-	-	-	-
1992/93	-	-	-	-	-	-	-	-	-
1993/94	-	-	-	97.9	69.3	41.2	17.0	-	225.4
1994/95	-	-	26.0	60.9	52.4	8.0	-	-	147.3
1995/96	30.7	48.3	-	65.5	46.4	67.1	-	-	258.0
1996/97	37.9	25.7	-	34.8	41.9	45.0	24.7	-	209.9
1997/98	43.8	31.7	15.8	81.3	53.7	40.5	-	-	266.9
1998/99	69.8	30.5	-	65.6	99.7	48.1	22.5	-	336.2
1999/00	19.3	82.6	27.3	71.0	76.3	91.7	32.0	-	400.2
2000/01	65.9	60.6	-	84.3	86.9	72.4	14.8	-	384.9
2001/02	8.8	94.0	-	58.5	80.6	62.1	18.1	-	322.1
2002/03	4.0	60.1	11.5	72.8	46.6	37.0	-	-	231.9
2003/04	-	13.9	-	28.9	69.5	57.7	2.5	-	172.5
2004/05	-	43.5 (2.0)	-	50.1 (2.4)	40.9 (2.0)	46.7 (2.4)	27.2	-	208.4 (8.8)
2005/06	-	21.2 (1.9)	-	13.6 (2.5)	64.1 (3.6)	40.4 (2.5)	35.5	-	174.9 (10.5)
2006/07	-	26.5 (1.6)	-	51.5 (2.9)	86.1 (3.7)	45.3 (4.5)	-	-	209.4 (12.7)
2007/08	-	30.2 (2.5)	-	63.5 (3.3)	69.6 (4.2)	65.8 (4.5)	-	-	229.0 (14.4)
2008/09	36.5	(3.2)	-	56.3 (4.0)	53.1 (4.1)	126.7 (4.1)	-	-	272.6 (15.4)
2009/10	-	31.4 (2.3)	-	24.7 (2.6)	109.2 (4.3)	58.3 (3.9)	-	-	223.6 (13.0)
2010/11	-	31.9 (2.5)	-	6.9 (2.9)	43.3 (3.0)	68.5 (2.6)	27.6	-	178.3 (11.0)
2011/12	16.0*	-	25.0	38.0 (2.0)	37.2 (2.1)	8.8*	-	-	125.0 (4.2)
2012/13	-	-	-	-	-	-	-	-	-
2013/14	-	-	-	-	-	(1.9)	-	-	(1.9)
2014/15	40.7	16.4	-	65.9	56.8	44.5 (2.8)	25.2*	-	249.4 (2.8)
2015/16	20.1*	-	54.4	49.8	32.4 (4.1)	42.7	18.4	-	217.8 (4.1)
2016/17	21.2	24.6	-	29.8	40.2	49.3 (2.9)	59.4*	-	224.6 (2.9)
2017/18	58.0*	-	25.2	60.0	49.2 (3.5)	42.9*	-	-	235.3 (3.5)

*Data amalgamated across more than one month due to confidentiality requirements (<5 licence holders), data from gear trials not included. Survey catches since 2011/12 comprise catches from both T90 and diamond net.

3.2.2. CPUE

The GLM used to standardise nominal commercial CPUE data had a relatively high goodness-of-fit (adjusted R^2 value = 0.78). Region, fishing year, month, lunar phase, effort, licence number and cloud cover were all significant variables in final model (Table 9). Effort was the most influential variable, explaining 73.2% of the total model deviance. All other variables explained a total of 5.0% of the total model deviance and 21.7% of the deviance was caused by unknown factors.

Table 9. Analysis of deviance (Type II test) for the GLM used to standardise annual commercial catch in the Gulf St Vincent Prawn Fishery. Abbreviations: SS, sum of squares; df, degrees of freedom; F, F-statistic.

Effect	SS	df	F
fishing year	4617.3	23	131.6***
month	973.8	7	91.2 ***
region	410.9	9	29.9***
lunar phase	75.3	1	49.4***
effort	92100.0	1	60382.5***
licence no	188.7	9	13.7***
cloud	53.2	1	34.9***
residuals	27334.5	17921	NA

significance: *** $p < 0.001$.

Estimates of standardised annual commercial CPUE since 1990/91 are higher than estimates of nominal CPUE and are likely attributed to effort being used as a fixed term in the GLM (Figure 8). Despite the differences in scale of annual nominal and standardised CPUE estimates, the trend in both CPUE metrics was similar over time. It should be noted that estimates of standardised annual commercial CPUE will differ in each year that data are reported as new data from the latest season (e.g. 2017/18) are added to the GLM.

Historically, estimates of standardised annual commercial CPUE were relatively stable from 1993/94 to 2004/05, ranging from 644 kg.block⁻¹.vessel-night⁻¹, 95% CI [628, 660] in 2002/03 to 881 kg.block⁻¹.vessel-night⁻¹, 95% CI [859, 903] in 1999/00 (Figure 8). From 2005/06, standardised commercial CPUE increased steadily, reaching a peak of 1,146 kg.block⁻¹.vessel-night⁻¹, 95% CI [1105, 1187] in 2008/09. From 2008/09, standardised commercial CPUE declined to a low of 588 kg.block⁻¹.vessel-night⁻¹, 95% CI [563, 613] in 2011/12. The fishery was then closed in 2012/13 and 2013/14.

Following reopening of the fishery in 2014/15, the estimate of standardised annual commercial CPUE was $890 \text{ kg.block}^{-1}.\text{vessel-night}^{-1}$, 95% CI [864, 917], which was within the current target range for this performance indicator of ≥ 750 and $< 900 \text{ kg.block}^{-1}.\text{vessel-night}^{-1}$. In 2015/16, standardised annual commercial CPUE increased further to $929 \text{ kg.vessel-night}^{-1}$, 95% CI [902, 957], which was in the high range defined for this performance indicator ($\geq 900 \text{ kg.block}^{-1}.\text{vessel-night}^{-1}$).

In 2017/18, standardised annual commercial CPUE was $895 \text{ kg.block}^{-1}.\text{vessel-night}^{-1}$, 95% CI [868, 922], an increase of 0.7% since 2016/17 ($889 \text{ kg.block}^{-1}.\text{vessel-night}^{-1}$, 95% CI [863, 916]) and within the target range defined for this performance indicator (Figure 8).




Figure 8. Comparison of standardised annual commercial CPUE and nominal annual commercial CPUE ($\text{kg.block}^{-1}.\text{vessel-night}^{-1}$) in the Gulf St Vincent Prawn Fishery from 1990/91–2017/2018. Error bars are \pm standard error for nominal CPUE, and upper and lower (95%) confidence intervals for standardised CPUE.

3.2.3. Spatial patterns in catch and effort

Following the use of FIS' in 2004/05 to spatially control the areas harvested in GSV, the number of prawn fishing blocks that were fished declined until 2009/10 (Figure 9). From

2014/15, the number of surveys was reduced, control rules for spatial management were transferred and adopted into the SVPB Industry Code of Practice (see Section 1.2.2), and the number of blocks that were fished subsequently increased. In 2017/18, fishing took place in 11 blocks in November–December and 43 blocks in the remaining months of the season. The number of blocks fished in 2017/18 was lower than 2016/17, yet remain at a relatively high level compared to that recorded between 2004/05 and 2009/10.




Figure 9. The number of blocks fished pre-Christmas (November and December) and all other months in the Gulf St Vincent Prawn Fishery from 1989/90–2017/18.

The spatial distribution of the GSVPF catch has varied annually. However, since 2014/15, a relatively high proportion of the catch has been taken in fishing blocks located in southern parts of GSV (Figure 10). In 2017/18, 65.2% (153.4 t) of the total annual catch was taken from southern GSV in the regions DPH, INV, RG3, RG4 and SDH. Relatively high catches were also recorded in northern parts of the gulf, with 52.7 t recorded from RG1, representing 22.4% of the total annual catch of the GSVPF (Figure 10).

Similar to previous seasons, relatively low catches were recorded from RG4, RG5 and RG6 in the eastern parts of GSV in 2017/18. A combined total of 14.3 t was taken from these regions, representing 6.1% of the total annual catch recorded in the fishery.




Figure 10. The annual catch per fishing block in the Gulf St Vincent Prawn Fishery from 2007/08–2017/18. Note, the fishery was closed in 2012/13 and 2013/14. Confidential data from <5 licence holders are depicted in grey and typically <5 t. Map on bottom left shows regions used to describe spatial distribution of fishing (see Figure 4 for more detail).

3.2.4. Prawn size

The annual catch of Western King Prawn in the GSVPF since 2005/06 comprised, on average, 27% extra-large prawns, 39% large prawns, 22% medium prawns, 6% small prawns, and 7% soft & broken prawns (Figure 11). The percentage of small and medium prawns in the catch increased from 2005/06 to 2010/11, and a corresponding decrease in the proportion of large and extra-large prawns in the catch was recorded over the same period.

Following the re-opening of the fishery in 2014/15, extra-large and large prawns comprised a total of 69% of the total catch (Figure 11, Table 10). The proportion of extra-large and large prawns comprising the catch has remained similar in each season since 2014/15, and in 2017/18, extra-large and large prawns comprised a total of 66.0% of the total catch.

The percentage of small prawns in the catch has also remained similar in the last four seasons, and in 2017/18, small prawns comprised 5.3% of the catch (Figure 11, Table 10). The proportion of soft and broken prawns has historically been relatively low and in 2017/18 comprised 5.7% of the total annual catch (Figure 11, Table 10).




Figure 11. Size-grade composition (%) of annual harvests in the Gulf St Vincent Prawn Fishery from 2005/06–2017/18. Grades include; soft and broken (S&B, not graded), small (S, 20+), medium (M, 16/20), large (L, 10/15) and extra-large (XL, U10).

Table 10. Size-grade composition (t) of annual harvests in the Gulf St Vincent Prawn Fishery from 2005/06–2017/18. Grades include; soft and broken (S&B, not graded), small (S, 20+), medium (M, 16/20), large (L, 10/15) and extra-large (XL, U10). Note: grade weight totals were estimated from daily fishing logbooks.

SEASON	XL	XL-Split			L	M	S	SB
		U10	U8	U6				
2005/06	29.0				30.1	8.7	3.4	5.9
2006/07	46.0				53.1	26.0	6.8	9.7
2007/08	40.2				56.1	25.6	4.2	10.1
2008/09	50.4				67.6	38.4	9.7	10.4
2009/10	36.2				59.3	49.9	13.6	12.6
2010/11	26.2				42.9	43.9	12.0	7.6
2011/12	25.6				35.4	23.7	8.8	5.6
2012/13								
2013/14								
2014/15	57.1	36.8	17.8	2.5	104.8	49.1	9.1	15.9
2015/16	57.3	31.6	22.1	3.6	78.7	41.3	11.9	16.0
2016/17	53.4	32.2	18.2	3.0	88.9	40.9	9.9	14.1
2017/18	52.1	30.5	19.2	2.4	88.7	49.1	11.4	12.1

4. DISCUSSION

4.1. Information sources used for assessment

The current management plan for the fishery provides the decision rules for classifying the status of the Western King Prawn stock in GSV (PIRSA 2017). It uses limit, trigger and target reference points defined for three performance indicators relating to stock abundance and recruitment (PIRSA 2017). The performance indicators relating to stock abundance are: 1) standardised annual commercial CPUE estimated from fishery-dependent data recorded in daily commercial logbooks; and 2) standardised FIS CPUE estimated from data collected in the May FIS. The FIS recruitment index (FRI) is also estimated from data collected in the May FIS.

4.1.1. *Fishery-independent data*

From December 2004, a comprehensive survey program has contributed towards assessment of the status of the Western King Prawn stock in GSV. The methods used to collect the data have remained relatively consistent and are considered high quality, despite adaptations to the survey program in recent years (McLeay *et al.* 2017). The locations sampled in the May FIS have remained consistent over time, and standardised FIS CPUE and the FRI are considered the best available indices of current and future relative biomass. Also, while the distribution, extent and timing of recruitment is highly variable in GSV, new recruits are expected to be adequately represented in the catch during the FIS in May.

The introduction of T90-mesh codends fitted with Nordmøre-grids into the GSVPF in 2012 was based on the results of FRDC Project 2009/069 (Dixon *et al.* 2013), which indicated that this trawl-gear modification improved catch selectivity, reduced levels of bycatch and facilitated the escapement of small prawns in the fishery. Under the current management plan, data collected from diamond-mesh codends in the May FIS are used to estimate standardised FIS CPUE and the FRI. Ongoing data collection from T90-mesh codends during the FIS will potentially enable calibration of data collected from diamond-mesh codends in the future.

4.1.2. *Fishery-dependent data*

The trends in standardised annual commercial CPUE estimated from data recorded in daily commercial logbooks resemble the trends in standardised FIS CPUE since 2004/05. However, there remain sources of uncertainty associated with data collected from the fishery: 1) the spatial distribution of fishing effort has changed over time due to changes in spatial management of the fishery; 2) estimates of standardised annual commercial CPUE are based on daily logbook data from 1991/92 because catch data

available prior to this time are either not available or considered unreliable; and consequently, (3) estimates of standardised CPUE from 1991/92 do not take into account the period when catches were considerably higher (1975/76–1983/84).

4.2. Status

Since its inception in 1967, the GSVPF has gone through a number of cycles characterised by steady or increasing catches and catch rates, subsequent declines in recruitment and fishery performance, and resulting closure periods (1991/92, 1992/93, 2012/13 and 2013/14). The current harvest strategy for the fishery (PIRSA 2017) was developed considering the history of fishery performance, previous stock assessments and reviews of the fishery conducted by Knuckey *et al.* (2011), Morgan and Cartwright (2013), and Dichmont (2014).

In 2017/18, the total commercial catch of Western King Prawn in the GSVPF was 235.3 t, with an additional 3.5 t taken in the May 2018 survey. This catch was similar to the average catch landed by the fishery between 2000/01 and 2017/18 (228.7 ± 15.8 t). Fishing was conducted over 291 vessel-nights, comprising 97% of the TACE of 300 vessel-nights. All 50 allocated pre-Christmas vessel-nights were fished.

Estimates of standardised annual commercial CPUE were in the high range (≥ 900 kg.block $^{-1}$.vessel-night $^{-1}$) in 2015/16, and within the target range (≥ 750 and < 900 kg.block $^{-1}$.vessel-night $^{-1}$) in 2016/17 and 2017/18 (PIRSA 2017). Estimates of standardised FIS CPUE have remained in the high range (≥ 30 kg.trawl-shot $^{-1}$) since 2013/14. Similarly, the FRI has remained in the high range (≥ 600 recruits.h $^{-1}$) for four of the last five annual surveys (2013/14, 2015/16, 2016/17 and 2017/18).

PIRSA has adopted the NFSRF stock status classification system to determine the status of all South Australian fish stocks (Flood *et al.* 2014). The NFSRF was used to assess the status of the Western King Prawn stock in GSV as ‘transitional-depleting’ in 2014/15 and ‘sustainable’ in 2015/16 using a ‘weight-of-evidence’ approach. The current harvest strategy provides a prescriptive approach to determining the status of the Western King Prawn stock in GSV (PIRSA 2017). Stock status is determined within the harvest strategy by assessing how each of the three performance indicators align against pre-defined limit, trigger and target reference points (Table 5) (PIRSA 2017).

In 2017/18:

- Standardised annual commercial CPUE was 895 kg.block $^{-1}$.vessel-night $^{-1}$.
- Standardised FIS CPUE was 38.6 kg.trawl-shot $^{-1}$; and
- The FRI was 670.3 recruits.h $^{-1}$.

When the 2017/18 estimates for each performance indicator are applied in the decision matrix used to determine the status of the Western King Prawn stock in GSV (Table 5), the stock is classified as '**sustainable**'. This is the same stock status classification as determined in 2016/17 (McLeay *et al.* 2017).

4.3. Future directions

The 'sustainable' classification of the Western King Prawn stock in GSV is a positive sign for the fishery. However, the history of serial declines in fishery performance necessitate that sources of uncertainty in the assessment are highlighted and addressed in the future.

The status of the Western King Prawn stock in GSV in 2017/18 has been determined under the newly adopted harvest strategy for the GSVPF (PIRSA 2017) and is based on fishery-dependent data available since 1991/92 and FIS data available since 2004/05. The assessment does not take into account the period prior to 1991/92 when catches were higher. Consequently, the assessment is made against a 'baseline' that has shifted from an era of relatively high productivity, and the efficacy of the current harvest strategy in buffering reductions in recruitment, or in maintaining its operational objectives in the long term is currently unknown. Under the management plan for the fishery, a review of the current harvest strategy will occur 2½ years after its implementation - 1 July 2017 (PIRSA 2017). A review should include an evaluation of the efficacy of the harvest strategy in achieving the operational objectives of the GSVPF. The bio-economic model of Noell *et al.* (2015) may provide a useful tool for evaluating harvest strategy performance under a range of plausible biological and/or economic scenarios.

The outputs of the GLM used to standardise commercial and FIS CPUE indicate large amounts of unexplained variance. Standardisation procedures for commercial CPUE would benefit from incorporating information that accounts for factors that affect CPUE but are not related to prawn biomass. Further analyses are also required to assess the influence of survey design and other variables on FIS CPUE. Any review of the FIS data would also benefit from comparisons of CPUE from T90 and diamond-mesh codends, and would help to facilitate a potential transition to using only T90 codends in surveys in the future.

More research is also required to understand how recruitment patterns in the GSVPF are affected by the interaction between oceanographic patterns and fishing during the main spawning period (November–December). A biophysical model was recently developed for the Spencer Gulf Prawn Fishery to predict patterns of larval dispersal and settlement by coupling knowledge of the biology and behaviour of Western King Prawn larvae to a hydrodynamic model (McLeay *et al.* 2015). A similar model could be beneficial

to the GSVPF to simulate how changes in spatial fishing patterns and/or environmental conditions could maximise both recruitment and catch during the pre-Christmas period when catch value is highest.

5. REFERENCES

- Beckmann, C. L., Noell C. J. and Hooper, G. E. (2015). Status of the Gulf St Vincent Prawn *Penaeus (Melicertus) latisulcatus* Fishery in 2014/15. Fishery Status Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/00174-4. SARDI Research Report Series No. 870. 25pp.
- Beckmann, C. L. and Hooper, G. E. (2016). Gulf St Vincent Prawn *Penaeus (Melicertus) latisulcatus* Fishery in 2015/16. Fishery Assessment Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/0082-6. SARDI Research Report Series No. 919. 44pp.
- Boxshall, S., Xiao Y. and Williams H. (1999). Gulf St Vincent Prawn. Fishery Assessment Report to PIRSA. SARDI Aquatic Sciences, South Australian Fisheries Assessment Series 99/05.
- Boxshall, S. and Williams H. (2000). Gulf St Vincent Prawn. SARDI Aquatic Sciences, South Australian Fisheries Assessment Series 00/05.
- Boxshall, S. and Johnson J. (2001). Gulf St Vincent Prawn. SARDI Aquatic Sciences, South Australian Fisheries Assessment Series 01/05.
- Broadhurst, M.K., Larsen R.B., Kennelly S.J. and McShane P. (1999). Use and success of composite square-mesh codends in reducing bycatch and in improving size-selectivity of prawns in Gulf St Vincent, South Australia. Fish. Bull. 97: 434–448.
- Bryars, S.R. and Havenhand, J.N. (2006). Effects of constant and varying temperatures on the development of blue swimmer crab (*Portunus pelagicus*) larvae: laboratory observations and field predictions for temperate coastal waters. J. Exp. Mar. Biol. Ecol. 329, 218-229
- Carrick, N.A. (1982). Spencer Gulf Prawn Fishery – surveys increase our knowledge. SAFIC. vol 6(1) pp 3–32.
- Carrick, N.A. (1996). Key factors which affect prawn recruitment and implications to harvesting prawn stocks. FRDC report 91/3, Canberra, Australia.
- Carrick, N. A. (2003). Spencer Gulf Prawn (*Melicertus latisulcatus*) Fishery. Fishery assessment report to PIRSA Fisheries. RD03/0079-2. Adelaide, South Australia: South Australian Research and Development Institute (Aquatic Sciences).

Courtney, A.J. and Dredge M.C.L. (1988) Female reproductive biology and spawning periodicity of two species of king prawns, *Penaeus longistylus* Kubo and *Penaeus latisulcatus* Kishinouye, from Queensland's east coast fishery. Aust. J. Mar. Freshwater Res. 39: 729–741.

Dichmont, C. (2014). A review of the stock assessment methods, processes and outputs for the Gulf St Vincent Prawn Fishery. A report prepared for Primary Industries and Regions SA (PIRSA). CSIRO, Brisbane, Australia, 39 pp.

Dixon C.D. and Sloan S. (2007). Management Plan for the South Australian Gulf St Vincent Prawn Fishery. South Australian Fisheries Management Series. No. 53. Adelaide, South Australia: Primary Industries and Resources South Australia.

Dixon, C.D., Roberts S.D. and Hooper, G.E. (2011). Gulf St Vincent Prawn, *Penaeus (Melicertus) latisulcatus*, Fishery 2009/10. Fishery Assessment report to PIRSA Fisheries. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/000782-4. SARDI Research Report Series No. 526. 121pp.

Dixon, C.D., Hooper, G.E and Burch P. (2012). Gulf St Vincent Prawn, *Penaeus (Melicertus) latisulcatus*, Fishery 2010/11. Fishery Assessment Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/000782-5.

Dixon, C., Raptis, J., Gorman, D., Roberts, S., Hooper, G., Bicknell, N., Sorokin, S., Newman, R., Noell, C., Benediktsson, T., Saint, J. and Hill, Wallace. (2013). A collaborative approach to novel bycatch research for rapid development, extension and adoption into a commercial trawl fishery. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2012/000250-1. SARDI Research Report Series No. 643. 47pp.

Flood, M., Stobutzki, I., Andrews, J., Begg, G., Fletcher, W., Gardner, C., Kemp, J., Moore, A., O'Brien, A., Quinn, R., Roach, J., Rowling, K., Sainsbury, K., Saunders, T., Ward, T., and Winning, M. (eds) (2012). Status of key Australian fish stocks reports 2012, Fisheries Research and Development Corporation, Canberra.

Flood, M., Stobutzki, I., Andrews, J., Ashby, C., Begg, G., Fletcher, R., Gardner, C., McDonald, B., Moore, A., Roelofs, A., Sainsbury, K., Saunders, T., Smith, T., Stewardson, C., Stewart, J., and Wise, B. (2014). Status of key Australian fish stock reports 2014. Fisheries Research and Development Corporation, Canberra.

Gorman, D. and Dixon, C. (2015). Reducing discards in a temperate prawn trawl fishery: a collaborative approach to bycatch research in South Australia. ICES J. Mar. Sci. 72 (9): 2609-2617.

Grey D.L., Dall W. and Baker A. (1983). "A guide to the Australian penaeid Prawns". Department of Primary Production, Northern Territory, Australia.

Hooper G.E., Roberts S.D. and Dixon C.D. (2009). Gulf St Vincent prawn (*Melicertus latisulcatus*) fishery 2007/08 status report. Fishery status report to PIRSA Fisheries. SARDI Aquatic Sciences Publication No. F2007/001074-2, Research Report Series No 328.

Hudinaga, M. (1942). Reproduction, development and rearing of *Penaeus japonicus* Bate. Jap. J. Zool. 10: 305–393.

Jackson, C.J. and Burford, M.A. (2003). The effects of temperature and salinity on growth and survival of larval shrimp *Penaeus semisulcatus* (Decapoda: Penaeoidea). J. Crust. Biol. 23(4), 819-826

Kangas M. (1999). Postlarval and juvenile western king prawns *Penaeus latisulcatus* Kishinouye studies in Gulf St Vincent, South Australia with reference to the commercial fishery. PhD thesis, University of Adelaide.

Kangas, M. I. and Jackson, B. (1997). Gulf St Vincent Prawn Fishery. South Australian Fisheries Assessment Series, Report No. 99/05. Adelaide, South Australia: South Australian Research and Development Institute (Aquatic Sciences).

King, M. G. (1977). The biology of the western king prawn, *Penaeus latisulcatus* Kishinouye, and aspects of the fishery in South Australia. MSc Thesis, University of Adelaide.

Knuckey, I.T., Kompas, T. and Bodsworth A. (2011). Review of the Gulf St Vincent Prawn Fishery. Ocean Grove, Victoria: Cobalt MRM Pty Ltd. Report for PIRSA Fisheries and Aquaculture- September 2011. (Internal report).

Lober, M. and Zeng, C. (2009). Effect of microalgae concentration on larval survival, development and growth of an Australian strain of giant freshwater prawn *Macrobrachium rosenbergii*. Aquaculture 289, 95-100

McLeay, L., Doubell, M., Roberts, S., Dixon, C., Andreacchio, L., James, C., Luick, J. and Middleton, J. South Australian Research and Development Institute (Aquatic Sciences) (2015). Prawn and crab harvest optimisation: a bio-physical management tool.

Final Report to the Fisheries Research and Development Corporation. Adelaide, June. 80pp.

McLeay, L.J., Beckmann, C. L. and Hooper, G. E. (2017). Gulf St Vincent Prawn *Penaeus (Melicertus) latisulcatus* Fishery 2016/17. Fishery Assessment Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/000782-7. SARDI Research Report Series No. 972. 48pp.

McShane, P. (1996). Transfer of by-catch reduction technology to South Australian prawn fisheries. FRDC Report 96/254.02

Morgan, G. (1995). Assessment, management and research support for the Gulf St Vincent Prawn Fishery. South Australian Fish. Man. Series 12.

Morgan, G. and Cartwright G. (2013). Review of the Gulf St Vincent Prawn Fishery management Framework. Adelaide, South Australia. (Internal report).

National Disease and Pest Outbreaks (2018). [White Spot Disease – current situation](#). Accessed 11 September 2018.

Noell, C.J., O'Neill, M.F., Carroll, J.D. and Dixon, C.D. (2015). A bio-economic model for South Australia's prawn trawl fisheries. Final Report. Prepared by the South Australian Research and Development Institute (Aquatic Sciences), Adelaide. CRC Project No. 2011/750. 115pp.

Penn, J.W. (1976). Tagging experiments with the western king prawn *Penaeus latisulcatus* Kishinouye. II. Estimation of population parameters. Aust. J. Mar. Freshwater Res. 27: 239–250.

Penn, J.W. (1980). Spawning and fecundity of the western king prawn, *Penaeus latisulcatus* Kishinouye, in Western Australian waters. Australian Journal of Marine and Freshwater Research 31, 21–35.

Penn, J.W., Hall N.G. and Caputi N. (1988). Resource assessment and management perspectives of the Penaeid prawn fisheries of Western Australia. In: J. Caddy (Ed) The scientific basis of Shellfish Management. John Wiley and Sons, New York.

PIRSA (2017). Management Plan for the South Australian Commercial Gulf St Vincent Prawn Fishery. Paper number 74. PIRSA Fisheries & Aquaculture. 70 pp.

Preston, N. (1985). The combined effects of temperature and salinity on hatching success and the survival, growth and development of the larval stages of *Metapenaeus bennettae* (Racek and Dall). J. Exp. Mar. Biol. Ecol. 85, 57-74

Primavera J.H. and Lebata M.J.H.L. (2000). Size and diel differences in activity patterns of *Metapenaeus ensis*, *Penaeus latisulcatus* and *P. merguiensis*. Mar. Freshwater Behav. Physiol. 33: 173–185.

Rasheed M.A. and Bull C.M. (1992). Behaviour of the western king prawn, *Penaeus latisulcatus* Kishinouye: Effect of food dispersion and crowding. Aust. J. Mar. Freshwater Res. 43: 745–752.

Roberts S.D., Dixon C.D. and Hooper G.E. (2007a). Gulf St Vincent Prawn (*Melicertus latisulcatus*) Fishery 2005/06. Fishery Assessment report to PIRSA Fisheries. SARDI Aquatic Sciences Publication No. F2007/000782-1, Research Report Series No 253.

Roberts S.D., Dixon C.D. and Hooper G.E. (2007b). Gulf St Vincent Prawn (*Melicertus latisulcatus*) Fishery 2006/07 Status Report. Fishery status report to PIRSA Fisheries. SARDI Aquatic Sciences Publication No. F2007/001074-1, Research Report Series No 254.

Roberts S.D., Dixon C.D. and Hooper G.E. (2008). Gulf St Vincent Prawn (*Melicertus latisulcatus*) Fishery 2006/07. Fishery Assessment report to PIRSA Fisheries. SARDI Aquatic Sciences Publication No. F2007/000782-2, Research Report Series No 300.

Roberts S.D., Dixon C.D. and Hooper G.E. (2009). Gulf St Vincent Prawn, *Penaeus (Mericertus) latisulcatus*, Fishery 2007/08. Fishery Assessment report to PIRSA Fisheries. SARDI Aquatic Sciences Publication No. F2007/000782-3, Research Report Series No 382.

Roberts, S.D., Deveney, M. and Sierp, M. (2010). Biosecurity and disease status of prawn nurseries in South Australia. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication Number F2010/000593-1. SARDI Research Report Series No. 449. 35p

Roberts, S.D., Dixon, C.D. and Andreacchio L. (2012). Temperature dependent larval duration and survival of the western king prawn, *Penaeus (Mericertus) latisulcatus* Kishinouye, from Spencer Gulf, South Australia. J. Exp. Mar. Biol. Ecol 411: 14–22.

Stewardson, C., Andrews, J., Ashby, C., Haddon, M., Hartmann, K., Hone, P., Horvat, P., Mayfield, S., Roelofs, A., Sainsbury, K., Saunders, T., Stewart, J., Stobutzki, I. and Wise, B. (2016). Status of Australian Fish Stocks Reports 2016. Canberra, Fisheries Research and Development Corporation.

Svane, I. (2003). Gulf St Vincent Prawn Fishery (*Melicertus latisulcatus*). Fishery assessment report to PIRSA for the Prawn Fishery Management Committee, October 2003. SARDI Aquatic Sciences RD03/0063-2.

Svane, I. and Roberts S.D. (2005). Gulf St Vincent Prawn Fishery (*Melicertus latisulcatus*). Fishery assessment report to PIRSA. SARDI Aquatic Sciences RD03/0063–3.

Svane, I. and Johnson J. (2003). Gulf St Vincent Prawn (*Melicertus latisulcatus*) Fishery. Fishery assessment report to PIRSA. SARDI Aquatic Sciences RD03/0063.

Tanner, J.E. (2003). The influence of prawn trawling on sessile benthic assemblages in Gulf St Vincent, South Australia. *Can. J. Fish. Aquat. Sci.* 60: 517–526.

Tanner, J.E. and Deakin, S. (2001). Active habitat selection for sand by juvenile western king prawns, *Melicertus latisulcatus* (Kishinouye). *J. Exp. Mar. Biol. Ecol.* 261: 199–209.

Wallner, B. (1985). An assessment of the South Australian West Coast western king prawn (*Penaeus latisulcatus*) fishery. Department of Fisheries, South Australia.

Xiao Y. (2004). Use of generalized linear models in analyzing the catch and effort data on the western king prawn *Penaeus latisulcatus* Kishinouye in the Gulf St Vincent, Australia. *Fisheries Research* 68: 67-82.

Xiao, Y. and McShane, P. (1998). Gulf St Vincent Prawn Fishery. SARDI South Australian Fishery Assessment Series 98/05.

Xiao, Y. and McShane, P. (2000). Estimation of instantaneous rates of fishing and natural mortalities from mark-recapture data on the western king prawn *Penaeus latisulcatus* in the Gulf St Vincent, Australia, by conditional likelihood. *Trans. Am. Fish. Soc.* 129: 1005-1017.

Zacharin, W. (1997). Management plan for the South Australian Gulf St Vincent prawn fishery. South Australian Fishery Management Series. Paper no. 30.