

# Inland Waters & Catchment Ecology

## Chowilla Icon Site Fish Assemblage Condition Monitoring 2024



J. Fredberg<sup>1</sup>, C. M. Bice<sup>1</sup> and B. P. Zampatti<sup>2</sup>

SARDI Publication No. F2008/000907-15  
SARDI Research Report Series No. 1247

SARDI Aquatic and Livestock Sciences  
PO Box 120 Henley Beach SA 5022

March 2025

# **Chowilla Icon Site Fish Assemblage Condition Monitoring 2024**

**J. Fredberg<sup>1</sup>, C. M. Bice<sup>1</sup> and B. P. Zampatti<sup>2</sup>**

**SARDI Publication No. F2008/000907-15  
SARDI Research Report Series No. 1247**

**March 2025**

*The South Australian Research and Development Institute respects Aboriginal people as the state's first people and nations. We recognise Aboriginal people as traditional owners and occupants of South Australian land and waters. We pay our respects to Aboriginal cultures and to Elders past, present and emerging.*

This publication may be cited as:

Fredberg, J.<sup>1</sup>, Bice, C.M.<sup>1</sup> and Zampatti, B. P.<sup>2</sup> (2025). Chowilla Icon Site Fish Assemblage Condition Monitoring 2024. South Australian Research and Development Institute (Aquatic and Livestock Sciences), Adelaide. SARDI Publication No. F2008/000907-15. SARDI Research Report Series No. 1247. 69pp.

<sup>1</sup>South Australian Research and Development Institute, Aquatic and Livestock Sciences

<sup>2</sup>CSIRO Environment

## DISCLAIMER

The contents of this publication do not purport to represent the position of the Commonwealth of Australia or the MDBA in any way and are presented for the purpose of informing and stimulating discussion for improved management of the Basin's natural resources. To the extent permitted by law, the copyright holders (including its employees and consultants) exclude all liability to any person for any consequences, including but not limited to all losses, damages, costs, expenses and any other compensation, arising directly or indirectly from using this report (in part or in whole) and any information or material contained in it. The authors warrant that they have taken all reasonable care in producing this report. The report has been through the SARDI internal review process, and has been formally approved for release by the Research Director, Aquatic and Livestock Sciences. Although all reasonable efforts have been made to ensure quality, SARDI does not warrant that the information in this report is free from errors or omissions. SARDI and its employees do not warrant or make any representation regarding the use, or results of the use, of the information contained herein as regards to its correctness, accuracy, reliability and currency or otherwise. SARDI and its employees expressly disclaim all liability or responsibility to any person using the information or advice. Use of the information and data contained in this report is at the user's sole risk. If users rely on the information they are responsible for ensuring by independent verification its accuracy, currency or completeness. The SARDI Report Series is an Administrative Report Series which has not been reviewed outside the department and is not considered peer-reviewed literature. Material presented in these Administrative Reports may later be published in formal peer-reviewed scientific literature.

## © 2025 SARDI & DEW

This work is copyright. Apart from any use as permitted under the *Copyright Act* 1968 (Cth), no part may be reproduced by any process, electronic or otherwise, without the specific written permission of the copyright owner. Neither may information be stored electronically in any form whatsoever without such permission. With the exception of the Commonwealth Coat of Arms, the Murray-Darling Basin Authority logo and photographs, all material presented in this document is provided under a Creative Commons Attribution 4.0 International licence (<https://creativecommons.org/licenses/by/4.0/>)




For the avoidance of any doubt, this licence only applies to the material set out in this document.

The details of the licence are available on the Creative Commons website (accessible using the links provided) as is the full legal code for the CC BY 4.0 licence (<https://creativecommons.org/licenses/by/4.0/legalcode>) Source: Licensed from the Department for Environment and Water (DEW) under a Creative Commons Attribution 4.0 International Licence. Enquiries regarding the licence and any use of the document are welcome to: Jan Whittle, Chowilla Icon Site Coordinator [jan.whittle@sa.gov.au](mailto:jan.whittle@sa.gov.au)

Author(s): J. Fredberg<sup>1</sup>, C. M. Bice<sup>1</sup> and B. P. Zampatti<sup>2</sup>

Reviewer(s): K. Frahn (SARDI) and J. Whittle (DEW)

Approved by: Q. Ye  
Program Leader – Inland Waters & Catchment Ecology

Signed: 

Date: 19 March 2025

Distribution: DEW, MDBA, SARDI Aquatic and Livestock Sciences, Parliamentary Library, State Library and National Library

Circulation: OFFICIAL

## ALL ENQUIRIES

South Australian Research and Development Institute – Aquatic and Livestock Sciences  
2 Hamra Avenue West Beach SA 5024  
PO Box 120 Henley Beach SA 5022  
P: (08) 8207 5400 F: (08) 8207 5415  
E: [pirs.sardiaquatics@sa.gov.au](mailto:pirs.sardiaquatics@sa.gov.au) W: <http://www.pir.sa.gov.au/research>

## TABLE OF CONTENTS

ACKNOWLEDGEMENTS .....	VIII
EXECUTIVE SUMMARY .....	1
1. INTRODUCTION .....	3
2. METHODS .....	5
<i>Abundance</i> .....	9
<i>Data analysis</i> .....	9
<i>Recruitment of fish species (Ecological Objective 11)</i> .....	13
3. RESULTS .....	17
<i>Hydrology during study period</i> .....	17
<i>Catch Summary</i> .....	18
<i>Abundance of native fish</i> .....	19
<i>Abundance of non-native fish</i> .....	19
<i>Temporal variation in fish abundance</i> .....	21
<i>Spatio-temporal differences in fish assemblage structure</i> .....	22
<i>Targeted Murray cod sampling</i> .....	26
<i>Diversity and extent of fish species (Ecological Objective 10)</i> .....	28
<i>Recruitment of small- to medium-bodied native species</i> .....	32
<i>Recruitment of large-bodied native species</i> .....	33
<i>Recruitment of non-native species</i> .....	35
4. DISCUSSION .....	38
5. CONCLUSIONS .....	43
REFERENCES .....	45
APPENDIX 1 .....	50

## LIST OF FIGURES

<b>Figure 1.</b> Map of the Chowilla Anabranh and Floodplain system and the adjacent River Murray main channel showing the fish condition monitoring sites 1–22. ....	5
<b>Figure 2.</b> Regions sampled to target Murray cod in June 2024: Pipeclay Creek (yellow), Salt Creek (red), Slaney and Little Slaney Creek (blue), Chowilla Creek downstream bridge (green), Chowilla Creek upstream bridge (purple) and River Murray main channel (orange). ....	8
<b>Figure 3.</b> Mean daily flow (ML.d <sup>-1</sup> ) in the River Murray at the South Australian Border (Site A42610010) January 2004–July 2024 (MDBA, unpublished data). Closed red circles indicate sampling events, the dotted line represents approximate bankfull discharge at Chowilla (~35,000 ML.d <sup>-1</sup> ) and the blue dashes represent Chowilla Regulator operations. ....	18
<b>Figure 4.</b> Mean ( $\pm$ SE) catch-per-unit-effort (CPUE) (fish.min <sup>-1</sup> ) of fish (all species combined) collected annually during standardised boat electrofishing surveys from 2005–2024 at 14–22 sites in the Chowilla Anabranh system and adjacent River Murray (dark grey = proportion native species, light grey = proportion of non-native species). ....	21
<b>Figure 5.</b> a) Dendrogram indicating fish assemblage clusters across sampling years from 2005–2024. b) Non-metric multi-dimensional scaling (MDS) plot of fish assemblages sampled from all years/sites combined. ....	23
<b>Figure 6.</b> Non-metric multi-dimensional scaling (MDS) plots of a) fast-flowing, b) slow-flowing, c) backwater and d) river mesohabitats sampled from all years/sites combined (excluding 2011). ....	24
<b>Figure 7.</b> Relative abundance (fish.min <sup>-1</sup> ) of Murray cod from targeted Murray cod surveys and condition monitoring combined since 2014. ....	28
<b>Figure 8.</b> Diversity indices for a) fast-flowing, slow-flowing, backwater and main channel mesohabitats and b) the calculated Icon Site Diversity Index (DI), at the Chowilla Icon Site from 2005–2024. ....	29
<b>Figure 9.</b> Extent Index (EI) scores for a) large-bodied native species and b) small- to medium-bodied native species at the Chowilla Icon Site from 2005–2024. Black dashed line represents extent equal to the reference, green dashed line extent 25% greater than reference and red dashed line extent 25% lesser than reference. ....	31
<b>Figure 10.</b> Recruitment Index ( <i>RI</i> ) values for unspotted hardyhead, Murray Rainbowfish, Australian smelt and bony herring from 2005–2024. Values for 2011 are not presented as sampling occurred at atypical time of year due to flooding. Dashed black line represents recruitment equal to the reference value and the dashed red line, recruitment 75% of the reference value. ....	32
<b>Figure 11.</b> Recruitment Index ( <i>RI</i> ) values for a) Murray cod ranging 400–600 mm TL and b) YOY Murray cod (<200 mm TL) from 2005–2024. Values for 2011 are not presented as sampling	

occurred at an atypical time of year due to flooding. Dashed black line represents recruitment equal to the reference value and the dashed red line, recruitment 75% of the reference value. 33

**Figure 12.** Recruitment Index (RI) values for golden perch from 2005–2024. Dashed black line represents recruitment equal to the reference value and the dashed red line, recruitment 75% of the reference value. .... 34

**Figure 13.** Length distribution of common carp captured at all sites sampled within Chowilla and the adjacent River Murray main channel from 2005–2024. .... 36

**Figure 14.** Length distribution of goldfish captured at all sites sampled within Chowilla and the adjacent River Murray main channel from 2005–2024. .... 37

## LIST OF TABLES

**Table 1.** Site number, location and mesohabitat type of fish condition monitoring sites surveyed within the Chowilla Anabranh system and adjacent River Murray main channel from 2005–2024. Asterisks denote years when sites were surveyed. (d/s = downstream, u/s = upstream)..... 6

**Table 2.** Summary of rarity scores (RS), interpretation of expectedness ratio (*ER*) and expectedness weight to be assigned to fish species at the Chowilla Icon Site..... 11

**Table 3a.** Rarity scores, expectedness ratio and expectedness weight for all native species sampled at the icon site within ‘fast-flowing’ mesohabitats..... 11

**Table 4.** Species, typical length of the YOY cohort during annual sampling (based upon knowledge of species biology), the mean proportion of the population comprised by the YOY cohort (*r<sub>standard</sub>*) and the recruitment index reference value (*RV*)..... 15

**Table 5.** Total and standardised (fish.site<sup>-1</sup>) abundances of fish captured from condition monitoring sites sampled in the Chowilla Anabranh system and adjacent River Murray 2005–2024. .... 20

**Table 6.** PERMANOVA results comparing the relative abundances of fish between years and mesohabitats from 2005–2024, excluding 2011. Significant *P* values are highlighted in bold.... 22

**Table 7.** PERMANOVA pair-wise comparisons between fish assemblages among different mesohabitats in Chowilla from 2005-2024, excluding 2011. Significant *p* values are highlighted in bold (B-Y corrected  $\alpha = 0.02$ )..... 22

**Table 8.** Indicator species analysis comparing the relative abundance of fish amongst years from 2005–2024, including 2011. (Year group 1 = 2005–2010, 2015–2016 and 2019–2022; Year group 2 = 2012–2014; Year Group 3 = 2011; Year Group 4 = 2017; Year Group 5 = 2023; and Year Group 6 = 2024). Significant indicators ( $\alpha = 0.05$ ) signifies that a species occurs in a higher relative abundance in a specific year group. Indicators that are not significant signify that a species

was either sampled in similar numbers in more than one year (widespread across years) or captured in low abundances (uncommon).....25

**Table 9.** Indicator species analysis comparing the relative abundance of fish in three of the four aquatic mesohabitats from 2005–2024, excluding 2011 (no species were found to be associated with slow-flowing mesohabitats). A significant difference ( $\alpha = 0.05$ ) indicates that a species occurs in a higher relative abundance in a specific mesohabitat. Values that are not significant indicate that a species was either sampled in similar numbers in more than one mesohabitat (widespread) or captured in low abundances (uncommon).....26

**Table 10.** Total and standardised catch-per-unit-effort (CPUE) (fish.min<sup>-1</sup>) of Murray cod from targeted sampling and condition monitoring in the Chowilla Anabranh system and adjacent River Murray between 2014–2024. ....27

## **ACKNOWLEDGEMENTS**

Thanks to the SARDI staff who assisted with field sampling, laboratory processing and previous report writing for this project; namely, Sandra Leigh, Ian Magraith, Paul Jennings, David Flear, Phillipa Wilson, Neil Wellman, David Short, Arron Strawbridge, Mark Sutcliffe, Leigh Thwaites, David Schmarr, Ruan Gannon, Travis Howson and Thiago Vasques Mari.

Jan Whittle, Tony Herbert, Alison Stokes, Todd Wallace, Richard Watts and Mark Schultz (Department for Environment and Water, DEW) managed the project at different stages, developed targets and provided input on site selection. Robbie Bonner, Tim Kruger, Warren Beer and Tony Waye (SA Water) provided space for us to live and generous hospitality at Lock 6. Thanks to Alison Stokes (DEW) for facilitating access to various parts of the Chowilla Game Reserve. Thanks also to Scott Huntley and Sylvia Zukowski of Nature Glenelg Trust (NGT) for use of their electrofishing vessel.

Thanks to Kate Frahn (SARDI) and Jan Whittle (DEW) who constructively reviewed a draft of this report. The report was formally approved for release by Associate Professor Qifeng Ye, Program Leader, Inland Waters & Catchment Ecology, SARDI Aquatic and Livestock Sciences. Funding for the 2024 condition monitoring was provided by the Murray-Darling Basin Authority (MDBA) Living Murray Initiative.

“The Living Murray is a joint initiative funded by the New South Wales, Victorian, South Australian and Commonwealth governments, coordinated by the Murray-Darling Basin Authority.”

## EXECUTIVE SUMMARY

The Chowilla Anabranch and Floodplain system is the largest remaining area of undeveloped floodplain habitat in the lower River Murray. Chowilla consists of a range of aquatic habitats that are now rare in the region, including permanently flowing creeks, and these habitats support a diverse native fish community. The Chowilla Floodplain, however, has become increasingly degraded due to changes to the natural flow regime, grazing and an extended period (2001–2010) of low flows in the Murray-Darling Basin. To 'enhance and restore' the environmental values of the Chowilla Floodplain, an Asset Environmental Management Plan was developed as part of the Chowilla Integrated Natural Resource Management Project. Subsequently, a Chowilla Floodplain Environmental Water Management Plan (MDBA 2012) has been developed with refined ecological objectives as follows:

- Ecological Objective 10: Maintain or increase the diversity and extent of distribution of native fish species.
- Ecological Objective 11: Maintain successful recruitment of small- and large-bodied native fish.

To assist with monitoring of Ecological Objectives 10 and 11, quantitative fish surveys have been undertaken annually in the Chowilla system since 2005. Sites have been selected to represent all aquatic mesohabitats present within the region. Since 2014, additional targeted surveys have been conducted for Murray cod (*Maccullochella peelii*).

In 2024, 18 of the 22 condition monitoring sites were sampled within Chowilla and the adjacent River Murray main channel; sampling was conducted via a hired electrofishing vessel due to the breakdown of the typical SARDI vessel. A total of 6,684 fish were sampled from 14 species. The fish assemblage consisted of 10 native and 4 non-native species, with bony herring (80% of total catch) (*Nematalosa erebi*), Australian smelt (2.5%) (*Retropinna semoni*) and unspotted hardyhead (*Craterocephalus stercusmuscarum fulvus*) (1.9%) the most abundant native species. Of the non-native species, common carp (*Cyprinus carpio*) were most abundant and comprised ~12% of the total catch.

The fish assemblage in 2024 was significantly different to all other years, but was most closely aligned to years characterised by persistent high flows (e.g. 2012–2014).

Following persistent high flows, fish assemblages at Chowilla are typically characterised by low overall abundance, but moderate–high abundances of medium to large-bodied native and non-native species including bony herring and common carp. Native small-bodied generalist species such as Australian smelt, unspocked hardyhead and Murray rainbowfish are typically less abundant in these years.

Annual data from 2005–2024 indicate that Objective 10 and 11 of the Icon Site management plan were met. Over the 20-year sampling period, species diversity by mesohabitat and for the overall site have been generally consistent, although a slight decrease was observed in 2024. The extent of most species throughout the available aquatic mesohabitats was maintained, but like diversity, in 2024, a decrease in extent for most species was also observed. Over time, fish species also exhibited fidelity to specific mesohabitats. The native species Murray cod, golden perch, silver perch (*Bidyanus bidyanus*), Australian smelt and freshwater catfish (*Tandanus tandanus*) characterised fast-flowing mesohabitats; Murray rainbowfish, unspocked hardyhead and redfin perch characterised river mesohabitats, and the non-native species, goldfish, characterised backwaters.

In 2024, recruitment was evident for all small-bodied species, but indices were below reference values for Australian smelt, unspocked hardyhead and Murray rainbowfish. The reference value, however, was achieved for the medium-bodied native species, bony herring. Recruitment of Murray cod to young-of-year (YOY; <200 mm TL) was also observed, as was recruitment to reproductive maturity (400–600 mm TL), however, this was less than the reference value. The golden perch recruitment index did not exceed the reference value in 2024.

**Keywords:** Chowilla, extent, diversity, recruitment, flow, native, non-native, Murray cod.

## 1. INTRODUCTION

The Chowilla Anabranched and Floodplain system (hereafter Chowilla) comprises the largest remaining area of undeveloped floodplain habitat in the lower River Murray. It encompasses a series of anabranching creeks, backwaters, wetlands and terminal lakes that bypass Lock and Weir No. 6 (hereafter Lock 6) on the River Murray. Chowilla is part of the Riverland Ramsar site and listed in the directory of important wetlands in Australia as a Wetland of national importance for nationally threatened species, habitats and communities (Environment Australia 2001). Chowilla is also recognised as an *Icon Site* under the Murray-Darling Basin Authority's (MDBA) *The Living Murray Program* (TLM) (MDBA 2016).

The lower River Murray, downstream of the Darling River junction, is regulated by 10 low level (~3 m high) weirs that have substantially altered the hydrology and hydraulics of the river. The combination of short distances (29–86 km) and low gradients (50 mm.km<sup>-1</sup>) between the weirs, and low regulated flows, has produced a shift from hydrodynamically variable lotic habitats to relatively stable lentic habitats more representative of a series of interconnected lakes (Walker 2006, Bice *et al.* 2017). Due to the ~3 m of head differential created by Lock 6, 20–90% of the River Murray flow is diverted through Chowilla under low flow conditions (i.e. main-channel flow to South Australia <10,000 ML.d<sup>-1</sup>) (Stace and Greenwood 2004). Consequently, Chowilla exhibits a range of permanent lotic (flowing water) habitats in what previously would have been a combination of perennial and ephemeral streams. These lotic habitats that characterise Chowilla, are now rare within the lower River Murray (Bice *et al.* 2017, Mallen-Cooper and Zampatti 2018).

Chowilla supports a wide range of aquatic organisms (O'Malley and Sheldon 1990), including a diverse native fish community (Lloyd 1990; Pierce 1990; Zampatti *et al.* 2011). The floodplain became increasingly degraded due to changes to the natural flow regime, grazing and drought (MDBC 2006). In response, and to 'enhance and restore' the environmental values of the Chowilla Floodplain system, the Department for Environment and Water (DEW) developed an Environmental Water Management Plan (EWMP) as part of the Chowilla Integrated Natural Resource Management Project (DWLBC 2006). Four preliminary management targets were developed for fish:

- Target 10. Maintain the diversity and extent of distribution of native fish species.

- Target 11. Reduce barriers to fish passage.
- Target 12. Maintain successful recruitment of small-bodied native fish every year.
- Target 13. Maintain successful recruitment of large-bodied fish at least once every five years.


Subsequently, the Chowilla Floodplain Environmental Water Management Plan (MDBA 2012) was developed with refined ecological objectives:

- Ecological Objective 10. Maintain or increase the diversity and extent of distribution of native fish species.
- Ecological Objective 11. Maintain successful recruitment of small- and large-bodied native fish.

Annual quantitative (standardised electrofishing) fish surveys have been undertaken at Chowilla since 2005. Data from these surveys are used to investigate spatial and temporal variability in the fish assemblage (i.e. species diversity, distribution and abundance) at Chowilla (Ecological Objective 10) and assess evidence for the recruitment of small- and large-bodied fishes (Ecological Objective 11). This report presents the results of fish condition monitoring undertaken in 2024 with reference to results from 2005–2023.

## 2. METHODS

Fish condition monitoring at Chowilla was initially undertaken in 2005 (Zampatti *et al.* 2008). Eighteen sites were selected, representing the range of permanent aquatic mesohabitats present within Chowilla (i.e. fast-flowing anabranches, slow-flowing anabranches, backwaters and the River Murray main channel) as described by Sheldon and Lloyd (1990). These sites were initially assigned to a mesohabitat category based on visual assessments (Table 1) and were later quantified and, if necessary, revised following the measurement of cross-sectional water velocity profiles in March 2007. Fast-flowing habitats were characterised as having mean cross-sectional velocities  $>0.18 \text{ m.s}^{-1}$ , slow-flowing habitats  $0.05\text{--}0.18 \text{ m.s}^{-1}$ , backwaters  $<0.05 \text{ m.s}^{-1}$  and River Murray main channel  $<0.1 \text{ m.s}^{-1}$  (Zampatti *et al.* 2008). Four additional sites in the New South Wales section of Chowilla were added to the monitoring program in 2008 and 2009 (Leigh *et al.* 2010), providing the current network of 22 sites (Figure 1).



**Figure 1.** Map of the Chowilla Anabranch and Floodplain system and the adjacent River Murray main channel showing the fish condition monitoring sites 1–22.


**Table 1.** Site number, location and mesohabitat type of fish condition monitoring sites surveyed within the Chowilla Anabranch system and adjacent River Murray main channel from 2005–2024. Asterisks denote years when sites were surveyed. (d/s = downstream, u/s = upstream).

Site No.	Location	Mesohabitat type	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
1	Chowilla Creek d/s Monoman Creek	Slow-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2	Chowilla Creek u/s of Boat Creek	Fast-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
3	Chowilla Creek d/s Slaney Creek	Fast-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
4	Boat Creek u/s vehicle bridge	Fast-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
5	Swiftys Creek d/s Bank I	Fast-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
6	Pipeclay Creek d/s Pipeclay Weir	Fast-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
7	Slaney Creek d/s Slaney Weir	Fast-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
8	Slaney Creek d/s Salt Creek junction	Fast-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
9	Slaney Billabong	Backwater	*	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
10	Hypurna Creek at Wilkadene	Slow-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
11	Punkah Creek d/s Punkah Island	Slow-flowing	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
12	Punkah Creek at Lake Littra	Slow-flowing	*	*	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
13	River Murray 5-7 km d/s Lock 6	Main River Channel	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
14	River Murray d/s Lock 6	Main River Channel	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
15	Isle of Mann backwater	Backwater	*	*	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
16	Monoman Creek at campsite 9	Backwater	*	*	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
17	Salt Creek at cliffs (NSW)	Slow-flowing					*	*		*	*	*	*	*	*	*	*	*	*	*	*	*
18	River Murray at Border Cliffs (NSW)	Main River Channel	*				*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
19	Salt Creek d/s Bank K (NSW)	Fast-flowing				*	*	*	*	*	*	*	*	*	*	*	*		*	*	*	
20	Salt Creek at NSW border (NSW)	Slow-flowing					*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
21	Salt Creek near Tareena Billabong	Slow-flowing					*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
22	Pilby Billabong	Backwater	*					*	*	*	*	*	*	*	*	*	*	*		*		*
<b>Total number of sites</b>			<b>18</b>	<b>16</b>	<b>15</b>	<b>14</b>	<b>21</b>	<b>22</b>	<b>21</b>	<b>22</b>	<b>22</b>	<b>21</b>	<b>22</b>	<b>21</b>	<b>22</b>	<b>21</b>	<b>22</b>	<b>21</b>	<b>21</b>	<b>21</b>	<b>22</b>	<b>18</b>

In 2024, 18 of the 22 sites were sampled from February 26 to March 21 (Figure 1, Table 1). Three sites were not sampled (Swifty's Creek, Bank K and Salt Creek at NSW border) due to electro-fishing vessel breakdown, while a further site was dry (Pilby Billabong) (Table 1). Annual condition monitoring from 2005–2024 (except for 2011) was conducted in March–April to maximise the likelihood that young-of-year (YOY) individuals from the preceding spring/summer spawning season were represented in the catch, enabling the recruitment of individual species to be assessed.

In 2024, the breakdown of the SARDI vessel necessitated the use of a different vessel to previous surveys, albeit of the same technical specifications. The implications of this alternative vessel use are discussed later in this report. Fish surveys were conducted using a vessel mounted 5 kW Smith Root Model GPP electrofishing system. At each site, 12 (6 on each bank) x 90 second (power on time) electrofishing shots were undertaken during daylight hours. All fish were dip-netted and placed in holding tanks. Any positively identified fish unable to be dip netted were recorded as “observed” and included in the total catch. Fish from each shot were identified, counted, measured for length ( $\pm 1$  mm, caudal fork length, FL or total length, TL) and released after processing. Where large numbers of an individual species were collected, a sub sample of 20 individuals were measured for length.

In addition to February/March sampling, in June/July 2024, targeted sampling was conducted for Murray cod (*Maccullochella peelii*) to provide additional data to assess Objective 11 (maintain successful recruitment of small- and large-bodied native fish). Spatio-temporally specific sampling for Murray cod was adopted in 2014 following collection of juvenile Murray cod (i.e. <200 mm TL) in targeted sampling in May 2013 that were not detected through standard condition monitoring in March 2013 (Wilson *et al.* 2014). From 25 June – 2 July 2024, five sites: Chowilla Creek downstream of bridge, Chowilla Creek upstream of bridge, Slaney Creek, Pipeclay Creek and River Murray main channel (Figure 2) were sampled using the methods described above. At each site, effort (power-on time) was increased, compared to standardised condition monitoring, and only specific habitats were targeted (e.g. large wood and flowing water).



**Figure 2.** Regions sampled to target Murray cod in June 2024: Pipeclay Creek (yellow), Slaney and Little Slaney Creek (blue), Chowilla Creek downstream bridge (green), Chowilla Creek upstream bridge (purple) and River Murray main channel (orange).

### *Abundance*

Fish abundance is not a specific target in the Environmental Water Management Plan; nevertheless, changes in abundance may reflect environmental conditions. The abundance of individual fish species was investigated over the twenty sampling years. Total and standardised abundances are presented for each species in each year. Total abundances for each fish species were calculated as the number of fish captured in addition to the number of fish observed. Standardised abundances were calculated by dividing the total number calculated for each species by the number of sites sampled in that year (i.e. fish.site<sup>-1</sup>).

### *Data analysis*

Differences in the relative abundance (CPUE, fish.min<sup>-1</sup>) of fish sampled between years were analysed using uni-variate single-factor PERMANOVA (permutational ANOVA) (Anderson and Ter Braak 2003), performed on similarity matrices calculated using Euclidean distances, in the package PRIMER v. 6.1.12 and PERMANOVA+ (Anderson *et al.* 2008).

Differences in fish assemblages from 2005–2024 (excluding 2011) and among mesohabitats (fast-flowing, slow-flowing, backwater and River Murray main channel) were analysed using two-factor permutational multivariate analysis of variance (PERMANOVA) (Anderson 2001; Anderson and Ter Braak 2003), Non-Metric Multi-Dimensional Scaling (MDS), cluster analysis (McCune *et al.* 2002) and indicator species analysis (Dufrene and Legendre 1997). All multi-variate PERMANOVA analyses and MDS ordinations were performed on Bray-Curtis (1957) similarity matrices. For these analyses, a significance level of  $\alpha = 0.05$  was adopted and the B-Y significance correction was used for pairwise comparisons (Benjamini and Yekutieli 2001).

Group average clustering was performed on pooled site data (individual species CPUE, fish.min<sup>-1</sup> for each year), and a cut off score of 85% similarity was used to determine the cluster groups based on species abundance. Indicator Species Analysis was then undertaken with the software package PCOrd v. 5.12 (McCune and Mefford 2005) to determine species that characterised assemblages in different clusters and species mesohabitat preferences. Indicator species analysis combines information on the concentration of species abundance in a particular group and the faithfulness of occurrence of a species in a particular group (McCune *et al.* 2002). A perfect indicator of a particular group should be faithful to that group (always present) and exclusive to that group (never occurring in other groups) (McCune *et al.* 2002). This test produces indicator values for each species in each group based on the standards of the perfect indicator.

Statistical significance of each indicator value is tested by a Monte Carlo (randomisation) technique, where the real data are compared against 5000 runs of randomised data (Dufrene and Legendre 1997). A species that is deemed not to be a significant indicator of a particular group is either uncommon or widespread. An uncommon species is found only in one group but in low numbers and a widespread species is found in more than one group in similar numbers (Dufrene and Legendre 1997). A species was classed as a widespread or uncommon non-significant species by examination of the raw data.

#### *Diversity and extent of fish species (Ecological Objective 10)*

In 2024, sites were grouped into aquatic mesohabitat categories (Table 1) with the diversity and extent of native fish species calculated using the methodology defined by Robinson (2013) and outlined below.

#### *Diversity*

The diversity reference and index were derived using an 'expected vs predicted' approach adapted from the SRA method of Robinson (2013). An expectedness weight was developed for each native species for each mesohabitat type based upon all sampling data from 2005–2024. Sampling conducted from 2005–2014 was used to calculate all reference values, as no suitable 'baseline' data on fish diversity and extent are available for the icon site prior to the TLM program. This period incorporates a range of hydrological conditions (drought and flood) and reference values derived from this dataset were deemed suitable for assessing site condition through time. For each mesohabitat, the proportion of sites at which a species was sampled was calculated for every sampling year, and the mean of this value across years (hereafter called 'expectedness ratio') was used to calculate the 'expectedness weight' for each species (Table 2). Rarity scores were also assigned to each native species based upon expert opinion (Robinson 2013) (Table 2).

**Table 2.** Summary of rarity scores (RS), interpretation of expectedness ratio (ER) and expectedness weight to be assigned to fish species at the Chowilla Icon Site.

RS	Interpretation (expectedness ratio, ER)	Expectedness weight
1	Either rare or cryptic species. Expected to be collected in up to 20% of sites in the Chowilla region.	0.10
3	Locally abundant species. Expected to be collected in 20 to 70 % of sites in the Chowilla region.	0.45
5	Common and abundant species. Expected to be collected in 70 to 100% of sites in the Chowilla region.	0.85
0	Native species not historically recorded in the Chowilla region. Not included in expectedness calculations.	0

Rarity scores, mean expectedness ratio (i.e. proportional presence of native fish within mesohabitats) and associated expectedness weights for all native fish species using the standardised method at the icon site are presented below in Tables 3a–d. These metrics are presented separately for each mesohabitat type. Non-native species are not included in diversity calculations.

**Table 3a.** Rarity scores, expectedness ratio and expectedness weight for all native species sampled at the icon site within 'fast-flowing' mesohabitats.

Species	Rarity score	Expectedness ratio	Expectedness weight
Australian smelt	5	0.96	0.85
Bony herring	5	1	0.85
Carp gudgeon complex	3	0.66	0.45
Dwarf flat-headed gudgeon	1	0.04	0.1
Flat-headed gudgeon	3	0.43	0.45
Freshwater catfish	3	0.23	0.45
Golden perch	5	1	0.85
Murray cod	3	0.57	0.45
Murray rainbowfish	5	0.90	0.85
Silver perch	3	0.39	0.45
Spangled perch	0	0	0
Unspecked hardyhead	5	0.77	0.85
<b>Predicted no. species</b>			<b>11</b>
<b>Expected no. species</b>			<b>6.6</b>

**Table 3b.** Rarity scores, expectedness ratio and expectedness weight for all native species sampled at the icon site within 'slow-flowing' mesohabitats.

Species	Rarity score	Expectedness ratio	Expectedness weight
Australian smelt	5	0.83	0.85
Bony herring	5	1	0.85
Carp gudgeon complex	3	0.64	0.45
Dwarf flat-headed gudgeon	1	0.05	0.1
Flat-headed gudgeon	3	0.61	0.45
Freshwater catfish	3	0.06	0.1
Golden perch	5	0.94	0.85
Murray cod	3	0.06	0.1
Murray rainbowfish	5	0.79	0.85
Silver perch	3	0.20	0.1
Spangled perch	0	0.02	0
Unspecked hardyhead	5	0.75	0.85
<b>Predicted no. species</b>			<b>11</b>
<b>Expected no. species</b>			<b>5.55</b>

**Table 3c.** Rarity scores, expectedness ratio and expectedness weight for all native species sampled at the icon site within 'river main channel' mesohabitats.

Species	Rarity score	Expectedness ratio	Expectedness weight
Australian smelt	5	0.9667	0.85
Bony herring	5	1	0.85
Carp gudgeon complex	3	0.7	0.45
Dwarf flat-headed gudgeon	1	0.1667	0.10
Flat-headed gudgeon	3	0.65	0.45
Freshwater catfish	3	0.267	0.45
Golden perch	5	0.96667	0.85
Murray cod	3	0.1833	0.10
Murray rainbowfish	5	1	0.85
Silver perch	3	0.283	0.45
Spangled perch	0	0.033	0
Unspecked hardyhead	5	0.9667	0.85
<b>Predicted no. species</b>			<b>11</b>
<b>Expected no. species</b>			<b>6.25</b>

**Table 3d.** Rarity scores, expectedness ratio and expectedness weight for all native species sampled at the icon site within 'backwater' mesohabitats.

Species	Rarity score	Expectedness ratio	Expectedness weight
Australian smelt	5	0.66	0.45
Bony herring	5	1	0.85
Carp gudgeon complex	3	0.70	0.45
Dwarf flat-headed gudgeon	1	0.08	0.10
Flat-headed gudgeon	3	0.483	0.45
Freshwater catfish	3	0.1	0.10
Golden perch	5	0.7083	0.45
Murray cod	3	0	0
Murray rainbowfish	5	0.7	0.45
Silver perch	3	0.125	0.10
Spangled perch	0	0	0
Unspecked hardyhead	5	0.9	0.85
<b>Predicted no. species</b>			<b>10</b>
<b>Expected no. species</b>			<b>4.25</b>

The diversity index (DI) (i.e. no. species sampled/expected no. species) was calculated for each site within a mesohabitat. The mesohabitat diversity index presented in the results (Figure 8) is the mean of these site-specific indices from all sites within a particular mesohabitat. In turn, the icon site score is the mean of the mesohabitat indices. Values of DI >1.0 indicate diversity greater than the reference, whilst values <1.0 indicate diversity less than the reference.

### Extent

An Extent Index (*EI*) was developed using the expectedness ratios calculated above (Tables 3a–d) as the ‘reference value’ (Robinson 2013). The expectedness ratio represents the mean distribution of individual native species across a mesohabitat type (i.e. proportion of sites within a mesohabitat where the species was sampled), annually across the entire study period (2005–2024).

The extent index is species-specific and is calculated as outlined below.

- *MH* = mesohabitat,
- *R<sub>year</sub>* = ratio of sites sampled in given year,
- *ER* = expected ratio for each mesohabitat type,
- *EI* = Icon Site Extent Index,
- $EI = \text{mean}(MH_1(R_{year}/ER_{MH1}) + MH_2(R_{year}/ER_{MH2}) + MH_3(R_{year}/ER_{MH3}) + MH_4(R_{year}/ER_{MH4}))$ ,
  - *EI* = 0.75–1.25 represent stable extent/distribution
  - *EI* >1.25 represents increased extent/distribution
  - *EI* <0.75 represents decreased extent/distribution

Species with rarity scores of 0 (i.e. spangled perch (*Leipotherapon unicolour*)) or 1 (i.e. dwarf flat-headed gudgeon (*Philynodon macrostomus*)) were excluded. Furthermore, Murray cod do not have an expectedness ratio in backwater mesohabitats, as they are rarely sampled in this mesohabitat type in Chowilla.

### *Recruitment of fish species (Ecological Objective 11)*

Recruitment indices were developed for six native species representative of the varying life history strategies exhibited by fish of the region (see Winemiller and Rose 1992). These life history groupings are based on longevity, biology/ecology and spatial scale of life history, and thus, the potential to be influenced by icon site-scale management. Reference and index values were calculated as per Robinson (2013) for: 1) four small- to medium-bodied generalist species, 2)

Murray cod (an equilibrium strategist) and 3) golden perch (*Macquaria ambigua*, a periodic strategist). Recruitment indices were not developed for non-native species such as common carp (*Cyprinus carpio*) and goldfish (*Carassius auratus*), but length frequency distributions are presented as a means to document recruitment.

### Small- and medium-bodied species

References and indices were developed for the following generalist species,

- Unspecked hardyhead (*Craterocephalus stercusmuscarum fulvus*);
- Murray rainbowfish (*Melantaenia fluviatilis*);
- Australian smelt (*Retropinna semoni*); and
- Bony herring (*Nematalosa erebi*).

The index for small-bodied fish incorporates both age/size structure and abundance. These species are short-lived (1–5 years) and are thus reliant upon annual recruitment. In most species, fish comprising the YOY cohort in autumn will contribute to the reproductively mature adult population the following spawning season. Abundance is included in the index, as reliance on an age/size structure alone may result in years where few fish are sampled being classified as years of ‘successful’ recruitment. For these species, length is an appropriate surrogate for true measures of age (e.g. otolith increment counts).

The reference value (Table 4) is the mean abundance of the YOY cohort from baseline data collected from 2005–2014, and is calculated as:

- $X$  = icon site abundance (fish.minute electrofishing<sup>-1</sup>.site<sup>-1</sup>),
- $r_{standard}$  = set proportion YOY\* (\*Values of  $r_{standard}$  were calculated as the mean proportion of the population comprised of young-of-the-year from 2005–2014.),
- Reference value (RV) =  $\text{mean}((X_{2005} * r_{standard}) + (X_{2006} * r_{standard}) + (X_{2007} * r_{standard}) + \dots + (X_{2014} * r_{standard}))$

**Table 4.** Species, typical length of the YOY cohort during annual sampling (based upon knowledge of species biology), the mean proportion of the population comprised by the YOY cohort ( $r_{standard}$ ) and the recruitment index reference value ( $RV$ ).

Species	Length YOY	$r_{standard}$	RV
Unspecked hardyhead	<40 mm FL	65%	2.36
Murray rainbowfish	<40 mm FL	25%	0.20
Australian smelt	<40 mm FL	40%	0.57
Bony herring	<100 mm FL	65%	12.64

The annual recruitment index for small-bodied species was calculated as:

- $X_{year}$  = annual abundance (fish.minute electrofishing<sup>-1</sup>.site<sup>-1</sup>),
- $r_{year}$  = annual proportion of YOY
- Annual recruitment value ( $AV$ ) =  $X_{year} * r_{year}$
- Recruitment index ( $RI$ ) =  $AV/RV$ 
  - Values of  $RI > 1.0$  represent enhanced recruitment relative to reference
  - Values of  $RI < 1.0$  represent diminished recruitment relative to reference

### Murray cod

The recruitment index for Murray cod incorporates length frequency data only. Abundance is not included due to the low numbers of fish typically sampled. Murray cod recruitment is measured as the proportion of fish ranging 400–600 mm TL. This length range corresponds to individuals 3–6 years of age in the lower River Murray (Zampatti *et al.* 2014) and subsequently the age at sexual maturity (Rowland 1998), and thus represents recruitment to the adult population. Recruitment to YOY was also assessed, as indicated by the proportion of fish <200 mm TL and provides a useful measure of survival of recently spawned fish.

The reference value is the mean proportion of the population comprised of fish 400–600 mm TL and YOY <200 mm TL over baseline data collected from 2005–2014. These values are 20% and 7% for fish 400–600 mm TL and <200 mm TL, respectively.

Golden perch

Golden perch length-at-age is highly variable (Anderson *et al.* 1992), therefore the recruitment index for golden perch incorporates abundance and age-structure data (as derived from otolith microstructure analyses). Recruitment of golden perch is measured as the abundance of individuals classified at age 0+ (i.e. YOY) and age 1+. Detectability of 1+ fish is greater than for 0+ fish and provides a more reliable estimate of recruitment. The reference value is calculated as the mean abundance of the age 0+/1+ cohorts from baseline data collected from 2005–2014, where:

- $X$  = icon site abundance (fish.minute electrofishing<sup>-1</sup>.site<sup>-1</sup>),
- $r_{standard}$  = set proportion of the combined 0+/1+ cohort\* (\*Values of  $r_{standard}$  were calculated as the mean proportion of the population comprised of 0+ and 1+ individuals from 2005–2014.)
- Reference value (RV) =  $\text{mean}((X_{2005} * r_{standard}) + (X_{2006} * r_{standard}) + (X_{2007} * r_{standard}) + \dots + (X_{2014} * r_{standard}))$   $r_{standard} = 19\%$ , RV = 0.10

The annual recruitment index for golden perch is then calculated as:

- $X_{year}$  = year specific abundance (fish.minute electrofishing<sup>-1</sup>.site<sup>-1</sup>),
- $r_{year}$  = year specific proportion of YOY
- Annual recruitment value (AV) =  $X_{year} * r_{year}$
- Recruitment index (RI) =  $AV/RV$ 
  - Values of  $RI > 1.0$  represent enhanced recruitment relative to reference
  - Values of  $RI < 1.0$  represent diminished recruitment relative to reference


### 3. RESULTS

#### *Hydrology during study period*

Discharge in the lower River Murray (QSA) over the course of the study has been highly variable (Figure 3). From 2001 to 2010, the MDB experienced one of the most severe droughts in recorded history (van Dijk *et al.* 2013), and thus, through 2005–2010, annual sampling was associated with discharge that was below average and insufficient to inundate floodplains (MDBA 2011). At the other end of the scale were years characterised by widespread and persistent flooding in 2010/11 (peak  $\sim 93,000 \text{ ML.d}^{-1}$ , duration  $\sim 11$  months) and 2022/23 (peak  $\sim 185,000 \text{ ML.d}^{-1}$ , duration  $\sim 8$  months), and smaller overbank flooding events in 2012 (peak  $\sim 60,000 \text{ ML.d}^{-1}$ , duration  $\sim 5$  months) and 2016/17 (peak  $\sim 95,000 \text{ ML.d}^{-1}$ , duration  $\sim 3$  months). For much of the intervening period (2013–2022), discharge remained within-channel, but with distinct peaks in discharge ( $12,000\text{--}38,000 \text{ ML.d}^{-1}$ ), often in spring–early summer. Except for 2011, discharge has been  $<25,000 \text{ ML.d}^{-1}$  and within-channel at the time of sampling.

Since 2014, in addition to changes to water level associated with hydrology, the Chowilla Regulator has been operated on six occasions (Figure 3). This includes mid-level events in 2014 (19.14 m AHD), high level events in 2016 (19.78 m AHD), 2021 (19.59 m AHD) and 2022 (19.56 m AHD); and a low-level in-channel rise event in 2018 (18.54 m AHD) and 2015 (17.85 m AHD).

In 2023/24, peak discharge and low-level overbank flooding occurred in early September 2023 ( $\sim 53,000 \text{ ML.d}^{-1}$ ) before discharge steadily decreased through to December 2023 ( $\sim 7,000 \text{ ML.d}^{-1}$ ) before a further peak in January 2024 ( $\sim 22,000 \text{ ML.d}^{-1}$ ) (Figure 3). During condition monitoring in February/March 2024, discharge had again decreased and ranged from  $\sim 8,000\text{--}12,000 \text{ ML.d}^{-1}$  (Figure 3).



**Figure 3.** Mean daily flow (ML.d<sup>-1</sup>) in the River Murray at the South Australian Border (Site A42610010) January 2004–July 2024 (MDBA, unpublished data). Closed red circles indicate sampling events, the dotted line represents approximate bankfull discharge at Chowilla (~35,000 ML.d<sup>-1</sup>) and the blue dashes represent Chowilla Regulator operations.

### Catch Summary

From 2005–2024, a total of 334,827 fish from 17 species (12 native and 5 non-native) were captured over twenty annual surveys (Table 5, Appendix 1–20). In 2024, 6,684 fish were sampled from 14 species (10 native and 4 non-native) (Table 5). In 2024, the most abundant native species were bony herring (80%), Australian smelt (2.5%) and unspoked hardyhead (1.9%), whilst the remaining species collectively comprised 1.5% of the total catch (Table 5). Common carp were the most abundant non-native species and comprised ~12% of the total catch (Table 5). Three species of conservation significance were collected. Two of these, Murray cod and silver perch (*Bidyanus bidyanus*) are listed as ‘vulnerable’ and ‘critically endangered’ respectively, under the

*EPBC Act*. The third, freshwater catfish, is protected under the South Australian *Fisheries Management Act 2007*.

#### *Abundance of native fish*

Golden perch was the most abundant large-bodied native species sampled in all years, with standardised abundance typically highest following significant overbank flooding (e.g. 2010/11) and lowest in low flow years (Table 5). Contrary to this pattern, in 2024, golden perch abundance was among the lowest recorded throughout the study despite following a year of persistent high flows. Low abundances of Murray cod and silver perch were generally captured each year (including 2024) but exhibited peaks in abundance in 2020–2023 and 2011, respectively (Table 5). Freshwater catfish were captured in very low numbers (i.e. <5 individuals) in most years and spangled perch were only captured in 2011, 2014 and 2015.

Bony herring were most abundant in 2016, 2020 and 2021, whilst the abundance of the small-bodied Murray rainbowfish was highest in 2011 and carp gudgeon, unspotted hardyhead and flat-headed gudgeon (*Philypnodon grandiceps*) were most abundant in 2005 (Table 5). Australian smelt was most abundant in 2021, whilst dwarf flat-headed gudgeon were not captured each year and when present, were sampled in low numbers (Table 5).

#### *Abundance of non-native fish*

Common carp and goldfish were the most abundant non-native species in most years (Table 5). Standardised abundances of common carp were greatest in 2011, 2017, 2022 and 2023, whilst goldfish abundance was highest in 2011, 2017 and 2023 (Table 5). Redfin perch were not captured each year and when present, were typically sampled in low numbers (i.e. ≤70 individuals). In 2023, however, the abundance of redfin perch increased dramatically and was the highest recorded since monitoring commenced in 2005, but in 2024 abundance once again decreased with only two individuals sampled (Table 5).


**Table 5.** Total and standardised (fish.site<sup>-1</sup>) abundances of fish captured from condition monitoring sites sampled in the Chowilla Anabranch system and adjacent River Murray 2005–2024.

Species	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	Grand Total
Golden perch	69	75	112	94	174	114	802	286	230	148	143	99	139	112	68	46	61	62	117	36	<b>2,987</b>
( <i>Macquaria ambigua</i> )	(3.8)	(4.7)	(7.5)	(6.7)	(8.3)	(5.2)	(38.2)	(13.0)	(10.5)	(7.0)	(6.5)	(4.7)	(6.3)	(5.3)	(3.1)	(2.2)	(2.9)	(3.0)	(5.3)	(2.0)	
Murray cod	13	11	14	15	21	15	7	9	7	7	14	13	5	18	13	33	38	30	22	18	<b>323</b>
( <i>Maccullochella peelii</i> )	(0.7)	(0.7)	(0.97)	(1.1)	(1.0)	(0.7)	(0.3)	(0.4)	(0.3)	(0.3)	(0.6)	(0.6)	(0.2)	(0.9)	(0.6)	(1.6)	(1.8)	(1.4)	(1.0)	(1.0)	
Silver perch	5	5	1	14	8	20	30	6	7	5	14	7	4	9	5	0	5	7	1	3	<b>156</b>
( <i>Bidyanus bidyanus</i> )	(0.3)	(0.3)	(0.1)	(1.0)	(0.4)	(0.9)	(1.4)	(0.3)	(0.3)	(0.2)	(0.6)	(0.3)	(0.2)	(0.4)	(0.2)	(0)	(0.2)	(0.3)	(0.0)	(0.2)	
Freshwater catfish	-	-	1	-	3	2	8	20	15	6	4	1	2	5	2	3	2	2	4	1	<b>81</b>
( <i>Tandanus tandanus</i> )	-	-	(0.1)	-	(0.1)	(0.1)	(0.4)	(0.9)	(0.7)	(0.3)	(0.2)	(0.1)	(0.1)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.1)	
Bony herring	3849	6229	6251	7782	10,629	17,948	2521	4433	5508	5225	10,314	19,221	11,045	11,151	15,530	22,611	22,229	11,192	12,692	5,351	<b>211,711</b>
( <i>Nematalosa erebi</i> )	(213.8)	(389.3)	(416.7)	(555.9)	(506.1)	(815.8)	(114.6)	(201.5)	(250.4)	(248.8)	(468.8)	(915.3)	(502)	(531)	(705.9)	(1076.7)	(1058.5)	(533.0)	(576.9)	(297.3)	
Australian smelt	526	189	740	803	1067	589	484	132	215	151	1029	916	2169	2377	680	1164	2720	1857	1919	173	<b>19,900</b>
( <i>Retropinna semoni</i> )	(29.2)	(11.8)	(49.3)	(57.4)	(50.8)	(26.8)	(22.0)	(6.0)	(9.8)	(7.2)	(46.8)	(43.6)	(98.6)	(113.2)	(30.9)	(55.4)	(129.5)	(88.4)	(87.2)	(9.6)	
Murray rainbowfish	458	378	123	213	231	240	686	50	200	235	652	490	195	557	204	293	372	310	273	38	<b>6,199</b>
( <i>Melantaenia fluviatilis</i> )	(25.4)	(23.6)	(8.2)	(15.2)	(11.0)	(10.9)	(31.2)	(2.3)	(9.1)	(11.2)	(29.6)	(23.3)	(8.9)	(26.5)	(9.3)	(13.9)	(17.7)	(14.8)	(12.4)	(2.1)	
Flat-headed gudgeon	93	6	20	18	70	21	11	20	69	35	65	14	4	33	6	102	88	32	1	5	<b>713</b>
( <i>Philypnodon grandiceps</i> )	(5.2)	(0.4)	(1.3)	(1.3)	(3.3)	(1.0)	(0.5)	(0.9)	(3.1)	(1.7)	(3.0)	(0.7)	(0.2)	(1.6)	(0.3)	(4.9)	(4.2)	(1.5)	(0.0)	(0.3)	
Dwarf flat-headed gudgeon	2	-	-	11	2	6	-	-	-	-	3	4	-	-	1	1	-	-	-	-	<b>30</b>
( <i>Philypnodon macrostomus</i> )	(0.1)	-	-	(0.8)	(0.1)	(0.3)	-	-	-	-	(0.1)	(0.2)	-	-	(0.0)	(0.1)	-	-	-	-	
Unspecked hardyhead	2659	1602	1574	1786	2145	1687	455	26	84	89	656	2441	1687	1494	525	922	1168	676	96	127	<b>21,899</b>
( <i>Craterocephalus stercusmuscarum fulvus</i> )	(147.7)	(100.1)	(104.9)	(127.6)	(102.1)	(76.7)	(20.7)	(1.2)	(3.8)	(4.2)	(29.8)	(116.2)	(76.7)	(71.1)	(23.9)	(43.9)	(55.6)	(32.2)	(4.4)	(7.1)	
Carp gudgeon spp.	398	113	104	73	84	153	92	2	28	222	137	251	181	68	93	109	102	26	9	3	<b>2,248</b>
( <i>Hypseleotris</i> spp.)	(22.1)	(7.1)	(6.9)	(5.2)	(4.0)	(7)	(4.2)	(0.1)	(1.3)	(10.6)	(6.2)	(12.0)	(8.2)	(3.2)	(4.2)	(5.2)	(4.9)	(1.2)	(0.4)	(0.2)	
Common carp*	234	466	277	185	400	357	11602	2023	1218	590	730	339	5164	539	700	412	484	3041	20,812	833	<b>50,456</b>
( <i>Cyprinus carpio</i> )	(13.0)	(29.1)	(18.5)	(13.2)	(19.1)	(16.2)	(527.4)	(92.0)	(55.4)	(28.1)	(33.2)	(16.1)	(234.7)	(25.7)	(31.8)	(19.6)	(23)	(144.8)	(946.0)	(49.1)	
Gambusia*	200	61	125	60	107	490	647	12	40	65	126	300	398	160	27	103	331	171	804	10	<b>4,237</b>
( <i>Gambusia holbrooki</i> )	(11.1)	(3.8)	(8.3)	(4.3)	(5.1)	(22.3)	(29.4)	(0.5)	(1.8)	(3.1)	(5.7)	(14.3)	(18.1)	(7.6)	(1.2)	(4.9)	(15.8)	(8.1)	(36.5)	(0.6)	
Goldfish*	202	296	177	156	551	217	3945	385	55	171	299	331	2517	188	146	90	200	207	2,487	34	<b>12,654</b>
( <i>Carassius auratus</i> )	(11.2)	(18.5)	(11.8)	(11.1)	(26.2)	(9.9)	(179.3)	(17.5)	(2.5)	(8.1)	(13.6)	(15.8)	(114.4)	(9.0)	(6.6)	(4.3)	(9.5)	(9.9)	(113.0)	(1.9)	
Redfin perch*	-	-	9	3	7	8	5	3	-	-	3	1	27	13	21	36	66	70	957	2	<b>1,231</b>
( <i>Perca fluviatilis</i> )	-	-	(0.6)	(0.2)	(0.3)	(0.4)	(0.2)	(0.1)	-	-	(0.1)	(0.1)	(1.2)	(0.6)	(1.0)	(1.7)	(3.1)	(3.3)	(43.5)	(0.1)	
Oriental weatherloach*	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	<b>2</b>
( <i>Misgurnus anguillicaudatus</i> )	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	(0.1)	-	
Spangled perch^	-	-	-	-	-	-	1	-	-	1	1	-	-	-	-	-	-	-	-	-	<b>3</b>
( <i>Leipotheapon unicolor</i> )	-	-	-	-	-	-	(0.05)	-	-	(0.05)	(0.05)	-	-	-	-	-	-	-	-	-	
<b>Total species</b>	<b>13</b>	<b>12</b>	<b>14</b>	<b>14</b>	<b>15</b>	<b>15</b>	<b>15</b>	<b>14</b>	<b>13</b>	<b>14</b>	<b>16</b>	<b>15</b>	<b>14</b>	<b>14</b>	<b>15</b>	<b>14</b>	<b>14</b>	<b>14</b>	<b>14</b>	<b>14</b>	<b>16</b>
<b>Total number of sites</b>	<b>18</b>	<b>16</b>	<b>15</b>	<b>14</b>	<b>21</b>	<b>22</b>	<b>21</b>	<b>22</b>	<b>22</b>	<b>21</b>	<b>22</b>	<b>21</b>	<b>22</b>	<b>21</b>	<b>22</b>	<b>21</b>	<b>21</b>	<b>21</b>	<b>22</b>	<b>18</b>	
<b>Total number of fish</b>	<b>8,708</b>	<b>9,431</b>	<b>9,528</b>	<b>11,213</b>	<b>15,499</b>	<b>21,867</b>	<b>21,296</b>	<b>7,407</b>	<b>7,676</b>	<b>6,950</b>	<b>14,190</b>	<b>24,428</b>	<b>23,537</b>	<b>16,724</b>	<b>18,021</b>	<b>25,925</b>	<b>27,866</b>	<b>17,683</b>	<b>40,196</b>	<b>6,684</b>	<b>334,827</b>
<b>Standardised total abundance (fish.site<sup>-1</sup>)</b>	<b>483.7</b>	<b>589.4</b>	<b>635.2</b>	<b>800.9</b>	<b>738.0</b>	<b>934.0</b>	<b>969.7</b>	<b>336.7</b>	<b>348.9</b>	<b>330.9</b>	<b>644.9</b>	<b>1163.2</b>	<b>1069.9</b>	<b>796.4</b>	<b>858.1</b>	<b>1234.5</b>	<b>1326.9</b>	<b>842.05</b>	<b>1827.0</b>	<b>371.33</b>	

\*Denotes non-native species, ^ denotes native species captured outside its 'normal' distribution range.

### Temporal variation in fish abundance

Between 2005 and 2024, annual relative abundance of fish (all species combined) varied significantly among years (Figure 4;  $Pseudo-F_{19, 402} = 9.9199$ ,  $p < 0.001$ ). Between 2005 and 2011, relative abundance gradually increased before decreasing substantially in 2012–2014 (Figure 4). Abundance again increased from 2015–2017, before slightly decreasing in 2018 and 2019, but increased again in 2020 and 2021 (Figure 4). In 2022, abundance decreased to levels similar to 2018–2019, before increasing substantially in 2023 to the highest levels recorded since sampling commenced in 2005 (Figure 4). In 2024, abundance again decreased with CPUE similar to those observed from 2012–2014. As a proportion of the total catch, native fish dominated in all years except for 2011, 2017 and 2023, when sampling followed high level overbank floods and common carp and goldfish comprised the majority of the catch (Figure 4; Table 5).



**Figure 4.** Mean ( $\pm$  SE) catch-per-unit-effort (CPUE) (fish.min<sup>-1</sup>) of fish (all species combined) collected annually during standardised boat electrofishing surveys from 2005–2024 at 14–22 sites in the Chowilla Anabranh system and adjacent River Murray (dark grey = proportion native species, light grey = proportion of non-native species).

### *Spatio-temporal differences in fish assemblage structure*

Two-factor PERMANOVA exhibited significant differences among years and mesohabitats, and no significant interaction (Table 6). This indicated that fish assemblages varied among years and mesohabitats, and that change through time was consistent among mesohabitats.


Pairwise comparisons revealed significant differences (B-Y corrected  $\alpha = 0.02$ ) in fish assemblages among mesohabitats for all comparisons (Table 7). In addition, MDS and cluster analysis indicated six distinct groupings of fish assemblages by sampling years (Figure 5a). Namely, assemblages sampled in years following within-channel flows in 2005–2010, 2015–2016 and 2018–2022 were similar, and distinctly different from assemblages sampled in years following high flow and flooding in 2011, 2012–2014, 2017, 2023 and 2024 (Figure 5b). A similar pattern of temporal variability was observed for fish assemblages among all mesohabitats (Figure 6a–6d).

**Table 6.** PERMANOVA results comparing the relative abundances of fish between years and mesohabitats from 2005–2024, excluding 2011. Significant *P* values are highlighted in bold.


Factor	<i>df</i>	<i>Pseudo-F</i>	<i>P</i>
Year	18, 381	12.667	<b>0.001</b>
Mesohabitat	3, 381	23.996	<b>0.001</b>
Year x mesohabitat	54, 381	0.93612	0.763

**Table 7.** PERMANOVA pair-wise comparisons between fish assemblages among different mesohabitats in Chowilla from 2005-2024, excluding 2011. Significant *p* values are highlighted in bold (B-Y corrected  $\alpha = 0.02$ ).

Pairwise comparison		<i>t</i>	<i>p</i> value
Mesohabitat	Mesohabitat		
Fast	Slow	6.3194	<b>0.001</b>
Fast	Backwater	5.8224	<b>0.001</b>
Fast	River	3.9568	<b>0.001</b>
Slow	Backwater	3.3399	<b>0.001</b>
Slow	River	4.7809	<b>0.001</b>
Backwater	River	2.931	<b>0.001</b>



**Figure 5.** a) Dendrogram indicating fish assemblage clusters across sampling years from 2005–2024. b) Non-metric multi-dimensional scaling (MDS) plot of fish assemblages sampled from all years/sites combined.



**Figure 6.** Non-metric multi-dimensional scaling (MDS) plots of a) fast-flowing, b) slow-flowing, c) backwater and d) river mesohabitats sampled from all years/sites combined (excluding 2011).

Indicator species analysis demonstrated that, in low flow years (Group 1: 2005–2010, 2015–2016, 2019–2022), fish assemblages were characterised by the medium-bodied bony herring (Table 8). Conversely, for two high flow groupings, fish assemblages were characterised by the large-bodied native freshwater catfish (Group 2: 2012–2014) and the large-bodied native silver perch and golden perch, and non-native goldfish (Group 3: 2011). The assemblage in 2011 was also characterised by the small-bodied Murray rainbowfish (Table 8). Group 4 (2017) was characterised by the small-bodied carp gudgeon spp. and unspotted hardyhead, whilst group 5 (2023) was characterised by non-native common carp, redfin perch, gambusia and oriental weatherloach, but also the small-bodied native species, Australian smelt (Table 8). No species were significant indicators of fish assemblages of group 6 (2024).

**Table 8.** Indicator species analysis comparing the relative abundance of fish amongst years from 2005–2024, including 2011. (Year group 1 = 2005–2010, 2015–2016 and 2019–2022; Year group 2 = 2012–2014; Year Group 3 = 2011; Year Group 4 = 2017; Year Group 5 = 2023; and Year Group 6 = 2024). Significant indicators ( $\alpha = 0.05$ ) signifies that a species occurs in a higher relative abundance in a specific year group. Indicators that are not significant signify that a species was either sampled in similar numbers in more than one year (widespread across years) or captured in low abundances (uncommon).

Species	Year Group	Indicator value	p value
Flat-headed gudgeon	1	15.0	0.1034
Dwarf flat-headed gudgeon	1	8.2	0.0600
Bony Herring	1	19.0	<b>0.0282</b>
Murray cod	1	8.9	0.5827
Freshwater catfish	2	13.1	<b>0.0208</b>
Silver perch	3	23.3	<b>0.0006</b>
Golden perch	3	26.5	<b>0.0002</b>
Spangled perch	3	3.4	0.3377
Murray rainbowfish	3	26.2	<b>0.0002</b>
Goldfish	3	26.6	<b>0.0002</b>
Carp gudgeon spp.	4	27.2	<b>0.0002</b>
Unspecked hardyhead	4	28.8	<b>0.0002</b>
Redfin perch	5	65.1	<b>0.0002</b>
Australian smelt	5	23.0	<b>0.0002</b>
Gambusia	5	29.4	<b>0.0002</b>
Common carp	5	25.3	<b>0.0002</b>
Oriental weatherloach	5	9.1	<b>0.0100</b>

Fast-flowing mesohabitats were characterised by greater abundances of golden perch, Murray cod, silver perch, Australian smelt and freshwater catfish (Table 9). Main river channel mesohabitats were characterised by Murray rainbowfish, unspecked hardyhead and redfin perch, and backwater mesohabitats were characterised by goldfish. No species were significant indicators of slow-flowing mesohabitats (Table 9).

**Table 9.** Indicator species analysis comparing the relative abundance of fish in three of the four aquatic mesohabitats from 2005–2024, excluding 2011 (no species were found to be associated with slow-flowing mesohabitats). A significant difference ( $\alpha = 0.05$ ) indicates that a species occurs in a higher relative abundance in a specific mesohabitat. Values that are not significant indicate that a species was either sampled in similar numbers in more than one mesohabitat (widespread) or captured in low abundances (uncommon).


Species	Mesohabitat	P - value
Golden perch	Fast	<b>0.0008</b>
Murray cod	Fast	<b>0.0002</b>
Silver perch	Fast	<b>0.0014</b>
Australian smelt	Fast	<b>0.0002</b>
Bony Herring	Fast	0.0532
Freshwater catfish	Fast	<b>0.0188</b>
Murray rainbowfish	River	<b>0.0002</b>
Spangled perch	River	0.5157
Unspecked hardyhead	River	<b>0.0002</b>
Redfin perch	River	<b>0.0250</b>
Gambusia	River	0.1784
Carp gudgeon spp.	River	0.0620
Common carp	Backwater	0.2226
Goldfish	Backwater	<b>0.0056</b>
Dwarf flat-headed gudgeon	Backwater	0.3553
Flat-headed gudgeon	Slow	0.1494
Oriental weatherloach	Slow	1.0000

#### *Targeted Murray cod sampling*

In 2024, targeted sampling yielded a total of 41 Murray cod, with a catch-per-unit-effort of 0.08 fish.min<sup>-1</sup>, indicating a decrease in abundance from the period 2020–2022 (Table 10; Figure 7). Eighteen Murray cod were captured during standard condition monitoring in February/March 2024 and the remaining 23 were captured during targeted sampling in June/July 2024. Relative abundances of Murray cod were highest in Slaney Creek (Table 10).

**Table 10.** Total and standardised catch-per-unit-effort (CPUE) (fish.min<sup>-1</sup>) of Murray cod from targeted sampling and condition monitoring in the Chowilla Anabranh system and adjacent River Murray between 2014–2024.

Year	Targeted cod surveys								Standard condition monitoring ( <i>n</i> = 18 sites)	Total
	Chowilla Creek d/s bridge	Chowilla Creek u/s bridge	River Murray Main channel	Slaney Creek	Pipeclay Creek	Bank K	Swifty's Creek	Salt Creek		
2024	0	2 (0.07)	2 (0.03)	19 (0.28)	0	- (-)	- (-)	- (-)	18 (0.05)	41 (0.08)
2023	2 (0.11)	7 (0.27)	0	15 (0.26)	0	- (-)	- (-)	0	22 (0.05)	48 (0.08)
2022	4 (0.13)	14 (0.30)	6 (0.59)	24 (0.34)	2 (0.12)	4 (0.18)	2 (0.16)	8 (0.41)	30 (0.07)	94 (0.15)
2021	1 (0.03)	4 (0.13)	4 (0.14)	25 (0.34)	4 (0.17)	7 (0.49)	2 (0.21)	5 (0.34)	38 (0.09)	90 (0.14)
2020	7 (0.53)	16 (0.30)	3 (0.36)	27 (0.47)	- (-)	- (-)	- (-)	- (-)	33 (0.09)	86 (0.17)
2019	1 (0.05)	5 (0.12)	3 (0.26)	22 (0.29)	0	2 (0.03)	2 (0.13)	2 (0.11)	13 (0.03)	50 (0.08)
2018	8 (0.23)	2 (0.04)	1 (0.02)	27 (0.31)	1 (0.08)	1 (0.05)	0	2 (0.09)	18 (0.05)	60 (0.09)
2017	1 (0.06)	3 (0.12)	0	12 (0.18)	1 (0.09)	0	1 (0.09)	0	5 (0.01)	23 (0.04)
2016	5 (0.10)	6 (0.10)	0	28 (0.24)	2 (0.22)	1 (0.03)	2 (0.14)	4 (0.13)	12 (0.03)	60 (0.07)
2015	0	1 (0.03)	2 (0.08)	18 (0.24)	2 (0.25)	0	1 (0.08)	0	14 (0.03)	38 (0.06)
2014	0	3 (0.05)	0	28 (0.29)	- (-)	2 (0.04)	- (-)	2 (0.10)	5 (0.01)	40 (0.06)




**Figure 7.** Relative abundance (fish.min<sup>-1</sup>) of Murray cod from targeted Murray cod surveys and condition monitoring combined since 2014.

#### *Diversity and extent of fish species (Ecological Objective 10)*

##### Diversity

From 2005–2024, except for 2008, 2010 and 2012–2014, diversity scores in all mesohabitats have been at or above reference values (Figure 8a). The mean of mesohabitat diversity indices for each year was calculated to provide an overall icon site diversity score (Figure 8b). In most years, except 2012, the icon site diversity score was greater than or equivalent to the reference value, with 2022 having the highest diversity score out of all sampling years. Decreases in diversity were observed in 2023–2024, but still remained above reference (Figure 8b).




**Figure 8.** Diversity indices for a) fast-flowing, slow-flowing, backwater and main channel mesohabitats and b) the calculated Icon Site Diversity Index (DI), at the Chowilla Icon Site from 2005–2024.

### Extent

For large-bodied species, golden perch distribution has been relatively stable across years (EI ranged from 0.75–1.0), while distribution of the remaining species has been variable (Figure 9a). For silver perch, EI was below the reference, suggesting limited extent in 2007, 2009, 2017, 2018, 2020 and 2022–2024. Freshwater catfish had limited distribution in 2005–2010, 2016, 2017, 2019, 2021, 2022 and 2024, but stable or increased distribution in 2011–2015 and in 2018, 2020 and 2023 (Figure 9a). Murray cod distribution varied greatly from year to year, with peaks in 2011, 2012 and 2020–2023, but a decline to below the reference in 2024 (Figure 9a).


The distribution of most small- to medium-bodied species among mesohabitats was stable over most sampling years (2005–2011, 2014–2016 and 2018–2024) (Figure 9b). Decreases in distribution for carp gudgeon (2012–2013 and 2023–2024), flat-headed gudgeon (2014–2017, 2022 and 2023) and unspecked hardyhead (2012–2014 and 2024) were typically associated with sampling that followed periods of high within-channel flows and overbank flooding in 2011–2013, 2016/17 and 2021–2024 (Figure 9b).



**Figure 9.** Extent Index (EI) scores for a) large-bodied native species and b) small- to medium-bodied native species at the Chowilla Icon Site from 2005–2024. Black dashed line represents extent equal to the reference, green dashed line extent 25% greater than reference and red dashed line extent 25% lesser than reference.

### Recruitment of small- to medium-bodied native species

For small- to medium-bodied species, annual recruitment was evident but indices varied among sampling years (Figure 10). Unspecked hardyhead recruitment was highest in 2005–2010, 2016, 2018 and 2021, but reduced in 2012–2015, 2017, 2019–2020, 2022–2024 (Figure 10). High recruitment of Murray rainbowfish was evident in 2005–2006, 2014–2015 and 2019–2021, but limited in 2017–2018 and 2022–2024 (Figure 10). Australian smelt recruitment was variable, with high recruitment in 2005, 2007–2010 and 2015–2021 but reduced recruitment in 2024 (Figure 10). Bony herring recruitment was above reference in 2007–2010 and 2015–2024, but limited from 2012–2014 (Figure 10).




**Figure 10.** Recruitment Index (*RI*) values for unspecked hardyhead, Murray Rainbowfish, Australian smelt and bony herring from 2005–2024. Values for 2011 are not presented as sampling occurred at atypical time of year due to flooding. Dashed black line represents recruitment equal to the reference value and the dashed red line, recruitment 75% of the reference value.

### Recruitment of large-bodied native species

#### Murray cod


Recruitment indices for Murray cod ranging from 400–600 mm TL were greater than reference in 2005–2006, 2012–2014, 2017 and 2021–2022, similar to reference in 2008–2009 and lower than reference in 2007, 2010, 2015–2016, 2018–2020 and 2023–2024 (Figure 11a). For YOY Murray cod (<200 mm TL), recruitment was greater than reference in 2009–2010, 2013–2014, 2016 and 2018–2024, similar to reference in 2006, 2008 and 2015 and less than reference in 2005, 2012 and 2017 (Figure 11b).



**Figure 11.** Recruitment Index ( $R_I$ ) values for a) Murray cod ranging 400–600 mm TL and b) YOY Murray cod (<200 mm TL) from 2005–2024. Values for 2011 are not presented as sampling occurred at an atypical time of year due to flooding. Dashed black line represents recruitment equal to the reference value and the dashed red line, recruitment 75% of the reference value.

### Golden perch


Golden perch recruitment was highest in the 2007, 2011–2012 and 2023 when the recruitment index values exceeded the reference value (Figure 12). In the remaining years, including 2024, YOY were mostly absent and therefore recruitment was negligible (Figure 12).




**Figure 12.** Recruitment Index (RI) values for golden perch from 2005–2024. Dashed black line represents recruitment equal to the reference value and the dashed red line, recruitment 75% of the reference value.

### *Recruitment of non-native species*

Length frequency distributions for common carp and goldfish indicate annual recruitment of age 0+ fish for both species in most years (i.e. common carp and goldfish ~100 mm in length) (Figure 13 and 14). Nevertheless, for common carp, recruitment was temporally variable and strong cohorts were most evident in 2006, 2009–2012, 2014–2017, 2019, and 2022–2024 whilst for goldfish, strong 0+ cohorts were evident in 2006, 2008–2009, 2014–2017, 2019, 2022 and 2023 (Figure 13 & 14).



**Figure 13.** Length distribution of common carp captured at all sites sampled within Chowilla and the adjacent River Murray main channel from 2005–2024.



**Figure 14.** Length distribution of goldfish captured at all sites sampled within Chowilla and the adjacent River Murray main channel from 2005–2024.

## 4. DISCUSSION

Condition monitoring of fish assemblages at Chowilla from 2005–2024 indicates Ecological Objectives 10 and 11 (as defined in the Chowilla Environmental Water Management Plan) are being met. In 2024, some species exhibited decreased extent, abundance and recruitment; this may be the result of prolonged high flow through 2022–2024 and four sites not being sampled, but also maybe related to issues with sampling. In 2024, vessel breakdown necessitated the use of a hired electrofishing boat. While this vessel had the same electrofishing unit and system as the SARDI vessel that had been used to collect the majority of long-term data, it is possible that sampling efficiency may have varied between the vessels. As such, continued monitoring will provide greater insight on true long-term trends on species extent, abundance and recruitment.

### *Abundance and assemblage structure*

In 2024, 14 fish species were sampled from Chowilla and the adjacent River Murray main channel. The fish assemblage consisted of 10 native and 4 non-native species, with bony herring, common carp, Australian smelt and unspotted hardyhead the most abundant species. Nonetheless, overall fish abundance (species pooled) was generally low. The fish assemblage in 2024 was significantly different to all other years, but was most closely aligned to years of sampling that followed persistent high flows (e.g. 2012–2014).

Fish assemblages at Chowilla immediately following overbank flooding (e.g. 2011, 2017 and 2023) are often characterised by high abundance and dominance of newly recruited common carp. A substantial decline in overall abundance often occurs in the following year, which is driven by declines in abundance of common carp. This pattern was observed in 2024. Furthermore, native small-bodied generalist species such as Australian smelt, unspotted hardyhead and Murray rainbowfish are typically less abundant in years following large scale flooding events, as evident in 2023–2024 and 2012–2014 (Wilson *et al.* 2015). These flow-related patterns of abundance are commonly observed in the lower River Murray main channel (Bice *et al.* 2014). Causal mechanisms are likely associated with the influence of hydraulics on in-stream habitat (e.g. aquatic macrophyte cover) and key life history processes (e.g. survival of early life stages and recruitment). Several of these small-bodied generalist species are typically associated with aquatic macrophytes (Bice *et al.* 2014). During periods of regulated, low flow and benign hydraulics, such habitats proliferate in the creeks of Chowilla, and in the lower River Murray main channel. These habitats, however, are diminished during and immediately following periods of high flow/flooding (Bice *et al.* 2014).

In contrast to small-bodied generalist species, golden perch were most abundant in 2011–2013, post widespread flooding in the MDB, and least abundant in 2005–2010, 2016 and 2018–2022 in association with low River Murray flows. Golden perch are periodic strategists that spawn and recruit in association with increased discharge (Humphries *et al.* 1999; Mallen-Cooper and Stuart 2003; Zampatti and Leigh 2013a), and substantial increases in abundance in 2011–2013 were a result of enhanced recruitment in association with overbank flooding (Zampatti and Leigh 2013b). In 2024, however, despite large-scale flooding occurring in the MDB in 2022/23 and persistent high flows throughout the remainder of 2023, golden perch recruitment was limited and abundance was the lowest recorded since condition monitoring commenced in 2005. In part, this may be due to the electrofishing vessel used in the 2024 surveys and differing sampling efficiency to the standard SARDI vessel. Indeed, sampling in the main river channel for the Flow-MER project suggested abundances of golden perch were similar to recent years (SARDI unpublished data). Additionally, this work did sample low numbers of 0+ and 1+ golden perch, which were not detected in Chowilla, suggesting a level of recruitment occurred in the lower River Murray in the past two years.

Freshwater catfish were also most abundant in years following elevated within-channel flows and overbank flooding in 2011–2013, but have remained in low abundance, albeit present every year, from 2014–2024. High abundances in 2011–2013, followed by a decline, was evident across the lower River Murray (Ye *et al.* 2015). Abundance of silver perch also peaked in 2011, but this species has only been sampled in low numbers (<10) in recent years.

Murray cod were generally captured in low abundances (7–38 individuals.year<sup>-1</sup>) in the Chowilla condition monitoring surveys (February–March), but targeted surveys in May–June 2014–2024 have increased catches (23–94 individuals.year<sup>-1</sup>). Based on integrated data, relative abundance peaked in 2020–2022, and has since gradually decreased. Nonetheless, abundance in 2024 remains moderate–high relative to 2014–2019.

Common carp were the most abundant non-native species. Carp abundances were greatest following years of increased discharge and flooding in 2011, 2017 and 2023. Throughout the southern MDB, increased water levels and floodplain inundation (natural and engineered) lead to increases in carp abundance (King *et al.* 2003; Stuart and Jones 2006; Bice and Zampatti 2011). In 2024, common carp abundances were substantially lower than 2023 due to much lower recruitment in 2024.

### *Diversity and extent (Ecological Objective 10)*

Species diversity in each mesohabitat and across years was similar, with mesohabitat diversity exceeding the reference value in most years. Furthermore, the extent of most species has mostly been maintained across all mesohabitats and sampling years. In 2024, however, a decrease in diversity and extent was observed for several species. Nonetheless, this result may in part, be due to reduced sampling efficiency with the provisional electrofishing vessel and may not represent a true pattern.

Distinct species are characteristic of certain mesohabitats. Over the 20-year sampling period (2005–2024), Murray cod, golden perch, silver perch, Australian smelt and freshwater catfish have characterised fast-flowing mesohabitats. Since 2018, the extent of Murray cod distribution throughout the Chowilla region has mostly increased, which can be attributed to YOY fish being sampled at additional sites and in different mesohabitats (e.g. Pipeclay Creek and Isle of Mann backwater). However, a decrease in extent for this species was evident between 2023–2024. Historically, species such as Murray cod, golden perch and silver perch were abundant in flowing riverine environments of the lower Murray (Cadwallader 1978; Mallen-Cooper and Brand 2007). The general loss of lotic habitats from the main channel of the lower Murray has corresponded with decreases in abundance of these species, thus highlighting the importance of flowing water mesohabitats in Chowilla (Mallen-Cooper and Zampatti 2018).

### *Recruitment of native species (Ecological Objective 11)*

Recruitment indices for the small-bodied species Murray rainbowfish, unspotted hardyhead, Australian smelt and the medium-bodied bony herring indicate some level of recruitment in all years. Recruitment was below reference values for Australian smelt in 2012–2014 and 2024, for unspotted hardyhead in 2017, 2019–2020 and 2022–2023, and for Murray rainbowfish in 2017–2018 and 2022–2024. These small-bodied generalist species are widespread and abundant throughout the lower River Murray (Baumgartner *et al.* 2008; Davies *et al.* 2008; Zampatti *et al.* 2008; Bice *et al.* 2014) and have flexible spawning and recruitment strategies that are not reliant on elevated flow (Baumgartner *et al.* 2013). Indeed, in the Chowilla region, these species displayed higher rates of recruitment in low flow years and as seen between 2023–2024, limited recruitment following high flow periods.

The YOY recruitment index for Murray cod was variable with distinct peaks in YOY recruitment in 2016, 2018 and 2020–2021. In these years, YOY Murray cod were also present throughout the

main channel of the lower River Murray (Ye *et al.* 2020); nevertheless, a mechanism for increased YOY recruitment during these low flow years remains unresolved. In 2017 and 2019, Murray cod YOY recruitment decreased substantially. Decreased recruitment coincided with a basin scale blackwater event in 2016/17 and the operation of the Chowilla Regulator the preceding spawning season (i.e. September–November 2016 and 2018), which may have promoted hydraulic conditions less conducive to larval survival and subsequent recruitment (Fredberg and Zampatti 2018; Gibbs *et al.* 2020). In 2024, the recruitment index decreased from the previous year, but was still above the reference value, suggesting recruitment of Murray cod to YOY within Chowilla was maintained.

Since 2005, the recruitment index for Murray cod 400–600 mm TL has indicated annual recruitment to reproductive maturity, except for 2015. Enhanced recruitment to maturity was observed in 2005–2006, 2012–2014, 2017 and 2021–2022. Years with high recruitment to maturity are correlated with increases in the abundance of YOY fish 3–4 years prior, thus indicating successful survival of these fish (Fredberg and Zampatti 2017). From 2015–2020 and 2023–2024, the annual recruitment to reproductive maturity index has been below the reference value, with the exception of 2017. Variability in recruitment of Murray cod at Chowilla warrants further investigation; mixed-effects modelling that assesses recruitment strength in association with various factors (e.g. hydrology, regulator operation, hydraulics) may be a viable approach in the future (Tonkin *et al.* 2020).

From 2005–2024, golden perch recruitment was episodic, with the recruitment index indicating the most prominent recruitment in 2011, and to a lesser extent in 2007, 2012 and 2022–2023 in association with antecedent overbank flooding or elevated within-channel flows in the lower River Murray. This is consistent with contemporary models of the flow-related ecology of golden perch that suggest spawning and recruitment of golden perch is stimulated by increases in discharge contained within the river channel or overbank (Humphries *et al.* 1999; Mallen-Cooper and Stuart 2003). In the lower River Murray, prominent recruitment of golden perch generally occurs when spring-summer flows exceed  $\sim 15,000$  ML.d<sup>-1</sup> (Zampatti and Leigh 2013a). Given this, we would predict limited recruitment of golden perch in the Chowilla region from 2015–2021, as flows in this period have predominantly remained low, with the exception of overbank flooding in 2016/17, which was associated with a hypoxic blackwater event. In 2024, no recruitment was again observed within Chowilla for golden perch despite increased recruitment of the species in 2023 after large-scale flooding in late 2022. Nonetheless, there was evidence of limited recruitment to

YOY for this species in the lower River Murray main channel during the same period (SARDI/CSIRO unpublished data).

#### *Recruitment of non-native species*

Increased recruitment of YOY common carp and goldfish in the Chowilla region generally occurs in sampling years following increased discharge and water levels in the lower River Murray (e.g. in 2011, 2014, 2017 and 2022–2023). Increases in water level may enhance recruitment of common carp and goldfish by increasing spawning effort and/or the availability of appropriate spawning and nursery habitat. In addition to years following high flow, in 2019, elevated recruitment was evident for common carp and goldfish and could potentially be associated with the operation of the Chowilla regulator from August–December 2018 and subsequent localised floodplain inundation. In contrast, sampling years that follow within-channel flow and no floodplain inundation (e.g. 2010) are typically characterised by low recruitment and abundance of common carp. In 2024, YOY recruitment of common carp was evident. This was somewhat expected as a residual response to the large-scale flooding event that occurred in the MDB in 2022/23 and was also seen throughout the lower River Murray and associated anabranches during the same period (SARDI unpublished data).

## 5. CONCLUSIONS

Annual monitoring of fish assemblages at Chowilla from 2005–2024 indicates the diversity and spatial distribution of native fishes have largely been maintained and that ecological objectives 10 and 11 of the Chowilla Environmental Water Management Plan are being met. Declines were observed in some metrics but may have been associated with issues of sampling efficiency related to the use of an alternative electrofishing boat.

### *Future Research Needs*

Twenty years of annual condition monitoring at Chowilla provides valuable information on the population dynamics of freshwater fish at Chowilla and the lower River Murray. Some underlying causal mechanisms of observed responses remain speculative and require hypothesis-based research in conjunction with condition monitoring to further test and refine our conceptual understanding of fish ecology in the lower River Murray, particularly in relation to contemporary management interventions at Chowilla (such as operation of the Chowilla Regulator).

Some research questions in order of priority include:

- Investigating factors influencing recruitment variability of Murray cod at Chowilla using mixed-effects modelling (*sensu* Tonkin *et al* 2020).
- Investigating the duration, timing, and hydrological parameters of elevated within-channel flows in relation to golden perch recruitment in the Chowilla region.
- The influence of site-scale management (e.g. regulator operation) on recruitment and abundance of common carp at Chowilla.
- The importance of Chowilla as a recruitment source for Murray cod and common carp in the lower River Murray, including the movement of adult and juvenile fish.
- The movement and habitat use of native (e.g. Murray cod) and exotic fish (e.g. common carp and redfin perch) in the Chowilla system and adjacent River Murray in relation to natural flows and engineered floodplain inundation.
- Response of fish assemblages (diversity and abundance) to altered hydrodynamics at the mesohabitat scale.

- Impact of river-scale blackwater events on the ecology and population dynamics of large-bodied native fish (e.g. golden perch and Murray cod) in the Chowilla region and adjacent River Murray.

## REFERENCES

- Anderson, J. R., Morison, A. K. and Ray, D. J. (1992). Validation of the use of thin-sectioned otoliths for determining the age and growth of golden perch, *Macquaria ambigua* (Perciformes : Percichthyidae), in the lower Murray Darling Basin, Australia. *Australian Journal of Marine and Freshwater Research* **43**, 1103-1128.
- Anderson, M. J. (2001). A new method for non-parametric analysis of variance. *Austral Ecology* **26**, 32-46.
- Anderson, M. J. and Ter Braak, C. J. F. (2003). Permutation tests for multi-factorial analysis of variance. *Journal of Statistical Computation and Simulation* **73**, 85-113.
- Anderson, M. J., Gorley, R. N. and Clarke, K. R. (2008). PERMANOVA+ for PRIMER: Guide to software and statistical methods, PRIMER-E: Plymouth, UK.
- Baumgartner, L. J., Stuart, I. G. and Zampatti, B. P. (2008). Determining diel variation in fish assemblages downstream of three weirs in a regulated lowland river. *Journal of Fish Biology* **72**, 218–232.
- Baumgartner, L. J., Conallin, J., Wooden, I., Campbell, B., Gee, R., Robinson, W. A. and Mallen-Cooper, M. (2013). Using flow guilds of freshwater fish in an adaptive management framework to simplify environmental flow delivery for semi-arid riverine systems. *Fish and Fisheries* **15**, 410–427.
- Benjamini, Y. and Yekutieli, D. (2001). The control of false discovery rate under dependency. *Annals of Statistics* **29**, 1165-1188.
- Bice, C.M. and Zampatti, B.P. (2011). Engineered water level management facilitates recruitment of non-native common carp, *Cyprinus carpio*, in a regulated lowland river. *Ecological Engineering* **37**, 1901-1904.
- Bice, C. M., Leigh, S. J., Nicol, J. M. and Zampatti, B. P. (2013). Changes in hydraulic complexity in the lower River Murray main channel in relation to flow variability. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2013/000214-1. SARDI Research Report Series No. 709. 32pp.
- Bice, C.M., Gehrig, S.L., Zampatti, B.P., Nicol, J.M., Wilson, P., Leigh, S.L., and Marsland, K. (2014) Flow induced alterations to fish assemblages, habitat and fish-habitat associations in a regulated lowland river. *Hydrobiologia* **722**, 205-222.
- Bice, C.M., Gibbs, M.S., Kilsby, N.N., Mallen-Cooper, M. and Zampatti, B.P. (2017). Putting the “river” back into the lower River Murray: quantifying the hydraulic impact of river regulation to guide ecological restoration, Transactions of the Royal Society of South Australia, DOI: 10.1080/03721426.2017.1374909.
- Bray, J. R. and Curtis, J. T. (1957). An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs* **27**, 325-349.
- Cadwallader, P.L. (1978). Some causes of the decline in range and abundance of native fish in the Murray-Darling river system. *Proceedings of the Royal Society of Victoria* **90**, 211–224.

Davies, P. E., Harris, J. H., Hillman, T. J. and Walker, K. F. (2008). SRA Report 1: A Report on the Ecological Health of Rivers in the Murray–Darling Basin, 2004–2007. Prepared by the Independent Sustainable Rivers Audit Group for the Murray–Darling Basin Ministerial Council.

DWLBC (2006). Asset Environmental Management Plan: Chowilla Floodplain (excluding Lindsay – Wallpolla) Significant Environmental Asset. Department of Water, Land and Biodiversity Conservation.

Dufrene, M. and Legendre, P. (1997). Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs* **67**, 345-366.

Environment Australia (2001). A Directory of Important Wetlands in Australia, Third Edition. Environment Australia, Canberra.

Fredberg, J. and Zampatti, B.P. (2017). Chowilla Icon Site Fish Assemblage Condition Monitoring 2016. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication F2008/000907-8. SARDI Research Report Series No. 940. 58pp.

Fredberg, J. and Zampatti, B.P. (2018). Murray cod (*Maccullochella peelii*) movement during regulator operation and passage efficiency of Slaney and Pipeclay Creek fishways, Chowilla, 2016–17. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2017/000349-1. SARDI Research Report Series No. 961. 53pp.

Gibbs, M. S., C. Bice, J. Brookes, D. Furst, L. Gao, K. Joehnk, M. Marklund, J. Nicol, H. Pethybridge, S. Szarvas, T. Wallace, N. Welti and B. Zampatti (2020). Ecological connectivity of the River Murray: Managing ecological outcomes and water quality risks through integrated river management, Goyder Institute for Water Research Technical Report Series No. 20/03.

Humphries, P., King, A. J. and Koehn, J. D. (1999). Fish, flows and floodplains: links between freshwater fishes and their environment in the Murray-Darling River system, Australia. *Environmental Biology of Fishes* **56**, 129–151.

King, A. J., Humphries, P. and Lake, P.S. (2003). Fish recruitment on floodplains: the roles of patterns of flooding and life history characteristics. *Canadian Journal of Fish and Aquatic Science* **60**, 773–786.

King, A.J. (2004). Ontogenetic patterns of habitat use by fishes within the main channel of an Australian floodplain river. *Journal of Fish Biology* **65**, 1582–1603.

Leigh, S. J., Zampatti, B. P. and Nicol, J. M. (2010). Chowilla Icon Site – Fish Assemblage Condition Monitoring 2005 - 2010. South Australian Research and Development Institute (Aquatic Sciences), Adelaide, 32 pp. SARDI Publication No. F2008/000907-2. SARDI Research Report Series No.517.

Leigh, S.J. and Zampatti, B.P. (2012). Chowilla Icon Site Fish Assemblage Condition Monitoring 2011. South Australia Research and Development Institute (Aquatic Sciences), Adelaide, 31 pp. SARDI Publication No. F2008/000907-3. SARDI Research Report Series No. 597.

Lloyd, L. (1990). Fish communities. In: Chowilla Floodplain Biological Study (O'Malley, C. & Sheldon, F., eds), pp. 183–193. Nature Conservation Society of South Australia, Adelaide.

- Mallen-Cooper, M. and Stuart, I. (2003). Age, growth and non-flood recruitment of two potamodromous fishes in a large semi-arid/temperate river system. *River Research and Applications* **19**, 697–719.
- Mallen-Cooper, M. and Brand, D.A. (2007). Non-salmonids in a salmonid fishway: what do 50 years of data tell us about past and future fish passage? *Fisheries Management and Ecology* **14**, 319–332.
- Mallen-Cooper, M, Zampatti B.P. (2018). History, hydrology and hydraulics: Rethinking the ecological management of large rivers. *Ecohydrology*. 2018; e1965. <https://doi.org/10.1002/eco.1965>.
- McCune, B. and Mefford, M.J. (2005). PC-ORD Multivariate Analysis of Ecological Data, Version 5. (MjM Software Design: Glendon Beach, Oregon, USA).
- MDBC (2006). The Living Murray Foundation Report on the significant Ecological Assets Targeted in the First Step Decision. pp 329. Murray-Darling Basin Commission.
- MDBA (2011). The Living Murray Annual Environmental Watering Plan 2011-2012. MDBA publication no. 170/11. Murray-Darling Basin Authority, Canberra.
- MDBA (2012). Chowilla Floodplain: Environmental Water Management Plan 2012 MDBA Publication No. 220/11. Murray-Darling Basin Authority, Canberra.
- MDBA (2016). The Living Murray Annual Environmental Watering Plan 2014-2015. MDBA publication no. 25/15. Murray-Darling Basin Authority, Canberra.
- Milton, D.A. and Arthington, A.H. (1985). Reproductive strategy and growth of Australian smelt, *Retropinna semoni* (Weber) (Pisces: Retropinnidae), and the olive perchlet, *Ambassis nigripinnis* (De Vis) (Pisces: Ambassidae), in Brisbane, south-eastern Queensland. *Australian Journal of Marine and Freshwater Research* **36**, 329–341.
- O'Malley, C. and Sheldon, F. (1990). Chowilla Floodplain Biological Study. pp 224. Nature Conservation Society of South Australia, Adelaide.
- Pierce, B. E. (1990). Chowilla Fisheries Investigations. South Australian Department of Fisheries, Adelaide.
- Robinson, W. A. (2013). The Living Murray: Towards assessing whole of Icon Site Condition. Report to the Murray-Darling Basin Authority, July 2013.
- Rowland, S. J. (1998). Aspects of the reproductive biology of Murray cod, *Macullochella peelii peelii*. *Proceedings of the Linnean Society of New South Wales* **120**: 147-162.
- Sharley, T. and Huggan, C. (1995). Chowilla Resource Management Plan. Murray-Darling Basin Commission, Canberra.
- Sheldon, F. and Lloyd, L. (1990) Physical limnology and aquatic habitats. In: Chowilla Floodplain Biological Study (O'Malley, C. and Sheldon, F., eds) pp 121-135. Nature Conservation Society of South Australia, Adelaide.

Stace, P. and Greenwood, A. (2004). Chowilla Anabranch System Surface Water Information Summary. Pg 88. Department of Water, Land and Biodiversity Conservation.

Stuart, I.G. and Jones, M. (2006). Large, regulated forest floodplain is an ideal recruitment zone for non-native carp (*Cyprinus carpio* L.). *Marine and Freshwater Research* **57**, 333–347.

Thiem, J. D., Wooden, I. J., Baumgartner, L. J., Butler, G. L., Forbes, J. P. and Conallin, J. (2017). Recovery from a fish kill in a semi-arid Australian river: Can stocking augment natural recruitment processes? *Austral Ecology* **42**, 218–226.

Tonkin, Z., J. Yen, J. Lyon, A. Kitchingman, J. D. Koehn, W. M. Koster, J. Lieschke, S. Raymond, J. Sharley, I. Stuart and C. Todd (2020). Linking flow attributes to recruitment to inform water management for an Australian freshwater fish with an equilibrium life-history strategy. *Science of The Total Environment*.

Van Dijk, A.I.J.M., Beck, H.E., Crosbie, R.S., de Jeu, R.A.M., Liu, G.M., Podger, B., Timbal, B. and Viney, N.R. (2013). The millennium drought in southeast Australia (2001–2009): Natural and human causes and implications for water resources, ecosystems, economy, and society. *Water Resources Research* **49**, 1040–1057.

Walker, K.F. (2006). Serial weirs, cumulative effects: the Lower River Murray, Australia. In 'Ecology of Desert Rivers'. (Ed. R Kingsford) pp.248–279. (Cambridge University Press).

Webb, J.A., Stewardson, M.J. and Koster, W.M. (2010). Detecting ecological responses to flow variation using Bayesian hierarchical models. *Freshwater Biology* **55**, 108–126.

Wilson, P., Leigh, S., Bice, C. and Zampatti, B. (2012). *Chowilla Icon Site Fish Assemblage Condition Monitoring 2012*. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2008/000907-4. SARDI Research Report Series No. 674. 48pp.

Wilson, P., Zampatti, B. P., Leigh, S. J. and Bice, C. M. (2014). *Chowilla Icon Site Fish Assemblage Condition Monitoring 2013*. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2008/000907-5. SARDI Research Report Series No. 785. 50pp.

Wilson, P.J., Zampatti, B.P. and Leigh, S.J. (2015). *Chowilla Icon Site Fish Assemblage Condition Monitoring 2014*. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2008/000907-6. SARDI Research Report Series No. 839. 54pp.

Wilson, P.J., Fredberg, J. and Zampatti, B.P. (2016). *Chowilla Icon Site Fish Assemblage Condition Monitoring 2015*. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2008/000907-7. SARDI Research Report Series No. 925. 58pp.

Winemiller, K. O. and K. A. Rose (1992). Patterns of life-history diversification in North American fishes: implications for population regulation. *Canadian Journal of Fisheries and Aquatic Sciences* **49**: 2196-2218.

Ye, Q., Bucater, L., Zampatti, B. P., Bice, C. M., Wilson, P. J., Sutor, L., Wegener, I. K., Short, D. A. and Fler, D. (2015). Population dynamics and status of freshwater catfish (*Tandanus tandanus*) in the lower River Murray, South Australia. Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2014/000903-1. SARDI Research Report Series No 841. 51pp.

Ye, Q., Giatas, G., Brookes, J., Furst, D., Gibbs, M., Oliver, R., Shiel, R., Zampatti, B., Aldridge, K., Bucater, L., Busch, B., Hipsey, M., Lorenz, Z., Maas, R., and Woodhead, J. (2020). Commonwealth Environmental Water Office Long-Term Intervention Monitoring Project 2014–2019: Lower Murray River Technical Report. A report prepared for the Commonwealth Environmental Water Office by the South Australian Research and Development Institute, Aquatic Sciences.

Zampatti, B. P., Leigh, S. J and Nicol, J M. (2008). Chowilla Icon site – fish assemblage condition monitoring 2005 – 2008. South Australian research and development institute (Aquatic Sciences), Adelaide, 38 pp. SARDI Publication No. F2008/000907-1. SARDI Research Report Series Number 319.

Zampatti, B.P., Leigh, S.J. and Nicol, J.M. (2011). Fish and Aquatic macrophyte communities in the Chowilla Anabranch system, South Australia: *A report on investigations from 2004 – 2007*. South Australian Research and Development Institute (Aquatic Sciences), Adelaide, 180 pp. SARDI Aquatic Sciences Publication Number: F2010/000719-1. SARDI research report series number: 525.

Zampatti, B.P. and Leigh, S.J. (2013a). Within-channel flows promote spawning and recruitment of golden perch, *Macquaria ambigua ambigua* – implications for environmental flow management in the River Murray, Australia. *Marine and Freshwater Research* **64**, 618-630.

Zampatti, B.P. and Leigh, S.J. (2013b). Effects of flooding on recruitment and abundance of golden perch (*Macquaria ambigua ambigua*) in the lower River Murray. *Ecological Management and Restoration* **14**, 135-143.

Zampatti, B.P., Bice, C.M., Wilson, P.J., and Ye, Q. (2014). Population dynamics of Murray cod (*Maccullochella peelii*) in the South Australian reaches of the River Murray: a synthesis of data from 2002–2013. South Australian Research and Development Institute (Aquatic Sciences), Adelaide, 42 pp. SARDI Aquatic Sciences Publication Number: F2014/000089-1. SARDI research report series number: 761.

Zampatti, B., Leigh, S. J., Bice, C. M. and Rogers, P. J. (2018). Multiscale movements of golden perch (Percichthyidae: *Macquaria ambigua*) in the River Murray, Australia. *Austral Ecology* **43**: 763-774.

**APPENDIX 1.** Total number of species captured at each site in 2005.

2005	Site Number																		Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	18	22	
Golden perch	7	7	2	5	1	4	9	10	0	6	1	3	2	4	2	3	3	0	69
Murray cod	0	2	0	1	2	1	2	4	0	0	0	0	0	0	0		1	0	13
Silver perch	1	0	0	0	1	0	0	0	0	0	0	0	2	0	1			0	5
Bony herring	503	75	183	27	390	217	433	93	61	184	164	148	124	325	104	727	90	1	3849
Australian smelt	35	9	36	5	166	18	50	29	29	19	26	20	48	15	1	17	3	0	526
Murray rainbowfish	15	16	10	17	46	10	27	18	0	13	6	7	124	83	17	4	45	0	458
Flathead gudgeon	2	1	10	0	2	2	4	3	0	3	0	4	10	12	0	2		38	93
Dwarf flathead gudgeon	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0			0	2
Unspecked hardyhead	131	8	166	20	57	307	48	18	34	79	23	34	413	712	300	101	157	51	2659
Carp gudgeon spp	3	5	24	5	4	21	14	4	3	7	0	8	24	23	0	97	6	150	398
Freshwater catfish	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0	0
Common carp	13	10	15	17	19	8	7	11	3	14	12	13	6	16	36	19	9	6	234
Gambusia	0	3	26	53	10	9	1	2	3	8	0	11	3	6	4	16	3	42	200
Goldfish	4	1	27	1	0	8	1	1	40	17	19	28	7	1	4	16	1	26	202
Redfin perch	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0	0
<b>Total species</b>	<b>10</b>	<b>11</b>	<b>10</b>	<b>10</b>	<b>11</b>	<b>11</b>	<b>11</b>	<b>11</b>	<b>8</b>	<b>10</b>	<b>7</b>	<b>10</b>	<b>12</b>	<b>10</b>	<b>9</b>	<b>10</b>	<b>10</b>	<b>7</b>	<b>13</b>
<b>Total fish/site</b>	<b>714</b>	<b>137</b>	<b>499</b>	<b>151</b>	<b>698</b>	<b>605</b>	<b>596</b>	<b>193</b>	<b>174</b>	<b>350</b>	<b>251</b>	<b>276</b>	<b>764</b>	<b>1197</b>	<b>469</b>	<b>1002</b>	<b>318</b>	<b>314</b>	<b>8708</b>

**APPENDIX 2.** Total number of species captured at each site in 2006.

2006 Species	Site Number																Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	
Golden perch	7	14	1	10	4	4	6	6	0	1	0	0	6	13	2	1	75
Murray cod	0	3	0	2	2	0	1	3	0	0	0	0	0	0	0	0	11
Silver perch	1	0	1	0	1	0	0	0	0	0	0	0	2	0	0	0	5
Bony herring	835	147	889	98	183	84	851	85	104	209	184	216	695	545	138	966	6229
Australian smelt	5	9	5	1	74	15	22	12	1	6	3	1	27	7	0	1	189
Murray rainbowfish	18	21	4	12	73	23	81	21	0	3	5	5	40	38	5	29	378
Flathead gudgeon	0	0	1	1	0	0	1	0	0	1	0	0	1	0	0	1	6
Dwarf flathead gudgeon	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Unspecked hardyhead	53	53	124	38	113	444	118	28	93	76	16	26	227	119	10	64	1602
Carp gudgeon spp	16	6	11	2	3	17	14	1	1	9	5	1	3	9	2	13	113
Freshwater catfish	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Common carp	13	6	57	9	24	23	19	11	13	47	52	48	25	21	47	51	466
Gambusia	3	1	1	23	0	5	6	4	0	5	0	7	5	0	1	0	61
Goldfish	17	1	13	3	0	17	3	0	15	27	40	24	27	64	8	37	296
Redfin perch	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<b>Total species</b>	<b>10</b>	<b>10</b>	<b>11</b>	<b>11</b>	<b>9</b>	<b>9</b>	<b>11</b>	<b>9</b>	<b>6</b>	<b>9</b>	<b>7</b>	<b>8</b>	<b>11</b>	<b>8</b>	<b>8</b>	<b>9</b>	<b>12</b>
<b>Total fish/site</b>	<b>968</b>	<b>261</b>	<b>1107</b>	<b>199</b>	<b>477</b>	<b>632</b>	<b>1122</b>	<b>171</b>	<b>227</b>	<b>384</b>	<b>305</b>	<b>328</b>	<b>1058</b>	<b>816</b>	<b>213</b>	<b>1163</b>	<b>9431</b>

**APPENDIX 3.** Total number of species captured at each site in 2007.

2007	Site Number															Grand Total
	1	2	3	4	5	6	7	8	10	11	12	13	14	15	16	
Golden perch	5	8	17	7	4	9	18	9	5	7	5	8	6	2	2	112
Murray cod	0	3	0	1	1	0	1	7	0	0	0	1	0	0	0	14
Silver perch	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Bony herring	201	145	170	132	2104	274	935	461	237	170	87	183	90	51	1010	6251
Australian smelt	7	38	5	11	142	92	319	44	12	0	0	55	9	0	6	740
Murray rainbowfish	6	13	13	15	14	3	26	6	6	0	1	8	7	2	3	123
Flathead gudgeon	0	0	4	2	0	1	0	0	2	1	1	3	3	2	1	20
Dwarf flathead gudgeon	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Unspecked hardyhead	13	28	15	89	100	145	353	17	34	5	5	298	158	215	99	1574
Carp gudgeon spp	6	3	2	5	0	16	50	0	7	1	0	5	2	0	7	104
Freshwater catfish	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1
Common carp	20	10	24	9	23	11	12	23	16	26	10	21	13	42	17	277
Gambusia	1	2	3	56	2	30	7	0	8	2	4	1	0	2	7	125
Goldfish	3	2	10	3	1	6	1	1	35	20	16	1	5	12	61	177
Redfin perch	0	0	0	0	0	0	0	0	0	0	0	1	4	2	2	9
<b>Total species</b>	<b>10</b>	<b>10</b>	<b>10</b>	<b>12</b>	<b>9</b>	<b>10</b>	<b>10</b>	<b>8</b>	<b>10</b>	<b>8</b>	<b>8</b>	<b>12</b>	<b>10</b>	<b>9</b>	<b>11</b>	<b>14</b>
<b>Total fish/site</b>	<b>263</b>	<b>252</b>	<b>263</b>	<b>331</b>	<b>2391</b>	<b>587</b>	<b>1722</b>	<b>568</b>	<b>362</b>	<b>232</b>	<b>129</b>	<b>585</b>	<b>297</b>	<b>330</b>	<b>1215</b>	<b>9528</b>

**APPENDIX 4.** Total number of species captured at each site in 2008.

2008 Species	Site Number														Grand Total
	1	2	3	4	5	6	7	8	9	10	11	13	14	19	
Golden perch	9	7	3	3	4	8	22	3	0	3	5	3	10	14	94
Murray cod	0	1	0	0	1	0	3	2	0	0	0	0	0	8	15
Silver perch	1	0	0	0	6	0	1	2	0	0	0	0	0	4	14
Bony herring	193	252	391	311	2573	297	1439	783	231	172	246	192	465	237	7782
Australian smelt	16	61	39	20	274	26	116	63	19	33	9	15	25	87	803
Murray rainbowfish	14	30	0	17	42	5	32	12	0	4	4	15	13	25	213
Flathead gudgeon	4	0	4	1	0	1	0	0	0	3	1	1	3	0	18
Dwarf flathead gudgeon	1	1	2	0	0	1	0	0	0	0	0	1	5	0	11
Unspecked hardyhead	40	95	53	274	220	252	423	23	23	56	7	103	164	53	1786
Carp gudgeon spp	8	4	3	6	5	5	16	1	1	10	0	2	11	1	73
Freshwater catfish	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Common carp	18	12	8	16	29	8	11	24	5	8	14	3	23	6	185
Gambusia	4	2	3	12	1	16	1	1	0	5	2	3	2	8	60
Goldfish	8	2	21	1	0	0	3	7	49	24	25	2	3	11	156
Redfin perch	0	0	0	0	0	0	1	0	0	0	0	0	2	0	3
<b>Total species</b>	<b>12</b>	<b>11</b>	<b>10</b>	<b>10</b>	<b>10</b>	<b>10</b>	<b>12</b>	<b>11</b>	<b>6</b>	<b>10</b>	<b>9</b>	<b>11</b>	<b>12</b>	<b>11</b>	<b>14</b>
<b>Total fish/site</b>	<b>316</b>	<b>467</b>	<b>527</b>	<b>661</b>	<b>3155</b>	<b>619</b>	<b>2068</b>	<b>921</b>	<b>328</b>	<b>318</b>	<b>313</b>	<b>340</b>	<b>726</b>	<b>454</b>	<b>11213</b>

**APPENDIX 5.** Total number of species captured at each site in 2009.

2009 Species	Site Number																					Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
Golden perch	8	17	4	13	5	10	20	14	0	3	3	5	9	11	3	1	7	11	11	9	10	174
Murray cod	0	4	0	0	0	0	3	1	0	0	0	0	2	0	0	0	0	0	5	0	6	21
Silver perch	0	0	0	0	2	1	2	0	0	0	0	0	0	0	0	0	0	0	2	0	1	8
Bony herring	474	817	947	276	808	970	1615	311	459	395	136	103	399	321	169	407	415	260	291	506	550	10629
Australian smelt	3	206	60	13	223	101	220	68	8	21	1	4	6	12	0	3	6	38	44	8	22	1068
Murray rainbowfish	4	10	0	36	15	15	21	1	0	6	1	2	8	36	5	0	3	34	15	3	16	231
Flathead gudgeon	3	1	9	0	1	0	0	0	2	7	2	2	0	15	2	11	4	3	5	0	3	70
Dwarf flathead gudgeon	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1
Unspecked hardyhead	106	105	22	342	95	258	76	47	93	100	10	22	87	209	75	152	35	138	43	64	66	2145
Carp gudgeon spp	11	0	3	2	7	2	5	0	2	4	0	1	0	18	2	5	1	7	5	3	6	84
Freshwater catfish	0	0	0	1	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	3
Common carp	23	17	22	14	11	7	6	16	75	7	10	23	11	16	19	16	23	31	9	23	21	400
Gambusia	1	2	0	17	2	14	0	2	3	9	2	1	1	12	4	9	4	6	7	7	4	107
Goldfish	40	5	41	2	4	10	0	1	69	17	24	17	45	28	61	57	34	9	7	42	38	551
Redfin perch	2	0	0	0	0	0	0	0	0	1	0	0	0	2	0	2	0	0	0	0	0	7
<b>Total species</b>	<b>11</b>	<b>10</b>	<b>8</b>	<b>10</b>	<b>12</b>	<b>10</b>	<b>9</b>	<b>9</b>	<b>8</b>	<b>10</b>	<b>9</b>	<b>10</b>	<b>9</b>	<b>12</b>	<b>9</b>	<b>10</b>	<b>10</b>	<b>10</b>	<b>12</b>	<b>10</b>	<b>12</b>	<b>15</b>
<b>Total fish/site</b>	<b>675</b>	<b>1184</b>	<b>1108</b>	<b>716</b>	<b>1174</b>	<b>1388</b>	<b>1968</b>	<b>461</b>	<b>711</b>	<b>570</b>	<b>189</b>	<b>180</b>	<b>568</b>	<b>681</b>	<b>340</b>	<b>663</b>	<b>532</b>	<b>537</b>	<b>444</b>	<b>666</b>	<b>743</b>	<b>15499</b>

**APPENDIX 6.** Total number of species captured at each site in 2010.

Species	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	9	11	4	4	4	14	14	7	3	1	1	1	6	17	5	0	1	3	7	0	2	0	114
Murray cod	0	4	1	1	0	0	3	2	0	0	0	0	0	0	0	0	0	0	4	0	0	0	15
Silver perch	0	9	1	0	2	0	0	4	0	0	0	0	1	0	1	0	0	0	2	0	0	0	20
Bony herring	940	274	781	177	1805	516	605	526	970	813	517	175	846	1334	820	5251	600	352	360	193	83	10	17947
Australian smelt	31	7	41	6	204	28	38	6	21	36	1	0	26	29	27	13	23	41	11	0	0	0	589
Murray rainbowfish	10	5	3	31	14	11	27	0	2	4	0	1	9	28	19	7	3	61	3	0	1	1	240
Flathead gudgeon	0	0	0	1	0	0	1	0	1	3	2	1	1	0	0	1	1	2	1	0	0	6	21
Dwarf flathead gudgeon	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	4	6
Unspecked hardyhead	38	14	34	72	15	20	16	1	26	38	18	2	124	706	101	52	32	86	21	2	3	267	1688
Carp gudgeon spp	10	2	5	2	2	4	4	1	8	8	5	0	6	8	1	1	2	11	1	0	0	72	153
Freshwater catfish	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	2
Common carp	41	22	30	13	11	12	8	9	35	12	15	11	43	15	21	14	5	15	3	16	4	2	357
Gambusia	8	2	12	25	2	5	3	1	10	34	23	4	6	22	39	35	7	44	17	2	2	187	490
Goldfish	42	5	6	3	3	5	0	3	14	2	9	8	23	19	16	11	14	6	2	4	2	20	217
Redfin perch	0	0	0	0	0	0	0	0	1	0	0	0	0	4	0	0	0	2	0	1	0	0	8
<b>Total species</b>	<b>9</b>	<b>11</b>	<b>11</b>	<b>12</b>	<b>10</b>	<b>9</b>	<b>10</b>	<b>10</b>	<b>12</b>	<b>10</b>	<b>9</b>	<b>8</b>	<b>11</b>	<b>11</b>	<b>10</b>	<b>9</b>	<b>10</b>	<b>12</b>	<b>12</b>	<b>6</b>	<b>7</b>	<b>9</b>	<b>15</b>
<b>Total fish/site</b>	<b>1129</b>	<b>355</b>	<b>918</b>	<b>336</b>	<b>2062</b>	<b>615</b>	<b>719</b>	<b>560</b>	<b>1092</b>	<b>951</b>	<b>591</b>	<b>203</b>	<b>1091</b>	<b>2183</b>	<b>1050</b>	<b>5385</b>	<b>688</b>	<b>624</b>	<b>432</b>	<b>218</b>	<b>97</b>	<b>569</b>	<b>21867</b>

**APPENDIX 7.** Total number of species captured at each site in 2011.

Species	2011																				Grand Total	
	Site Number																					
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	18	19	20	21	22	
Golden perch	50	30	22	25	54	39	41	20	16	37	46	61	80	26	11	10	98	39	49	46	2	802
Murray cod	0	1	1	0	0	0	1	0	0	0	0	1	3	0	0	0	0	0	0	0	0	7
Silver perch	2	7	2	0	2	1	0	0	0	1	1	0	1	3	0	0	7	3	0	0	0	30
Bony herring	62	31	76	7	70	74	209	59	171	61	41	213	142	161	65	644	226	83	59	54	13	2521
Australian smelt	14	72	75	2	4	37	50	7	6	0	44	47	10	28	0	45	1	1	18	22	1	484
Murray rainbowfish	69	22	15	35	26	44	16	9	26	55	44	22	52	93	18	8	47	27	40	14	4	686
Flathead gudgeon	1	1	2	0	0	0	0	0	0	0	1	0	0	0	1	1	3	0	1	0	0	11
Unspecked hardyhead	33	5	15	0	4	10	12	6	31	3	11	2	26	16	113	31	13	0	0	5	119	455
Carp gudgeon spp	1	1	8	0	0	0	0	1	8	1	2	1	2	0	0	13	3	1	5	0	45	92
Freshwater catfish	0	0	0	0	1	0	0	0	1	0	0	0	1	2	0	0	0	0	0	0	3	8
Common carp	536	587	225	1228	268	389	220	154	443	464	622	700	195	578	848	394	1052	613	806	1088	192	11602
Gambusia	13	22	5	23	70	8	34	4	8	19	7	8	2	9	2	48	21	1	47	9	287	647
Goldfish	210	56	70	310	46	112	27	56	116	155	545	248	77	158	33	408	103	364	344	493	14	3945
Redfin perch	2	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	2	0	0	0	5
Spangled perch	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1
<b>Total species</b>	<b>12</b>	<b>12</b>	<b>12</b>	<b>7</b>	<b>10</b>	<b>9</b>	<b>9</b>	<b>9</b>	<b>10</b>	<b>9</b>	<b>11</b>	<b>10</b>	<b>12</b>	<b>11</b>	<b>8</b>	<b>10</b>	<b>11</b>	<b>10</b>	<b>9</b>	<b>9</b>	<b>10</b>	<b>15</b>
<b>Total fish/site</b>	<b>993</b>	<b>835</b>	<b>516</b>	<b>1630</b>	<b>545</b>	<b>714</b>	<b>610</b>	<b>316</b>	<b>826</b>	<b>796</b>	<b>1364</b>	<b>1303</b>	<b>591</b>	<b>1075</b>	<b>1091</b>	<b>1602</b>	<b>1574</b>	<b>1134</b>	<b>1369</b>	<b>1732</b>	<b>680</b>	<b>21296</b>

**APPENDIX 8.** Total number of species captured at each site in 2012.

2012 Species	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	11	6	18	25	8	25	15	15	1	21	23	4	13	18	7	11	5	11	17	10	20	2	286
Murray cod	0	0	0	0	0	1	3	3	0	0	0	0	1	0	0	0	0	0	0	0	1	0	9
Silver perch	0	0	1	0	1	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	6
Bony herring	701	29	176	43	293	211	338	206	134	165	102	100	196	93	277	440	99	567	94	47	55	67	4433
Australian smelt	2	1	2	0	14	9	77	2	4	2	0	0	1	1	0	0	0	7	6	3	0	1	132
Murray rainbowfish	0	1	0	4	0	4	14	0	0	0	0	0	6	3	1	1	1	4	8	1	0	2	50
Flathead gudgeon	0	0	3	0	0	0	1	0	0	0	1	0	1	0	2	0	6	2	2	0	2	0	20
Unspecked hardyhead	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	23	26
Carp gudgeon spp	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2
Freshwater catfish	0	2	0	2	5	1	0	1	0	1	0	0	0	4	1	0	0	0	3	0	0	0	20
Common carp	94	30	99	272	18	100	16	144	28	158	54	58	155	72	53	165	78	91	56	72	177	33	2023
Gambusia	0	0	1	2	0	0	0	0	0	0	0	0	0	2	0	1	0	4	0	0	0	2	12
Goldfish	3	10	20	21	7	34	9	17	2	44	20	14	22	109	1	19	6	0	13	10	2	2	385
Redfin perch	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	3
<b>Total species</b>	<b>5</b>	<b>7</b>	<b>8</b>	<b>8</b>	<b>7</b>	<b>8</b>	<b>10</b>	<b>7</b>	<b>5</b>	<b>6</b>	<b>5</b>	<b>4</b>	<b>10</b>	<b>9</b>	<b>7</b>	<b>7</b>	<b>7</b>	<b>8</b>	<b>8</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>14</b>
<b>Total fish/site</b>	<b>811</b>	<b>79</b>	<b>320</b>	<b>371</b>	<b>346</b>	<b>385</b>	<b>475</b>	<b>388</b>	<b>169</b>	<b>391</b>	<b>200</b>	<b>176</b>	<b>397</b>	<b>303</b>	<b>342</b>	<b>638</b>	<b>197</b>	<b>687</b>	<b>199</b>	<b>143</b>	<b>258</b>	<b>132</b>	<b>7407</b>

**APPENDIX 9.** Total number of species captured at each site in 2013.

2013 Species	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	3	16	12	11	9	28	27	8	1	16	6	9	10	0	12	10	7	4	12	20	8	1	230
Murray cod	0	1	1	0	0	0	1	3	0	0	0	0	0	0	0	0	0	0	0	0	1	0	7
Silver perch	0	2	0	0	1	0	0	0	0	0	0	2	0	0	1	0	0	0	1	0	0	0	7
Bony herring	422	173	247	52	203	576	269	89	175	487	119	160	1012	182	93	313	187	138	121	126	214	150	5508
Australian smelt	6	3	2	0	62	28	25	30	23	1	3	3	3	11	3	0	4	2	1	3	1	1	215
Murray rainbowfish	0	2	0	24	36	39	25	0	1	8	0	0	7	8	10	0	0	29	7	0	1	3	200
Flathead gudgeon	3	0	1	5	0	0	2	0	0	7	0	0	0	1	0	0	42	3	0	1	2	2	69
Unspecked hardyhead	0	1	0	8	1	14	0	0	8	1	0	0	5	1	3	0	1	4	0	0	0	37	84
Carp gudgeon spp	0	0	0	14	0	0	1	0	0	0	0	0	0	0	0	0	2	0	0	0	1	10	28
Freshwater catfish	0	0	2	3	4	0	0	1	0	0	0	0	0	2	0	1	0	1	0	0	1	0	15
Common carp	74	41	77	61	47	84	54	48	11	63	58	69	19	14	97	61	53	41	24	124	55	43	1218
Gambusia	0	0	0	11	0	8	1	0	0	3	1	0	0	3	0	0	0	8	0	0	0	8	43
Goldfish	0	0	2	8	3	2	3	0	7	0	8	3	1	0	6	1	0	1	0	0	1	6	52
<b>Total species</b>	<b>5</b>	<b>8</b>	<b>8</b>	<b>10</b>	<b>9</b>	<b>8</b>	<b>10</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>6</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>8</b>	<b>5</b>	<b>7</b>	<b>10</b>	<b>6</b>	<b>5</b>	<b>10</b>	<b>10</b>	<b>13</b>
<b>Total fish/site</b>	<b>508</b>	<b>239</b>	<b>344</b>	<b>197</b>	<b>366</b>	<b>779</b>	<b>408</b>	<b>179</b>	<b>226</b>	<b>586</b>	<b>195</b>	<b>246</b>	<b>1057</b>	<b>222</b>	<b>225</b>	<b>386</b>	<b>296</b>	<b>231</b>	<b>166</b>	<b>274</b>	<b>285</b>	<b>261</b>	<b>7,676</b>

**APPENDIX 10.** Total number of species captured at each site in 2014.

2014																							
	Site Number																						
Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	Grand Total
Golden perch	7	13	13	14	10		17	7		8	5	1	2	2	5	12	5	3	7	5	11	1	148
Murray cod		2						5															7
Silver perch		1			1									1	1				1				5
Bony herring	381	266	156	51	1290		420	76	117	183	154	211	332	272	63	313	231	119	182	211	161	36	5225
Australian smelt	8	13	1		62		7	19	8	3	4	5	5	6			2		3	2	3		151
Murray rainbowfish	5	7	7	18	18		19	6	2	20	6	3	25	41	17	3	3	5	5		7	18	235
Flathead gudgeon				2			5										8	2			1	17	35
Unspecked hardyhead	1	2			1		3		4	8	1			20	18	8		1				22	89
Carp gudgeon spp	1			2			6			1			1	4	2	3	2			1	2	197	222
Freshwater catfish		1		1	1							1					1	1					6
Common carp	13	15	21	38	13		51	29	19	60	37	14	33	16	40	43	19	27	9	37	29	27	590
Gambusia				2	1		2			3				1	6			4				46	65
Goldfish	11	1	2	1	1		8	1	1	22	12	9	17	8	9	33	5	14	3	1	2	10	171
Spangled perch														1									1
<b>Total species</b>	<b>8</b>	<b>10</b>	<b>6</b>	<b>9</b>	<b>10</b>		<b>10</b>	<b>7</b>	<b>6</b>	<b>9</b>	<b>7</b>	<b>7</b>	<b>7</b>	<b>11</b>	<b>8</b>	<b>8</b>	<b>9</b>	<b>9</b>	<b>7</b>	<b>6</b>	<b>8</b>	<b>9</b>	<b>14</b>
<b>Total fish/site</b>	<b>427</b>	<b>321</b>	<b>200</b>	<b>129</b>	<b>1398</b>		<b>538</b>	<b>143</b>	<b>151</b>	<b>308</b>	<b>219</b>	<b>244</b>	<b>415</b>	<b>372</b>	<b>160</b>	<b>416</b>	<b>276</b>	<b>176</b>	<b>210</b>	<b>257</b>	<b>216</b>	<b>374</b>	<b>6950</b>

**APPENDIX 11.** Total number of species captured at each site in 2015.

2015	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	6	11	7	10	5	11	13	6	3	16	7	1	3	3		5	6		6	11	13		143
Murray cod		4	1	1	3		3	1											1				14
Silver perch		1			1	1	2		1				2		4				2				14
Bony herring	983	719	332	94	514	610	1335	451	215	250	414	383	1317	719	168	406	266	254	377	245	192	70	10314
Australian smelt	38	88	19	10	155	187	189	61	12	37	44	9	37	13	14	6	21	11	57	9	6	6	1029
Murray rainbowfish	61	50	11	38	70	19	74	18	5	35	15		43	32	9	13	4	69	68	12	4	2	652
Flathead gudgeon	1											1		2			50	2	5	1	3		65
Dwarf flathead gudgeon		1		1												1							3
Unspecked hardyhead	26		9	71	3	17	12	2	50	15	2	4	64	190	57	14	24	59	7	5	1	24	656
Carp gudgeon spp	5	2	6	10	1	4	8	3	9	10	2		8	18	1	10	5	2	2			31	137
Freshwater catfish				1	2												1						4
Common carp	78	13	73	28	14	19	11	36	68	59	29	31	38	19	30	59	25	7	26	36	20	11	730
Gambusia			3	26			1		5	33	2	2	3		4	4	7	16	3	1		16	126
Goldfish	5	3	21	8		3	2	1	25	44	39	14	59	11	5	24	15	1	5	4	2	8	299
Redfin perch													2						1				3
Spangled Perch																	1						1
<b>Total species</b>	<b>9</b>	<b>10</b>	<b>10</b>	<b>12</b>	<b>10</b>	<b>9</b>	<b>11</b>	<b>9</b>	<b>10</b>	<b>9</b>	<b>9</b>	<b>8</b>	<b>10</b>	<b>9</b>	<b>9</b>	<b>10</b>	<b>12</b>	<b>10</b>	<b>12</b>	<b>9</b>	<b>8</b>	<b>8</b>	<b>16</b>
<b>Total fish/site</b>	<b>1203</b>	<b>892</b>	<b>482</b>	<b>298</b>	<b>768</b>	<b>871</b>	<b>1650</b>	<b>579</b>	<b>393</b>	<b>499</b>	<b>554</b>	<b>445</b>	<b>1574</b>	<b>1007</b>	<b>292</b>	<b>542</b>	<b>425</b>	<b>422</b>	<b>559</b>	<b>324</b>	<b>241</b>	<b>168</b>	<b>14190</b>

**APPENDIX 12.** Total number of species captured at each site in 2016.

2016 Species	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	3	6	4	4	8	6	10	13		1	3	2	2	7	1	2	8	2	4	1	12		99
Murray cod		1			3		4	2				2							1				13
Silver perch		3		1										1					1		1		7
Bony herring	1507	1294	1147	172	4153	901	4211	429	367	73	340	250	1059	255	10	1550	344	96	579	173	311		19221
Australian smelt	52	72	13	29	145	96	104	45	24	14	9	10	84	20	22	22	51	27	49	9	19		916
Murray rainbowfish	32	17	3	61	37		29	12	2	1	4	1	49	77	23	18	7	28	63	1	25		490
Flathead gudgeon				1						1				3			5	1	2		1		14
Dwarf flathead gudgeon			1											1			1				1		4
Unspecked hardyhead	83	32	7	353	38	25	60	7	287	12	7	1	309	657	221	78	69	35	113	20	27		2441
Carp gudgeon spp	22		8	41	3	5	19	2	17	19	4	2	14	19	1	35	13	1	9	2	15		251
Freshwater catfish						1																	1
Common carp	20	5	15	14	9	10	5	9	23	15	24	18	21	24	18	27	15	9	26	12	20		339
Gambusia	1	1	8	108	9	1	9		8	11	16	2	3	3	16	4	18	17	27	23	15		300
Goldfish	8		10	18	2		1	2	33	22	55	6	15	28		50	31	5	16	15	14		331
Redfin perch																	1						1
<b>Total species</b>	<b>9</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>10</b>	<b>8</b>	<b>10</b>	<b>9</b>	<b>8</b>	<b>10</b>	<b>9</b>	<b>10</b>	<b>9</b>	<b>12</b>	<b>8</b>	<b>9</b>	<b>12</b>	<b>10</b>	<b>12</b>	<b>9</b>	<b>12</b>	<b>0</b>	<b>15</b>
<b>Total fish/site</b>	<b>1728</b>	<b>1431</b>	<b>1216</b>	<b>802</b>	<b>4407</b>	<b>1045</b>	<b>4452</b>	<b>521</b>	<b>761</b>	<b>169</b>	<b>462</b>	<b>294</b>	<b>1556</b>	<b>1095</b>	<b>312</b>	<b>1786</b>	<b>563</b>	<b>221</b>	<b>890</b>	<b>256</b>	<b>461</b>	<b>0</b>	<b>24428</b>

**APPENDIX 13.** Total number of species captured at each site in 2017.

2017		Site Number																				Grand Total	
Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21		22
Golden perch	1	19	4	12	16	4	5	13	2	6	1	2	11	2	6	12	5	6	7	1	4		139
Murray cod		1						3													1		5
Silver perch		1			2								1										4
Bony herring	1031	282	87	117	1757	445	975	102	496	670	243	215	1786	288	223	650	382	136	600	139	136	285	11045
Australian smelt	31	235	17	9	202	470	437	47	6	31	14	16	152	162	2	2	160	3	153	3	14	3	2169
Murray rainbowfish	3	49		9	6	17	13	2	4	5			15	20	3	4	2	15	16		9	3	195
Flathead gudgeon						1											2					1	4
Unspecked hardyhead	25	90	18	50	177	99	124	12	14	18	4	3	596	207	31	13	43	36	36	8	31	52	1687
Carp gudgeon spp	2	9	5	31	5	9		1		4	1	1	5	13	2	3	9	6	10	12		53	181
Freshwater catfish					1													1					2
Common carp	243	206	148	348	185	138	75	56	124	242	97	67	559	128	458	261	176	250	135	284	392	592	5164
Gambusia		7	7	194	18	10	3	4	5	5	8	9	1	13	24	2	8	21	4	20	9	26	398
Goldfish	223	56	88	129	67	129	21	13	70	156	135	46	270	103	40	186	66	20	35	225	302	137	2517
Redfin perch		2	1		1	2	1			3			1	1		2	4		2	3	4		27
<b>Total species</b>	<b>8</b>	<b>11</b>	<b>8</b>	<b>9</b>	<b>11</b>	<b>10</b>	<b>8</b>	<b>10</b>	<b>8</b>	<b>9</b>	<b>8</b>	<b>8</b>	<b>10</b>	<b>10</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>10</b>	<b>10</b>	<b>9</b>	<b>10</b>	<b>9</b>	<b>14</b>
<b>Total fish/site</b>	<b>1559</b>	<b>955</b>	<b>374</b>	<b>899</b>	<b>2436</b>	<b>1322</b>	<b>1653</b>	<b>253</b>	<b>721</b>	<b>1137</b>	<b>503</b>	<b>359</b>	<b>3396</b>	<b>937</b>	<b>789</b>	<b>1135</b>	<b>857</b>	<b>494</b>	<b>998</b>	<b>695</b>	<b>902</b>	<b>1152</b>	<b>23537</b>

**APPENDIX 14.** Total number of species captured at each site in 2018.

2018 Species	Site Number																					Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
Golden perch	7	4	1	13	8	8	6	5		2		4	6	3	6	6	1	6	2	9	15	112
Murray cod		2	1	1	1			5	4				1					2	1			18
Silver perch		1		1	5								1							1		9
Bony herring	547	1002	530	195	1982	1064	1352	400	132	97	243	155	903	520	86	809	219	229	509	102	75	11151
Australian smelt	166	67	178	9	74	731	79	151	161	25	26	64	147	166	2	19	64	7	178	35	28	2377
Murray rainbowfish	6	2	1	19	103	27	17	8		15		1	30	168	36	20		50	47	3	4	557
Flathead gudgeon									4				1	4	1		5	3	8	6	1	33
Unspecked hardyhead	14	14	6	136	103	116	45		74	36			170	557	128	12	12	24	45	1	1	1494
Carp gudgeon spp	3	1	3	9	1	4	1		1	6			1	8	2	8	4	4	7	3	2	68
Freshwater catfish	1			2	2																	5
Common carp	34	8	28	27	37	11	16	29	22	30	17	14	22	47	49	45	19	9	8	33	34	539
Gambusia			3	86	2	6			2		1	5	2		6	1	3	8	17	18		160
Goldfish	14		5	5		1		1	4	14		3	10	1	8	110	1	7		1	3	188
Redfin perch															1		2	9		1		13
<b>Total species</b>	<b>9</b>	<b>9</b>	<b>10</b>	<b>12</b>	<b>11</b>	<b>9</b>	<b>8</b>	<b>7</b>	<b>8</b>	<b>8</b>	<b>4</b>	<b>7</b>	<b>12</b>	<b>9</b>	<b>11</b>	<b>9</b>	<b>10</b>	<b>12</b>	<b>11</b>	<b>11</b>	<b>9</b>	<b>14</b>
<b>Total fish/site</b>	<b>792</b>	<b>1101</b>	<b>756</b>	<b>503</b>	<b>2318</b>	<b>1968</b>	<b>1521</b>	<b>598</b>	<b>400</b>	<b>225</b>	<b>287</b>	<b>246</b>	<b>1294</b>	<b>1474</b>	<b>325</b>	<b>1030</b>	<b>330</b>	<b>358</b>	<b>823</b>	<b>212</b>	<b>163</b>	<b>16724</b>

**APPENDIX 15.** Total number of species captured at each site in 2019.

2019 Species	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	5	3		4	6	3	3	3		3	1	2	2	3	5	1	1	7	8	1	7		68
Murray cod	1		1			1	1	6				1							2				13
Silver perch				1	2										1				1				5
Bony herring	1353	998	346	107	708	1362	1245	458	337	175	165	201	2525	904	249	1867	668	226	982	276	337	41	15530
Australian smelt	31	88	12	10	55	177	108	21	29	5	15	13	15	26		11	2		30	1	6	25	680
Murray rainbowfish	5			15	46	12	20			5	2		8	8	22	4	1	32	15	1	5	3	204
Flathead gudgeon																	1					5	6
Dwarf flathead gudgeon																						1	1
Unspecked hardyhead	10	3	3	40	27	83	22		16	57	3	3	18	57	113	15	20	21	1	11	1	1	525
Carp gudgeon spp	10			5		5	10	1	1	3	1		3	6		25		2	3		1	17	93
Freshwater catfish				1	1																		2
Common carp	36	8	29	40	31	24	13	12	20	92	52	20	10	1	33	52	17	18	40	15	33	104	700
Gambusia			1							1	2				1	2	2	3	3	1		11	27
Goldfish		1	5	9	1	1		7	2	10	15	3	3	2		9	2		6	2	1	67	146
Redfin perch		1	1	2						2			2	1		4	3		1	3	1		21
<b>Total species</b>	<b>8</b>	<b>6</b>	<b>7</b>	<b>10</b>	<b>9</b>	<b>9</b>	<b>8</b>	<b>7</b>	<b>6</b>	<b>9</b>	<b>9</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>7</b>	<b>10</b>	<b>7</b>	<b>12</b>	<b>9</b>	<b>9</b>	<b>10</b>	<b>15</b>	
<b>Total fish/site</b>	<b>1451</b>	<b>1102</b>	<b>398</b>	<b>234</b>	<b>877</b>	<b>1668</b>	<b>1422</b>	<b>508</b>	<b>405</b>	<b>353</b>	<b>256</b>	<b>243</b>	<b>2586</b>	<b>1008</b>	<b>424</b>	<b>1990</b>	<b>717</b>	<b>309</b>	<b>1092</b>	<b>311</b>	<b>392</b>	<b>275</b>	<b>18021</b>

**APPENDIX 16.** Total number of species captured at each site in 2020.

2020 Species	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	5	6	2	3	2	1	1					2	1		9	2	4	2			6		46
Murray cod	1	1	1	2	3	6	10	6				1			1	0	0	1					33
Silver perch																							
Bony herring	1109	1313	1262	335	1972	2906	4504	1033	336	624	306	364	1984	1380	313	799	871	280		222	615	83	22611
Australian smelt	26	62	114	37	46	146	62	96	148	74	24	71	26	120	2	13	32	7		33	24	1	1164
Murray rainbowfish	7	1	5	59	45	8	10	1		2	1		5	20	84	3	3	19		3	17		293
Flathead gudgeon				1		2				1		1		3	4	1	7			1	6	75	102
Dwarf flathead gudgeon																						1	1
Unspecked hardyhead				77	5	17	7		9	4		1	5	348	408	3	7	17		12	2		922
Carp gudgeon spp	2		1	3		7	7		1	6	1		1	20	7	4	8	2		5	4	30	109
Freshwater catfish					1	1									1								3
Common carp	9	11	23	11	26	17	11	17	15	50	8	17	9	7	44	24	6	7		9	31	60	412
Gambusia				11		2	2			12	1			3	51		7			10		4	103
Goldfish	2	1	3	2				1	2	16				1	2	2	19			9	2	28	90
Redfin perch					1					3	5	2		1	1	1	2	14	1		5		36
<b>Total species</b>	<b>8</b>	<b>7</b>	<b>8</b>	<b>11</b>	<b>8</b>	<b>11</b>	<b>9</b>	<b>6</b>	<b>6</b>	<b>9</b>	<b>6</b>	<b>7</b>	<b>7</b>	<b>10</b>	<b>13</b>	<b>11</b>	<b>12</b>	<b>9</b>	<b>0</b>	<b>10</b>	<b>9</b>	<b>8</b>	<b>14</b>
<b>Total fish/site</b>	<b>1161</b>	<b>1395</b>	<b>1411</b>	<b>541</b>	<b>2101</b>	<b>3113</b>	<b>4614</b>	<b>1154</b>	<b>514</b>	<b>794</b>	<b>343</b>	<b>457</b>	<b>2032</b>	<b>1903</b>	<b>927</b>	<b>853</b>	<b>978</b>	<b>336</b>	<b>0</b>	<b>309</b>	<b>707</b>	<b>282</b>	<b>25925</b>

**APPENDIX 17.** Total number of species captured at each site in 2021.

2021 Species	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	1	5	1	5	5	2						4	1	3	2	4	8	6	2		12	61	
Murray cod		5		4	5	2	7	5				3		1					6			38	
Silver perch					2			2								1						5	
Bony herring	796	2261	355	227	3451	2100	3176	534	47	566	470	239	285	1817	5	617	645	580	3111	509	438	22229	
Australian smelt	120	130	43	30	175	216	232	113	110	138	41	41	79	260	8	46	180	79	573	72	34	2720	
Murray rainbowfish	10	17	28	55	17	4	14	1		3	2	7	29	13	4	5	20	41	77	1	24	372	
Flathead gudgeon	5		1				3	1	6	3		2	1	15	27	4	7	3	6	2	2	88	
Dwarf flathead gudgeon																						0	
Unspecked hardyhead	23	13	15	41	56	187	10		32	14		3	62	323	265	34	12	56	14	7	1	1168	
Carp gudgeon spp	2		1	2		9	5	1		1		1	15	39	5	3	1	4	4	5	4	102	
Freshwater catfish					1														1			2	
Common carp	24	10	25	15	34	14	15	22	7	47	20	16	23	12	32	42	5	23	38	24	36	484	
Gambusia	2	3	1	17	2	2			3	7	1		6	12	230		6	32	2	5		331	
Goldfish	10		7	11		8		2		46	8	14	8	2	18	25	8	11	9	9	4	200	
Redfin perch	3	1			1	1				5	3				2	6	25	1	8	4	6	66	
<b>Total species</b>	<b>11</b>	<b>9</b>	<b>10</b>	<b>10</b>	<b>11</b>	<b>11</b>	<b>8</b>	<b>9</b>	<b>6</b>	<b>10</b>	<b>7</b>	<b>10</b>	<b>10</b>	<b>11</b>	<b>11</b>	<b>11</b>	<b>11</b>	<b>11</b>	<b>13</b>	<b>10</b>	<b>10</b>	<b>14</b>	
<b>Total fish/site</b>	<b>996</b>	<b>2445</b>	<b>477</b>	<b>407</b>	<b>3749</b>	<b>2545</b>	<b>3462</b>	<b>681</b>	<b>205</b>	<b>830</b>	<b>545</b>	<b>330</b>	<b>509</b>	<b>2497</b>	<b>598</b>	<b>787</b>	<b>917</b>	<b>836</b>	<b>3851</b>	<b>638</b>	<b>561</b>	<b>27866</b>	

**APPENDIX 18.** Total number of species captured at each site in 2022.

2022 Species	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	2	4		3	3	3		2	1	5	2	2	2	8	5	1	2	7	2	7	1		62
Murray cod	1	5		2	2		11	4		1		1	2	1									30
Silver perch	1			1	3						1	1											7
Bony herring	383	256	269	70	372	489	657	185	485	871	526	410	744	944	715	859	827	321	201	622	986		11192
Australian smelt	29	92	64	9	183	356	87	30	12	38	60	172	216	128	3	5	99	28	68	54	124		1857
Murray rainbowfish	11	10	7	29	6	24	57	3	5	4	6	1	21	27	9	7	11	18	26	7	21		310
Flathead gudgeon												1					4			17	10		32
Dwarf flathead gudgeon																							0
Unspecked hardyhead	3	4	6	67	8	83	10	1	26	29	2	8	30	123	169	6	18	43	20	6	14		676
Carp gudgeon spp	1		1		1		2		3	4			1	5		3	1			1	3		26
Freshwater catfish																		1	1				2
Common carp	149	78	110	182	64	57	70	47	103	145	105	268	244	102	147	165	58	89	192	249	417		3041
Gambusia		1	3	22	5	3		1	1	1		4	89	1	10	3	6	2	2	2	15		171
Goldfish	1	1	14	7	2	8	1		7	5	6	19	30	30	2	8	17	4		9	36		207
Redfin perch	5		2			5		1	2	3	4	2	6	1	1	17	2	3		7	9		70
<b>Total species</b>	<b>11</b>	<b>9</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>9</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>9</b>	<b>12</b>	<b>11</b>	<b>11</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>10</b>	<b>8</b>	<b>11</b>	<b>11</b>		
<b>Total fish/site</b>	<b>586</b>	<b>451</b>	<b>476</b>	<b>392</b>	<b>649</b>	<b>1028</b>	<b>895</b>	<b>274</b>	<b>645</b>	<b>1106</b>	<b>712</b>	<b>889</b>	<b>1385</b>	<b>1370</b>	<b>1061</b>	<b>1074</b>	<b>1045</b>	<b>516</b>	<b>512</b>	<b>981</b>	<b>1636</b>		<b>17683</b>

**APPENDIX 19.** Total number of species captured at each site in 2023.

2023 Species	Site Number																						Grand Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
Golden perch	3	11	4	6	5	12		11		8	7	6	8	7	1	2	3	10	4	3	6		117
Murray cod		2				1	2	12				1				1		2		1			22
Silver perch		1																					1
Bony herring	610	1976	358	25	516	366	511	1817	219	184	156	145	###	627	299	1606	256	196	176	93	315	101	12692
Australian smelt	190	55	122	2	12	508	114	133	77	52	67	3	154	88	4	64	94	6	30	4	102	38	1919
Murray rainbowfish	10	5	2	22		1	20		4			1	6	17	2	95	11	2	68		5	2	273
Flathead gudgeon	1																						1
Dwarf flathead gudgeon																							0
Unspecked hardyhead	4	3	2			1			4				16	22	4	23	5	1	7	1	1	2	96
Carp gudgeon spp	1								1				3			1		2		1			9
Freshwater catfish				1	1														1		1		4
Common carp	533	740	585	2008	309	283	280	708	620	1216	1826	1088	492	609	1887	672	438	1133	584	3081	1070	650	20812
Gambusia	23	11	24	65	49	20	25	31	19	88	74	2	66	13	2	32	37	115	38	7	48	15	804
Goldfish	184	64	151	8	7	29	20	49	59	28	159	10	62	188	20	970	45	205	96	13	83	37	2487
Oriental weatherloach																		1		1			2
Redfin perch	49	20	65	22	26	35	22	52	35	65	75	62	31	50	16	44	44	78	45	68	48	5	957
<b>Total species</b>	<b>11</b>	<b>11</b>	<b>9</b>	<b>9</b>	<b>8</b>	<b>10</b>	<b>8</b>	<b>8</b>	<b>9</b>	<b>7</b>	<b>7</b>	<b>9</b>	<b>10</b>	<b>9</b>	<b>9</b>	<b>11</b>	<b>10</b>	<b>11</b>	<b>11</b>	<b>10</b>	<b>10</b>	<b>8</b>	
<b>Total fish/site</b>	<b>1608</b>	<b>2888</b>	<b>1313</b>	<b>2159</b>	<b>925</b>	<b>1256</b>	<b>994</b>	<b>2813</b>	<b>1038</b>	<b>1641</b>	<b>2364</b>	<b>1318</b>	<b>2978</b>	<b>1621</b>	<b>2235</b>	<b>3510</b>	<b>934</b>	<b>1750</b>	<b>1050</b>	<b>3272</b>	<b>1679</b>	<b>850</b>	<b>40196</b>

**APPENDIX 20.** Total number of species captured at each site in 2024.

2024		Site Number																	
Species	1	2	3	4	6	7	8	9	10	11	12	13	14	15	16	17	18	21	Grand Total
Golden perch	1	5	1	1	4		5		2	2	1	1	4		2	1		6	36
Murray cod		6	1	1	2		8												18
Silver perch		1		2															3
Bony herring	376	320	353	44	374	1119	97	188	125	160	190	487	389	14	446	398	35	236	5351
Australian smelt	3	32			46	17	17	4	9	3	2	12	3		1	22	1	1	173
Murray rainbowfish	1	1		5	5	2		1	1			13	2	2	1		2	2	38
Flathead gudgeon				1			1	2								1			5
Dwarf flathead gudgeon																			0
Unspecked hardyhead					2	2		1	1			21	99	1					127
Carp gudgeon spp								1					1			1			3
Freshwater catfish																		1	1
Common carp	75	70	35	42	40	17	77	111	49	52	29	32	17	38	53	20	23	103	883
Gambusia					1							4	5						10
Goldfish	2		1		1			7	1			6		2	10			4	34
Oriental weatherloach																			0
Redfin perch		1															1		2
<b>Total species</b>	<b>6</b>	<b>8</b>	<b>5</b>	<b>7</b>	<b>9</b>	<b>5</b>	<b>6</b>	<b>8</b>	<b>7</b>	<b>4</b>	<b>4</b>	<b>8</b>	<b>8</b>	<b>5</b>	<b>6</b>	<b>6</b>	<b>5</b>	<b>7</b>	
<b>Total fish/site</b>	<b>458</b>	<b>436</b>	<b>391</b>	<b>96</b>	<b>475</b>	<b>1157</b>	<b>205</b>	<b>315</b>	<b>188</b>	<b>217</b>	<b>222</b>	<b>576</b>	<b>520</b>	<b>57</b>	<b>513</b>	<b>443</b>	<b>62</b>	<b>353</b>	<b>6684</b>