

Inland Waters & Catchment Ecology

Lower Lakes Vegetation Condition Monitoring – 2020-21



Jason Nicol, Kate Frahn, Susan Gehrig and Kelly Marsland

**SARDI Publication No. F2009/000370-12
SARDI Research Report Series No. 1108**

**SARDI Aquatics Sciences
PO Box 120 Henley Beach SA 5022**

September 2021



**Government
of South Australia**
Department of Primary
Industries and Regions



Australian Government



Government of South Australia
Department for Environment
and Water

SARDI



**SOUTH AUSTRALIAN
RESEARCH AND
DEVELOPMENT
INSTITUTE**

Lower Lakes Vegetation Condition Monitoring – 2020-21

**Jason Nicol, Kate Frahn, Susan Gehrig and
Kelly Marsland**

**SARDI Publication No. F2009/000370-12
SARDI Research Report Series No. 1108**

September 2021

This publication may be cited as:

Nicol, J.M., Frahn, K.A., Gehrig, S.L. and Marsland, K.B. (2021). Lower Lakes Vegetation Condition Monitoring 2020 - 21. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2009/000370-12. SARDI Research Report Series No. 1108. 107pp.

Cover Photo: Emergent vegetation at Clayton Bay (Jason Nicol).

DISCLAIMER

The contents of this publication do not purport to represent the position of the Commonwealth of Australia or the MDBA in any way and are presented for the purpose of informing and stimulating discussion for improved management of the Basin's natural resources. To the extent permitted by law, the copyright holders (including its employees and consultants) exclude all liability to any person for any consequences, including but not limited to all losses, damages, costs, expenses and any other compensation, arising directly or indirectly from using this report (in part or in whole) and any information or material contained in it. The authors warrant that they have taken all reasonable care in producing this report. The report has been through the SARDI internal review process, and has been formally approved for release by the Research Director, Aquatic Sciences. Although all reasonable efforts have been made to ensure quality, SARDI does not warrant that the information in this report is free from errors or omissions. SARDI and its employees do not warrant or make any representation regarding the use, or results of the use, of the information contained herein as regards to its correctness, accuracy, reliability and currency or otherwise. SARDI and its employees expressly disclaim all liability or responsibility to any person using the information or advice. Use of the information and data contained in this report is at the user's sole risk. If users rely on the information they are responsible for ensuring by independent verification its accuracy, currency or completeness. The SARDI Report Series is an Administrative Report Series which has not been reviewed outside the department and is not considered peer-reviewed literature. Material presented in these Administrative Reports may later be published in formal peer-reviewed scientific literature.

© 2021 SARDI

This work is copyright. Apart from any use as permitted under the *Copyright Act* 1968 (Cth), no part may be reproduced by any process, electronic or otherwise, without the specific written permission of the copyright owner. Neither may information be stored electronically in any form whatsoever without such permission. With the exception of the Commonwealth Coat of Arms, the Murray-Darling Basin Authority logo and photographs, all material presented in this document is provided under a Creative Commons Attribution 4.0 International licence (<https://creativecommons.org/licenses/by/4.0/>). For the avoidance of any doubt, this licence only applies to the material set out in this document.



The details of the licence are available on the Creative Commons website (accessible using the links provided) as is the full legal code for the CC BY 4.0 licence (<https://creativecommons.org/licenses/by/4.0/legalcode>).

Source: Licensed from the Department of Environment and Water (DEW) under a Creative Commons Attribution 4.0 International Licence. Enquiries regarding the licence and any use of the document are welcome to: Adrienne Rumbelow, LLCMM Icon Site Coordinator adrienne.rumbelow@sa.gov.au.

Author(s): Jason Nicol, Kate Frahn, Susan Gehrig and Kelly Marsland

Reviewer(s): Greg Ferguson (SARDI), Kirsty Wedge and Adrienne Rumbelow (DEW)

Approved by: Assoc. Prof Qifeng Ye
Science Leader – Inland Waters & Catchment Ecology

Signed: 

Date: 20 September 2021

Distribution: DEW, MDBA, SARDI Aquatic Sciences, Parliamentary Library, State Library and National Library

Circulation: OFFICIAL

ALL ENQUIRIES

South Australian Research and Development Institute - SARDI Aquatic Sciences

2 Hamra Avenue West Beach SA 5024

PO Box 120 Henley Beach SA 5022

P: (08) 8207 5400 **F:** (08) 8207 5415 **E:** pirsa.sardiaquatics@sa.gov.au

W: <http://www.pir.sa.gov.au/research>

TABLE OF CONTENTS

TABLE OF CONTENTS	IV
LIST OF FIGURES	V
LIST OF TABLES.....	IX
LIST OF APPENDICES	X
ACKNOWLEDGEMENTS	XI
EXECUTIVE SUMMARY	1
1. INTRODUCTION	4
1.1. Background.....	4
1.2. Aquatic and littoral vegetation target revision	7
1.3. Objectives.....	13
2. METHODS.....	14
2.1. Study site, hydrology and salinity.....	14
2.2. Understorey vegetation survey protocol.....	19
2.3. Data Analysis	21
3. RESULTS	22
3.1. Change through time of the Lower Lakes plant community from spring 2008 to autumn 2021	22
3.2. TLM targets	32
4. DISCUSSION AND MANAGEMENT IMPLICATIONS	75
4.1. Impacts of water level and salinity.....	75
4.2. Change in plant community, spring 2008 to autumn 2021	76
4.3. The Living Murray targets and condition scores	77
4.4. Further studies.....	81
REFERENCES	83
APPENDICES.....	91

LIST OF FIGURES

Figure 1: Map of Lakes Alexandrina and Albert and Goolwa Channel showing the location of lakeshore and wetland vegetation monitoring sites (site numbers correspond to Table 7) and major flow control structures present in winter 2010 (where sites are in close proximity they may not be visible on map).....	16
Figure 2: Daily mean water levels in Goolwa Channel (Signal Point), Lake Alexandrina (Milang) and Lake Albert (Meningie) from August 2008 to May 2021 (Department of Environment and Water 2021b).....	17
Figure 3: Daily mean surface water electrical conductivity (EC) in Goolwa Channel (Signal Point), Lake Alexandrina (Milang) and Lake Albert (Meningie) from August 2008 to May 2021 (Department of Environment and Water 2021a).....	18
Figure 4: Vegetation surveying protocol for lakeshore sites: plan view showing placement of quadrats relative to the shoreline.	21
Figure 5: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in Lake Alexandrina (Sp denotes spring; Au denotes autumn).	24
Figure 6: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in Lake Albert (Sp denotes spring; Au denotes autumn).....	25
Figure 7: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in Goolwa Channel (Sp denotes spring; Au denotes autumn).....	27
Figure 8: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in permanent wetlands (Sp denotes spring; Au denotes autumn).....	28
Figure 9: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in seasonal wetlands (Sp denotes spring; Au denotes autumn).....	30
Figure 10: Mean native amphibious species richness per quadrat for Lake Alexandrina, Lake Albert and Goolwa Channel at a. +0.6 m and b. +0.8 m AHD over the condition monitoring program (error bars = ± 1 S.E.).	32
Figure 11: Percentage of quadrats containing a combined cover of <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 75% in the littoral zone of Lake Alexandrina from spring 2008 to autumn 2021.....	33
Figure 12: Percentage of quadrats containing a combined cover of <i>Paspalum distichum</i> and <i>Cenchrus clandestinus</i> greater than 50% in the littoral zone of Lake Alexandrina from spring 2008 to autumn 2021.....	34
Figure 13: Percentage of quadrats containing a cover native amphibious species greater than 5% in the littoral zone of Lake Alexandrina from spring 2008 to autumn 2021.....	35

Figure 14: Percentage of quadrats containing a cover of native emergent species other than <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 5% in the littoral zone of Lake Alexandrina from spring 2008 to autumn 2021.....	36
Figure 15: Percentage of quadrats containing a combined cover of <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 50% in the aquatic zone of Lake Alexandrina from spring 2008 to autumn 2021.....	37
Figure 16: Percentage of quadrats containing a cover of native emergent species other than <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 5% in the aquatic zone of Lake Alexandrina from spring 2008 to autumn 2021.....	38
Figure 17: Percentage of quadrats containing a cover of native submergent species greater than 5% in the aquatic zone of Lake Alexandrina from spring 2008 to autumn 2021.....	39
Figure 18: Whole of habitat condition score for Lake Alexandrina from spring 2008 to autumn 2021.....	40
Figure 19: Percentage of quadrats containing a combined cover of <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 75% in the littoral zone of Lake Albert from spring 2008 to autumn 2021.....	41
Figure 20: Percentage of quadrats containing a combined cover of <i>Paspalum distichum</i> and <i>Cenchrus clandestinus</i> greater than 50% in the littoral zone of Lake Albert from spring 2008 to autumn 2021.....	42
Figure 21: Percentage of quadrats containing a cover of native amphibious species greater than 5% in the littoral zone of Lake Albert from spring 2008 to autumn 2021.....	43
Figure 22: Percentage of quadrats containing a combined cover of <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 50% in the aquatic zone of Lake Albert from spring 2008 to autumn 2021.....	44
Figure 23: Percentage of quadrats containing a cover of native emergent species other than <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 5% in the aquatic zone of Lake Albert from spring 2008 to autumn 2021.....	45
Figure 24: Percentage of quadrats containing a cover of native submergent species greater than 5% in the aquatic zone of Lake Albert from spring 2008 to autumn 2021.....	46
Figure 25: Whole of habitat condition score for Lake Albert from spring 2008 to autumn 2021..	47
Figure 26: Percentage of quadrats containing a combined cover of <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 75% in the littoral zone of Goolwa Channel from spring 2008 to autumn 2021.....	48

Figure 27: Percentage of quadrats containing a combined cover of <i>Paspalum distichum</i> and <i>Cenchrus clandestinus</i> greater than 50% in the littoral zone of Goolwa Channel from spring 2008 to autumn 2021.....	49
Figure 28: Percentage of quadrats containing a cover native amphibious species greater than 5% in the littoral zone of Goolwa Channel from spring 2008 to autumn 2021.....	50
Figure 29: Percentage of quadrats containing a cover of native emergent species other than <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 5% in the littoral zone of Goolwa Channel from spring 2008 to autumn 2021.	51
Figure 30: Percentage of quadrats containing a combined cover of <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 50% in the aquatic zone of Goolwa Channel from spring 2008 to autumn 2021.....	52
Figure 31: Percentage of quadrats containing a cover of native emergent species other than <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 5% in the aquatic zone of Goolwa Channel from spring 2008 to autumn 2021.	53
Figure 32: Percentage of quadrats containing a cover of native submergent species greater than 5% in the aquatic zone of Goolwa Channel from spring 2008 to autumn 2021.....	54
Figure 33: Percentage of quadrats containing a cover of native submergent species greater than 5% in the deep water zone of Goolwa Channel from spring 2008 to autumn 2021.....	55
Figure 34: Whole of habitat condition score for Goolwa Channel from spring 2008 to autumn 2021.	57
Figure 35: Percentage of quadrats containing a combined cover of <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 50% in the littoral zone of permanent wetlands from spring 2008 to autumn 2021.....	58
Figure 36: Percentage of quadrats containing a combined cover of <i>Paspalum distichum</i> and <i>Cenchrus clandestinus</i> greater than 50% in the littoral zone of permanent wetlands from spring 2008 to autumn 2021.	59
Figure 37: Percentage of quadrats containing a cover native amphibious species greater than 5% in the littoral zone of permanent wetlands from spring 2008 to autumn 2021.	60
Figure 38: Percentage of quadrats containing a cover of native emergent species other than <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 5% in the littoral zone of permanent wetlands from spring 2008 to autumn 2021.....	61
Figure 39: Percentage of quadrats containing a combined cover of <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 50% in the aquatic zone of permanent wetlands from spring 2008 to autumn 2021.	62

Figure 40: Percentage of quadrats containing a cover of native emergent species other than <i>Typha domingensis</i> and <i>Phragmites australis</i> greater than 5% in the aquatic zone of permanent wetlands from spring 2008 to autumn 2021.....	63
Figure 41: Percentage of quadrats containing a cover of native submergent species between 5 and 50% in the aquatic zone of permanent wetlands from spring 2008 to autumn 2021.	64
Figure 42: Whole of habitat condition score for permanent wetlands from spring 2008 to autumn 2021.....	65
Figure 43: Percentage of quadrats containing a combined cover of <i>Paspalum distichum</i> and <i>Cenchrus clandestinus</i> greater than 50% around the edge of seasonal wetlands from spring 2008 to autumn 2021.....	66
Figure 44: Percentage of quadrats containing a cover native amphibious species greater than 5% around the edge of seasonal wetlands from spring 2008 to autumn 2021.....	67
Figure 45: Percentage of quadrats containing a cover of native emergent species greater than 5% around the edge of seasonal wetlands from spring 2008 to autumn 2021.....	68
Figure 46: Percentage of quadrats containing a cover of native amphibious species greater than 5% on the bed of seasonal wetlands from spring 2008 to autumn 2021.....	69
Figure 47: Percentage of quadrats containing a cover of native emergent species greater than 5% on the beds of seasonal wetlands from spring 2008 to autumn 2021.	70
Figure 48: Percentage of quadrats containing a cover of native submergent species greater than 25% in spring on the beds of seasonal wetlands from 2008 to 2020.	71
Figure 49: Whole of habitat condition score for seasonal wetlands from spring 2008 to autumn 2021.....	72
Figure 50: Whole of lakes condition score from spring 2008 to autumn 2021.....	74

LIST OF TABLES

Table 1: Revised vegetation targets for Lake Alexandrina.....	9
Table 2: Revised vegetation targets for Lake Albert.....	10
Table 3: Revised vegetation targets for Goolwa Channel.....	10
Table 4: Revised vegetation targets for permanent wetlands (Dunns Lagoon, Hunters Creek, Angas River Mouth and Bremer River Mouth).....	11
Table 5: Revised vegetation targets for seasonal wetlands in a. spring and b. autumn (Goolwa Channel Drive, Milang Wetland, Narrung Wetland, Loveday Bay Wetland, Point Sturt Wetland and Teringie).....	12
Table 6: Icon site scores for aquatic and littoral vegetation for the Lower Lakes with the condition rating used in Matter 8 Report Cards (Department for Environment and Water 2019).....	12
Table 7: List of understorey vegetation site numbers (relative to map provided in Figure 1), site name, location, habitat type (wetland or lakeshore), number of survey sites and the year sites were established (SAMDBNRM denotes, South Australian Murray-Darling Basin Natural Resources Management Board).	19
Table 8: Modified Braun-Blanquet (1932) scale estimating cover/abundance as per Heard and Channon (1997).....	20
Table 9: Whole of lakes condition score and Matter 8 condition rating (Department for Environment and Water 2019) from spring 2008 to autumn 2021.....	74

LIST OF APPENDICES

Appendix 1: Species list, functional classification (Gehrig and Nicol 2010b), life history strategy and conservation status (state conservation status from listings in Barker <i>et al.</i> (2005) and regional conservation status from listings in Lang and Kaeheneuhl (2001) from all sites and survey dates (*denotes exotic taxon, **denotes proclaimed pest plant in South Australia, ***denotes weed of national significance # denotes listed as rare in South Australia).	91
Appendix 2: GPS coordinates (UTM format, map datum WGS84) for lakeshore and wetland understorey vegetation monitoring sites (site numbers correspond with site numbers in Figure 1).	95
Appendix 3: Taxa present (green shading) in Lake Alexandrina spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).	96
Appendix 4: Taxa present (green shading) in Lake Albert spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).....	99
Appendix 5: Taxa present (green shading) in Goolwa Channel spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).	101
Appendix 6: Taxa present (green shading) in permanent wetlands spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).	103
Appendix 7: Taxa present (green shading) in temporary wetlands spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).	106

ACKNOWLEDGEMENTS

The authors would like to thank Arron Strawbridge, Rod Ward, Kate Mason, Geoff Braddock, Robert Mrongovius, Margaret Mrongovius, Charles Andre, Jo Andre, John Rowe, Lorraine Rowe, Bev Stone, Alex Stone and the Raukkan Natural Resources Management Team for field assistance. Adrienne Rumbelow, Rod Ward, Jason Tanner, Greg Ferguson, Rebecca Turner, Carol Schmidt and Chris Bice for comments on early drafts of this report. Geoff and Libby Braddock, Henry Harvey, Chris and Beth Cowan, Derek Walker, the Raukkan Natural Resources Management Team, Charles and Jo Andre and Jamie Gibbs for access to their properties. Geoff and Libby Braddock, Charles and Jo Andre, Alex and Bev Stone for lunches, tea, coffee and scones. This project was funded by The Living Murray; a joint initiative funded by the New South Wales, Victorian, South Australian, and Commonwealth governments and coordinated by the Murray-Darling Basin Authority. This project has been managed by the Department for Environment and Water, through the Lower Lakes, Coorong and Murray Mouth icon site staff.

This report was formally reviewed by Adrienne Rumbelow, and Kirsty Wedge (DEW), Dr Greg Ferguson (SARDI Aquatic Sciences) and Dr Mike Steer, Research Director, SARDI Aquatic Sciences. The report was formally approved for release by A/Prof. Qifeng Ye, Science Leader, Inland Waters and Catchment Ecology, SARDI Aquatic Sciences.

EXECUTIVE SUMMARY

The Lower Lakes, Coorong and Murray Mouth region is one of six icon sites under “The Living Murray” (TLM) program and is an indicator site under the “Basin Plan”. The Condition Monitoring Plan for the Lower Lakes, Coorong and Murray Mouth (LLCMM) icon site (Maunsell Australia Pty Ltd 2009) identified that existing monitoring programs would not adequately assess TLM target V3, *maintain or improve aquatic and littoral vegetation in the Lower Lakes*; therefore, a monitoring program that expanded and built upon existing programs was established in spring 2008. A review undertaken by Robinson (2015) suggested that the initial aquatic and littoral vegetation target for the Lower Lakes could be improved by developing a series of quantitative targets for the site nested under the original target (now included as an objective and reported on using the same index as the Basin Plan environmental outcome reporting (Matter 8). The updated quantitative targets and methodologies are outlined in the Condition Monitoring Plan (Revised) 2017 for the Lower Lakes, Coorong and Murray Mouth Icon Site (Department for Environment, Water and Natural Resources 2017). To develop the quantitative targets the Lower Lakes were divided into five different habitats (Lake Alexandrina, Lake Albert, Goolwa Channel, permanent wetlands and seasonal wetlands), with each habitat comprising zones based on elevation. Targets were developed for species and functional groups in each zone and habitat (see Table 1 to Table 5 for detailed description of targets) and the progress of target achievement through time was assessed. This report presents the findings of the first 12 years of a monitoring program established to evaluate TLM Target Objective V3 from spring 2008 to autumn 2021.

Vegetation surveys were conducted at selected wetlands and lakeshore sites across lakes Alexandrina and Albert, Goolwa Channel, lower Finniss River, lower Currency Creek and the mouths of the Angas and Bremer Rivers. Sites established in spring 2008 and 2009 (Goolwa Channel monitoring sites) were re-surveyed. At each site, transects were established perpendicular to the shoreline and three, 1 x 3 m quadrats, separated by one metre were positioned at regular elevation intervals (defined by plant community) for wetlands or elevations (+0.8, +0.6, +0.4, +0.2, 0 and -0.5 m AHD) for lakeshores. The cover and abundance of each species present in quadrats were estimated using a modified Braun-Blanquet (1932) cover abundance score. Vegetation surveys were undertaken in spring (October 2008, 2009, November 2010, October 2011, 2012, 2013, 2015, in temporary wetlands in December 2016, all sites in November 2017, 2018, 2019 and December 2020) and autumn (March 2009 to 2017, April 2018 and May 2021). The first two years of the monitoring program coincided with a period of record

low water levels in the Lower Lakes. During this period, significant engineering interventions (i.e. construction of the Clayton Regulator and Narrung Bund and pumping of water for the environment into Narrung Wetland) also influenced plant communities and were assessed as part of the monitoring program. In August 2010, water levels in Lake Alexandrina rapidly rose to normal pool level and in September 2010, the Clayton Regulator and Narrung Bund were breached, reconnecting these areas with Lake Alexandrina. Water levels between +1.1 and +0.4 m AHD, and connectivity throughout the system, continued from 2010 throughout the remainder of the monitoring program.

Over the 12 years of condition monitoring (spring 2008 to autumn 2021), a total of 164 taxa (including 77 exotics, two weeds of national significance, five proclaimed pest plants in South Australia and one species listed as rare in South Australia) were recorded throughout the Lower Lakes (Appendix 1). Lake Alexandrina was the most species rich of the habitats with 120 taxa (including 56 exotics) recorded between spring 2008 and autumn 2021, followed by permanent wetlands (98 taxa, including 36 exotics), then temporary wetlands (95 taxa, including 43 exotics), Goolwa Channel (80 taxa, including 30 exotics) and Lake Albert the least species rich with 61 taxa (including 32 exotics).

Changes through time of the plant community in each habitat indicated a shift in floristic composition during the condition monitoring program (nMDS ordination 2008–2021). Furthermore, for each habitat (except seasonal wetlands), there was greater change in the plant community between the early surveys that reduced through time resulting in less change in vegetation between the more recent surveys. The large changes in vegetation between the early surveys were due to the colonisation of terrestrial taxa between 2008 and 2010 and subsequent extirpation and colonisation of submergent, emergent and amphibious taxa after spring 2010. The reduced rate of change between the recent surveys suggests that a stable plant community may be developing. However, sustained small changes over time may result in a significant shift in the plant community in the future. In the seasonal wetlands there were strong seasonal patterns in the plant community after spring 2010 due to seasonal inundation. Submergent species were abundant in spring, when seasonal wetlands are inundated but absent in autumn, replaced by amphibious and emergent taxa.

Achievement of the targets varied among habitats through time; generally, very few targets were achieved in all habitats when water levels were low but shortly after water levels were reinstated the number of targets achieved generally increased. After spring 2010, patterns in achieved

targets were variable. In lakes Alexandrina and Albert, the number of targets achieved remained stable until the last six years when the abundance of several desirable taxa increased such that additional targets were achieved. In Goolwa Channel, the number of targets achieved increased until spring 2011, then decreased due to the increase in abundance of *Typha domingensis* and *Phragmites australis* and a decrease in submergent species. There was an increase in submergents in the deep water zone in autumn 2018 resulting in the target being met for this zone and an increase in the habitat condition score; however, this target was not met in spring 2018 and the habitat condition score decreased but more targets were met in spring 2020 (highest habitat condition score recorded) and autumn 2021 resulting in an increase in habitat condition score. In permanent wetlands, the number of targets achieved has remained constant spring 2010 to spring 2019 after which there was an increase and in temporary wetlands it peaked in autumn 2011 although it has been highly variable. The Whole of Icon Site Score (WOISS) for assessing the condition of the Lower Lakes has remained relatively stable from autumn 2011 to spring 2018 after which there was an increasing trend. The Lakes are now tracking in good condition using the Matter 8 condition scale. Progress towards achievement for most targets has shown they are tracking towards being achieved in the future but still not achieved to date. Therefore, under current hydrological conditions it is likely that the number of targets achieved in the future will further increase (and habitat and WOISS condition will improve) resulting in the TLM Objective V3: *maintain or improve aquatic and littoral vegetation in the Lower Lakes* continuing to be achieved.

Keywords: Lake Alexandrina, Lake Albert, aquatic vegetation.

1. INTRODUCTION

1.1. Background

The Lower Lakes, Coorong and Murray Mouth region is one of six icon sites under “The Living Murray” (TLM) program and is as an indicator site under the “Basin Plan”. The Condition Monitoring Plan for the Lower Lakes, Coorong and Murray Mouth Icon Site (herein referred to as the “icon site”) outlined a series of 17 condition targets for the icon site (Maunsell Australia Pty Ltd 2009). This report includes results from the first 12 years of the understory component of the condition monitoring program designed to evaluate TLM Target V3 (now referred to as objective V3): *maintain or improve aquatic and littoral vegetation in the Lower Lakes* (Marsland and Nicol 2009; Gehrig *et al.* 2010; 2011b; 2012; Frahn *et al.* 2013; 2014; Nicol *et al.* 2016a; 2017; 2019a; b; Nicol *et al.* 2020).

Marsland and Nicol (2006) identified that monitoring programs in existence in 2006 could not adequately assess TLM target V3; therefore, a monitoring program that expanded and built upon the existing programs was established in 2008 (Marsland and Nicol 2009). The understory vegetation monitoring program, described in this report, uses the same methods and sites as the community wetland monitoring program established by the former River Murray Catchment Water Management Board but includes additional sites in lakeshore habitats (in lakes Alexandrina and Albert), the lower reaches of the Finnis River, Currency Creek and Goolwa Channel (herein referred to as Goolwa Channel) and wetlands that were not part of the original program (Marsland and Nicol 2009). In 2009, eight extra sites in Goolwa Channel were added to assess the impact of the Goolwa Channel Water Level Management Project (Gehrig and Nicol 2010a; Gehrig *et al.* 2011a), and data from this project were subsequently included in the TLM Condition Monitoring Program (Gehrig *et al.* 2010; 2011b; 2012; Frahn *et al.* 2013; 2014; Nicol *et al.* 2016a; 2017; 2019a; b; Nicol *et al.* 2020).

The 2009 Condition Monitoring Plan for the Lower Lakes, Coorong and Murray Mouth (LLCMM) Icon Site proposed ‘indicators for monitoring’ that comprised individual taxa and discrete communities: *Melaleuca halmaturorum*, *Myriophyllum* spp. *Gahnia filum*, *Schoenoplectus* spp., *Typha domingensis*, *Phragmites australis* and samphire communities (Maunsell Australia Pty Ltd 2009). However, discussions concluded that the entire understory aquatic and littoral vegetation assemblage would be monitored with a separate technique used for the dominant tree species *Melaleuca halmaturorum* (which is monitored sporadically and not included in this report).

Monitoring aquatic and littoral understorey vegetation involves surveys in spring (high lake levels) and autumn (low lake levels) to determine the current condition, seasonal changes and medium- to long-term changes in floristic composition.

From 1996 to 2010, the Murray-Darling Basin experienced the most severe drought in recorded history (van Dijk *et al.* 2013). Below average stream flows coupled with upstream extraction and river regulation resulted in reduced inflows into South Australia (van Dijk *et al.* 2013), which between January 2007 and August 2010, were insufficient to maintain the pool level downstream of Lock and Weir number 1. Subsequently water levels in lakes Alexandrina and Albert dropped to unprecedented lows (<-0.75 m AHD), fringing wetlands became disconnected and desiccated and extensive areas of acid sulfate soils were exposed; particularly in Lake Albert and the lower reaches of the Finniss River and Currency Creek (Merry *et al.* 2003; Fitzpatrick *et al.* 2009a; 2009b; 2010; 2011).

Prior to 2007, fringing wetlands in the Lower Lakes region contained diverse communities of emergent, amphibious and submergent taxa (Renfrey *et al.* 1989; Holt *et al.* 2005; Nicol *et al.* 2006). For example, in 2004, *Ruppia polycarpa*, *Althenia* (formerly named *Lepilaena*) sp., *Nitella* sp. and *Myriophyllum* sp. were common in Narrung Wetland; *Myriophyllum salsgineum* and *Vallisneria australis* were common in Dunn's Lagoon; *Ruppia polycarpa*, *Ruppia tuberosa*, *Myriophyllum* sp. and *Potamogeton pectinatus* were common in Teringie Wetland and *Myriophyllum caput-medusae* was common in Shadows Lagoon and Boggy Creek (Holt *et al.* 2005). Furthermore, in 2005, *Ranunculus trichophyllus*, *Vallisneria australis* and *Myriophyllum caput-medusae* were common in Pelican Lagoon; *Ruppia polycarpa* was common in Point Sturt Wetland; *Ruppia tuberosa* and *Myriophyllum caput-medusae* were common in Poltalloch; *Ranunculus trichophyllus* and *Ruppia polycarpa* were common in Loveday Bay Wetland (Jenny's Lagoon) and *Myriophyllum caput-medusae*, *Myriophyllum salsgineum*, *Ruppia megacarpa*, *Ruppia tuberosa* and *Potamogeton pectinatus* were common in Hunters Creek (Nicol *et al.* 2006).

By spring 2008, submergent taxa had been extirpated (except for a small number of *Ruppia tuberosa* plants in Hunters Creek, in Lake Alexandrina near Raukkan and in Loveday Bay Wetland). The charophyte *Lamprothamnium macropogon* was also present in Loveday Bay Wetland. Amphibious taxa had declined in abundance and diversity, stands of emergent taxa were disconnected from remaining water and fringing habitats were dominated by terrestrial taxa and bare soil (Marsland and Nicol 2009). Furthermore, submergent taxa had not colonised the remaining open water areas (Marsland and Nicol 2009).

The loss of submergent vegetation, decline in abundance and diversity of amphibious taxa and disconnection of fringing emergent macrophytes had serious implications for ecosystem dynamics of the Lower Lakes. This is because aquatic vegetation is a critical ecosystem component in the Lower Lakes; plants are major primary producers (e.g. dos Santos and Esteves 2002; Camargo *et al.* 2006; Noges *et al.* 2010), improve water quality (e.g. Webster *et al.* 2001; James *et al.* 2004), provide habitat for invertebrates (e.g. Wright *et al.* 2002; Papas 2007; Bassett *et al.* 2012; Bell *et al.* 2013; Walker *et al.* 2013; Matuszak *et al.* 2014), birds (e.g. Brandle *et al.* 2002; Phillips and Muller 2006) and threatened fish (Wedderburn *et al.* 2007; Bice *et al.* 2008) and stabilise shorelines (Abernethy and Rutherford 1998; PIRSA Spatial Information Services 2009).

To mitigate impacts of acid sulfate soils, three regulators were constructed in the Lower Lakes: the Narrung Bund (completed in early 2008), the Clayton Regulator and the Currency Creek Regulator (both completed in August 2009) (Figure 1). However, only the impacts of the Narrung Bund and Clayton Regulator will be discussed in this report due to the Currency Creek Regulator spillway remaining inundated after the Clayton regulator was constructed. The regulators disconnected Goolwa Channel and Lake Albert from Lake Alexandrina, which enabled water levels within each site to be managed independently. An additional hydrological intervention was undertaken at Narrung Wetland, with 250 megalitres (ML) of water for the environment from Lake Alexandrina being pumped into the wetland in October 2009 to provide suitable conditions for the growth of submergent taxa (particularly *Ruppia tuberosa* and charophytes).

In August 2010, flows into South Australia increased, and as a result water levels in Lake Alexandrina were reinstated to historical levels (~+0.75 m AHD) and significant flow through the Murray Barrages (five flow control structures located at Goolwa, Tauwichee, Ewe Island, Boundary Creek and Mundoo to prevent saltwater intrusion in the Lower Lakes; Figure 1) was possible for the first time since spring 2005 (although there was a small water release in 2006-07 to operate fishways). Furthermore, the Clayton Regulator and Narrung Bund were breached in September 2010, and Lake Alexandrina was reconnected with Goolwa Channel and Lake Albert. After spring 2010, water levels were restored to historical levels ranging from +0.9 m AHD in spring when lakes were surcharged to +0.4 m AHD in autumn during periods of managed draw-down. The impacts of the regulators, pumping, unregulated River Murray flows and managed draw down on salinity and water levels throughout the condition monitoring program are outlined in section 2.1.

The period of low flow and subsequent low water levels, regulator construction, pumping, unregulated River Murray flows, regulator breaching, entitlement flows and managed draw-down have resulted in large changes to the hydrological and salinity regime of the Lower Lakes since 2007. Salinity (e.g. Hart *et al.* 1991; Nielsen *et al.* 2003; Nielsen *et al.* 2007; Nielsen and Brock 2009) and water regime (determined by lake levels) (e.g. Brock and Casanova 1997; Blanch *et al.* 1999b; 1999a; 2000; Nicol *et al.* 2003) are two of the primary drivers of plant community composition in freshwater ecosystems. Historically, the various components of the system were connected with relatively stable water levels ranging from +0.4 to +0.8 m AHD and surface water electrical conductivity lower than 2,000 $\mu\text{S}\cdot\text{cm}^{-1}$ (Kingsford *et al.* 2009; Kingsford *et al.* 2011). Between January 2007 and August 2010, surface water salinity, water regime and connectivity of the study area varied dramatically from historical patterns; however, since September 2010, these factors have largely reflected historical patterns.

1.2. Aquatic and littoral vegetation target revision

A review undertaken by Robinson (2015) suggested that the initial aquatic and littoral vegetation target for the Lower Lakes (TLM V3): *maintain or improve aquatic and littoral vegetation in the Lower Lakes* (Maunsell Australia Pty Ltd 2009) could be improved by developing a series of quantitative targets for the site. In response to this, targets were developed for the aquatic and littoral vegetation of the Lower Lakes. The targets were based largely on expert opinion; however, pre-drought vegetation information was available for wetlands through the 2004 (Holt *et al.* 2005) and 2005 (Nicol *et al.* 2006) River Murray wetlands baseline surveys, biological surveys of conservation reserves around the Murray Mouth (Brandle *et al.* 2002), habitat mapping for the entire system (Seaman 2003) and for Hindmarsh Island (Renfrey *et al.* 1989). Generally, these studies showed there was a diverse submergent, emergent and amphibious plant community in wetlands, along low energy shorelines in lakes Alexandrina and Albert and in aquatic habitats, on and around Hindmarsh Island, prior to 2007. Whilst these studies represent the only documented baseline (prior to 2007) for the Lower Lakes, they were snapshots that did not provide an indication of temporal variability. The updated quantitative targets and methodologies are outlined in the Condition Monitoring Plan (Revised) 2017 for the Lower Lakes, Coorong and Murray Mouth Icon Site (Department for Environment, Water and Natural Resources 2017). Within this Plan, the original target V3 is now referred to as an objective, with the quantitative targets nested below the objective.

The vegetation condition monitoring review divided the Lower Lakes into different habitats based on hydrology and geomorphology. Five habitats were identified: Lake Alexandrina, Lake Albert, Goolwa Channel, permanent wetlands and seasonal (temporary) wetlands. Within lakes Alexandrina and Albert and Goolwa Channel, three zones were identified based on elevation: the littoral zone (+0.8 to +0.6 m AHD), the aquatic zone (+0.4 to 0 m AHD) and the deep water zone (deeper than 0 m AHD). Permanent wetlands are typically shallow and have no deep water zone; hence they were divided into littoral and aquatic zones. Seasonal wetlands were divided into two zones: the wetland edge and wetland bed. In addition, there was a seasonal component for temporary wetlands with targets for spring (high water level) and autumn (low water level).

Due to the number of plant species present in the Lower Lakes, native species were classified into functional groups based on water regime using the classification in Gehrig and Nicol (2010; Appendix 1).

Exotic species and potentially invasive native species (e.g. *Typha domingensis* and *Phragmites australis*) were also monitored. The dominant exotic species in the Lower Lakes are *Cenchrus clandestinus* (formerly named *Pennisetum clandestinum*) and *Paspalum distichum* (Frahn *et al.* 2014). Both are low profile rhizomatous and stoloniferous, warm season growing grasses (Jessop *et al.* 2006) that grow well in the littoral zone throughout the Lower Lakes, except in areas where there is high soil salinity (Frahn *et al.* 2014). Native emergent and amphibious species are often absent when these species are abundant (Frahn *et al.* 2014). *Typha domingensis* and *Phragmites australis* are tall rhizomatous emergent species that are common throughout the Lower Lakes (Frahn *et al.* 2014) and are adapted to stable water levels (Blanch *et al.* 1999b; 2000). They are an important component of the vegetation in the Lower Lakes; however, they often form monospecific stands and it is undesirable for these species to occupy large areas of the littoral and aquatic zones.

Targets for aquatic and littoral understorey vegetation were based on a minimum proportion of quadrats in each habitat and zone having a minimum cover score of desirable species and a maximum number of quadrats having a maximum cover score of undesirable species in any given survey. Species were classified into water regime functional groups to assess targets except the undesirable species: *Paspalum distichum*, *Cenchrus clandestinus*, *Phragmites australis* and *Typha domingensis*.

Vegetation targets for Lake Alexandrina are presented in Table 1. The general objectives of the targets were to improve the abundance of diverse reed beds (shorelines with a diverse assemblage of emergent, submergent and amphibious species) and limit the amount of shoreline dominated by invasive species and to a lesser extent shorelines dominated by *Typha domingensis* and *Phragmites australis*. The deep water zone in Lake Alexandrina is generally unsuitable for submergent or emergent species; hence, there were no vegetation targets for this zone, but it was recognised that this zone needs to be inundated to maintain the hydrological connection between zones and prevent acid sulfate soil development (Fitzpatrick *et al.* 2009a; 2009b; 2010).

Table 1: Revised vegetation targets for Lake Alexandrina.

Zone	Target
Littoral +0.8 to +0.6 m AHD	<40% of quadrats in any given survey containing >75% combined cover (Braun-Blanquet score 5) of <i>Typha</i> and <i>Phragmites</i>
	<20% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Cenchrus</i> and <i>Paspalum</i>
	Minimum of 50% of quadrats in any given survey contain native amphibious species with a combined cover of $\geq 5\%$ (BB score 2 or greater)
	Minimum of 50% of quadrats in any given survey contain native emergent species (other than <i>Typha</i> and <i>Phragmites</i>) with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
Aquatic +0.4 to 0 m AHD	<40% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Typha</i> and <i>Phragmites</i>
	Minimum of 20% of quadrats in any given survey contain native emergent species (other than <i>Typha</i> and <i>Phragmites</i>) with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
	Minimum of 35% of quadrats in any given survey contain native submergent species with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
Deep Water <0 m AHD	Permanent inundation

Targets for Lake Albert are presented in Table 2. The targets for Lake Albert were similar to those for Lake Alexandrina except that there was an expectation of a lower proportion of diverse reed beds and lower proportions of submergent, amphibious and emergent species (except *Typha domingensis* and *Phragmites australis*).

Table 2: Revised vegetation targets for Lake Albert.

Zone	Target
Littoral +0.8 to +0.6 m AHD	<40% of quadrats in any given survey containing >75% combined cover (Braun-Blanquet score 5 or greater) of <i>Typha</i> and <i>Phragmites</i>
	<20% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Cenchrus</i> and <i>Paspalum</i>
	Minimum of 35% of quadrats in any given survey contain native amphibious species with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
	Minimum of 35% of quadrats in any given survey contain native emergent species (other than <i>Typha</i> and <i>Phragmites</i>) with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
Aquatic +0.4 to 0 m AHD	<40% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Typha</i> and <i>Phragmites</i>
	Minimum of 20% of quadrats in any given survey contain emergent species (other than <i>Typha</i> and <i>Phragmites</i>) with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
	Minimum of 20% of quadrats in any given survey contain submergent species with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
Deep Water <0 m AHD	Permanent inundation

Targets for Goolwa Channel are presented in Table 3. Targets for Goolwa Channel were also similar to Lake Alexandrina but there was an expectation that submergent species were present in the deep water zone and a higher proportion of quadrats dominated by *Typha domingensis* and *Phragmites australis*.

Table 3: Revised vegetation targets for Goolwa Channel.

Zone	Target
Littoral +0.8 to +0.6 m AHD	<50% of quadrats in any given survey containing >75% combined cover (Braun-Blanquet score 5 or greater) of <i>Typha</i> and <i>Phragmites</i>
	<20% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Cenchrus</i> and <i>Paspalum</i>
	Minimum of 50% of quadrats in any given survey contain native amphibious species with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
	Minimum of 50% of quadrats in any given survey contain native emergent species (other than <i>Typha</i> and <i>Phragmites</i>) with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
Aquatic +0.4 to 0 m AHD	<50% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Typha</i> and <i>Phragmites</i>
	Minimum of 20% of quadrats in any given survey contain native emergent species (other than <i>Typha</i> and <i>Phragmites</i>) with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
	Minimum of 40% of quadrats in any given survey contain native submergent species with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)
Deep Water <0 m AHD	Minimum of 20% of quadrats in any given survey contain native submergent species with a combined cover of $\geq 5\%$ (Braun-Blanquet score 2 or greater)

Targets for permanent wetlands are presented in Table 4. Prior to 2007, many wetlands contained a diverse assemblage of submergent, emergent and amphibious species (Holt *et al.* 2005; Nicol *et al.* 2006), which was reflected in the targets. The proportion of quadrats dominated by *Typha domingensis* and *Phragmites australis* is lower than Goolwa Channel, Lake Alexandrina and Lake Albert and proportion of quadrats with submergents is higher (Table 5). However, there is a maximum target of 50% cover for submergent species in the aquatic zone, which was related to small bodied fish habitat (S. Wedderburn pers. com.). The deep water zone is not included because wetlands are generally shallow and this zone is not present in most wetlands.

Table 4: Revised vegetation targets for permanent wetlands (Dunns Lagoon, Hunters Creek, Angas River Mouth and Bremer River Mouth).

Zone	Target
Littoral >+0.6 m AHD	<35% of quadrats in any given survey containing >75% combined cover (Braun-Blanquet score 5 or greater) of <i>Typha</i> and <i>Phragmites</i>
	<20% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Cenchrus</i> and <i>Paspalum</i>
	Minimum of 50% of quadrats in any given survey contain native amphibious species with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)
	Minimum of 50% of quadrats in any given survey contain native emergent species (other than <i>Typha</i> and <i>Phragmites</i>) with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)
Aquatic <+0.6 m AHD	<40% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Typha</i> and <i>Phragmites</i>
	Minimum of 20% of quadrats in any given survey contain native emergent species (other than <i>Typha</i> and <i>Phragmites</i>) with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)
	Minimum of 50% of quadrats in any given survey contain native submergent species with a combined cover of 5 to 50% (Braun-Blanquet score 2 to 4)

Targets for seasonal wetlands are presented in Tables 5a and b. Prior to 2007, seasonal wetlands in spring generally contained high numbers of submergent species (submergent r-selected species (*sensu* Casanova 2011) such as *Ruppia tuberosa*, *Ruppia polycarpa*, *Althenia cylindrocarpa* and charophytes) (Holt *et al.* 2005; Nicol *et al.* 2006). This is reflected in the spring target of 50% of quadrats containing greater than 25% cover of submergent species (Table 5a) because the regular wetting and drying cycle present in these wetlands favours this functional group (Casanova 2011). Furthermore, the wetting and drying cycle will favour amphibious species that require exposed sediment to germinate but persist as adults whilst standing water is present (Nicol *et al.* 2003; Casanova 2011). *Typha domingensis* and *Phragmites australis* are generally not abundant in seasonal wetlands (Frahn *et al.* 2014); hence, there were no targets relating to these species.

Table 5: Revised vegetation targets for seasonal wetlands in a. spring and b. autumn (Goolwa Channel Drive, Milang Wetland, Narrung Wetland, Loveday Bay Wetland, Point Sturt Wetland and Teringie).

a.

Zone	Target
Edge	<20% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Cenchrus</i> and <i>Paspalum</i>
	Minimum of 50% of quadrats in any given survey contain native amphibious species with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)
	Minimum of 50% of quadrats in any given survey contain native emergent species with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)
Bed	Minimum of 20% of quadrats in any given survey contain native emergent species with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)
	Minimum of 50% of quadrats in any given survey contain native submergent species with a combined cover of ≥25% (Braun-Blanquet score 3 or greater)
	Minimum of 25% of quadrats in any given survey contain native amphibious species with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)

b.

Zone	Target
Edge	<20% of quadrats in any given survey containing >50% combined cover (Braun-Blanquet score 4 or greater) of <i>Cenchrus</i> and <i>Paspalum</i>
	Minimum of 50% of quadrats in any given survey contain native amphibious species with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)
	Minimum of 50% of quadrats in any given survey contain native emergent species with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)
Bed	Minimum of 20% of quadrats in any given survey contain native emergent species with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)
	Minimum of 25% of quadrats in any given survey contain native amphibious species with a combined cover of ≥5% (Braun-Blanquet score 2 or greater)

In addition to quantitative aquatic and littoral vegetation targets, habitat and whole of icon site scores (WOISS) were developed to assess the condition of the lakes. The habitat condition score represents the proportion of targets achieved in a particular habitat and the WOISS represents the proportion of targets achieved in the different habitats. The WOISS is also used for Basin Plan environmental outcome reporting (Matter 8) and will be used to report on objective V3 using the same scale (Table 6) (DEW in prep.). A Matter 8 condition rating of good represents aquatic and littoral vegetation being maintained and a rating of very good represents condition is improving.

Table 6: Icon site scores for aquatic and littoral vegetation for the Lower Lakes with the condition rating used in Matter 8 Report Cards (Department for Environment and Water 2019).

Icon site score	Matter 8 condition rating
0.80-1.00	Very good
0.60-0.79	Good
0.40-0.59	Fair
<0.40	Poor

1.3. Objectives

The surveys undertaken in spring 2020 and autumn 2021 builds on data collected between spring 2008 and spring 2019 and provides information regarding the change in plant communities over this period. However, in spring 2016 surveys were only undertaken in seasonal wetland habitats with all sites surveyed in autumn and spring 2017. From autumn 2018 onwards, surveys were not undertaken in Milang and Waltowa wetlands.

The monitoring program includes a period of record low water levels in Lake Alexandrina, several engineering interventions, two large unregulated River Murray flow events (one in 2010/11 that reinstated historical water levels and one in spring/summer 2016), two in-channel flow pulses, entitlement flows, water for the environment provisions (that maintained historical water levels), managed draw-down to +0.5 m AHD in late summer and autumn 2018 and surcharge in spring 2019 to +0.9 m AHD. Therefore, this monitoring program collected information regarding the change in aquatic and littoral plant communities in response to draw-down, desiccation, increased water levels due to regulated inundation, natural flooding, spring surcharging of the lakes and managed drawn-down, and provides an insight into recovery of the system under hydrological restoration. The aims of this project are to:

- continue the statistically robust, quantitative understorey aquatic and littoral vegetation monitoring program in the Lower Lakes to assess TLM Objective V3 (Department for Environment, Water and Natural Resources 2017);
- report on the revised vegetation targets for each habitat and determine habitat condition using the WOISS ;
- monitor the recovery of the aquatic plant community after hydrological restoration following extended drought, draw-down, fragmentation and desiccation of aquatic habitats;
- investigate the longevity of the managed draw-down in late summer and autumn 2018.

2. METHODS

2.1. Study site, hydrology and salinity

Vegetation surveys were undertaken in the Goolwa Channel, Lake Alexandrina, Lake Albert and 11 associated wetlands (Figure 1). Between 2008 and 2010, a range of interventions were undertaken in the Lower Lakes to regulate water levels and mitigate acid sulfate soils; primarily the construction of the Narrung Bund and Clayton Regulator (Figure 1). Construction of the Narrung Bund was completed in early 2008 and this disconnected Lake Albert from Lake Alexandrina (Figure 1). Water was then pumped from Lake Alexandrina into Lake Albert to maintain water levels above -0.5 m AHD. Construction of the Clayton Regulator was completed in August 2009, resulting in impounded water from the Finniss River and Currency and Tookayerta Creeks (Figure 1). In addition, water was pumped into Goolwa Channel (Figure 2) from Lake Alexandrina to raise water levels to +0.7 m AHD in spring 2009. Both structures were breached in spring 2010, and from then on water levels were dependent on inflows and barrage operations. Water level and surface water electrical conductivity in the Lower Lakes from August 2008 to April 2020 are presented in Figure 2 and Figure 3 respectively. Details regarding interventions and their impacts on water level and salinity from 2008 to 2010 are outlined in Frahn *et al.* (2014).

Since the Clayton Regulator and Narrung Bund were breached in spring 2010, water levels in the Lower Lakes returned to historical levels and remained at these levels for the remainder of the survey period (Figure 2). Salinity in Lake Alexandrina and Goolwa Channel decreased rapidly after the Clayton Regulator and Narrung Bund were breached; however, salinity remains elevated (but slowly decreasing) in Lake Albert and there have been several short salinity spikes in Goolwa Channel during periods of reverse head (the water level in the Coorong is higher than Lake Alexandrina) (Figure 3).

Since 2011/12, water level management objectives for the Lower Lakes have focused on annual fluctuations within a range from +0.4 to +0.9 m AHD to achieve ecological benefits. In late summer and autumn 2018 water levels were drawn-down to around +0.5 m AHD for the longest period since the Millennium Drought. Water levels were also drawn down to a similar level in autumn 2019 and surcharged to +0.9 m AHD in spring 2019 with a similar pattern in autumn 2019 and spring 2020. However, in autumn 2021 water levels were drawn-down to around +0.6 m AHD (Figure 2). It is expected that these management actions will promote the establishment of

amphibious species in the littoral zone. Figure 2 indicates that lakes levels have generally followed this pattern with the exception of water levels reaching maximums of > 0.9 m AHD in spring 2016, 2019 and 2020.




Figure 1: Map of Lakes Alexandrina and Albert and Goolwa Channel showing the location of lakeshore and wetland vegetation monitoring sites (site numbers correspond to Table 7) and major flow control structures present in winter 2010 (where sites are in close proximity they may not be visible on map).




Figure 2: Daily mean water levels in Goolwa Channel (Signal Point), Lake Alexandrina (Milang) and Lake Albert (Meningie) from August 2008 to May 2021 (Department of Environment and Water 2021b).




Figure 3: Daily mean surface water electrical conductivity (EC) in Goolwa Channel (Signal Point), Lake Alexandrina (Milang) and Lake Albert (Meningie) from August 2008 to May 2021 (Department of Environment and Water 2021a).

2.2. Understorey vegetation survey protocol

Monitoring of understorey vegetation was conducted at 11 wetland and 25 lakeshore sites each spring and autumn from October 2008 to March 2014, March 2015, October 2015, March 2016, December 2016 (temporary wetlands only), March 2017, November 2017, April 2018 (except Milang and Waltowa wetlands), November 2018, November 2019, December 2020 and May 2021 (Table 7). Sites were grouped based on habitat (lakeshore, permanent wetland or seasonal wetland) and location (Lake Alexandrina, Lake Albert or Goolwa Channel). GPS coordinates for each site are listed in Appendix 2.

Table 7: List of understorey vegetation site numbers (relative to map provided in Figure 1), site name, location, habitat type (wetland or lakeshore), number of survey sites and the year sites were established (SAMDBNRM denotes, South Australian Murray-Darling Basin Natural Resources Management Board).

Site #	Site Name	Location	Habitat	No. Survey Sites	Year Established
1	Bremer Mouth Lakeshore	Lake Alexandrina	lakeshore	1	2008
2	Brown Beach 1	Lake Albert	lakeshore	1	2008
3	Brown Beach 2	Lake Albert	lakeshore	1	2008
4	Clayton Bay	Goolwa Channel	lakeshore	1	2009
5	Currency Creek 3	Goolwa Channel	lakeshore	1	2008
6	Currency Creek 4	Goolwa Channel	lakeshore	1	2008
7	Goolwa North	Goolwa Channel	lakeshore	1	2009
8	Goolwa South	Goolwa Channel	lakeshore	1	2009
9	Hindmarsh Island Bridge 01	Goolwa Channel	lakeshore	1	2009
10	Hindmarsh Island Bridge 02	Goolwa Channel	lakeshore	1	2009
11	Lake Reserve Rd	Lake Alexandrina	lakeshore	1	2008
12	Loveday Bay	Lake Alexandrina	seasonal wetland	4	2009
13	Loveday Bay Lakeshore	Lake Alexandrina	lakeshore	1	2009
14	Lower Finniss 02	Goolwa Channel	lakeshore	1	2009
15	Milang (existing SAMDBNRM Board community monitoring site)	Lake Alexandrina	seasonal wetland	4	pre-2008
16	Milang Lakeshore	Lake Alexandrina	lakeshore	1	2009
17	Pt Sturt Lakeshore	Lake Alexandrina	lakeshore	1	2008
18	Pt Sturt Water Reserve	Lake Alexandrina	lakeshore	1	2008
19	Teringie Lakeshore	Lake Alexandrina	lakeshore	1	2008
20	Upstream of Clayton Regulator	Lake Alexandrina	lakeshore	1	2009
21	Wally's Landing	Goolwa Channel	lakeshore	1	2009
22	Warrenjie 1	Lake Albert	lakeshore	1	2009
23	Lower Finniss 03	Goolwa Channel	lakeshore	1	2009
24	Narrung Lakeshore	Lake Alexandrina	lakeshore	1	2008
25	Nurra Nurra	Lake Albert	lakeshore	1	2008
26	Warrenjie 2	Lake Albert	lakeshore	1	2009
27	Angas Mouth	Lake Alexandrina	permanent wetland	1	2008
28	Bremer Mouth	Lake Alexandrina	permanent wetland	1	2008
29	Dunns Lagoon	Lake Alexandrina	permanent wetland	4	2008
30	Goolwa Channel Drive	Lake Alexandrina	seasonal wetland	3	2008
31	Hunters Creek	Lake Alexandrina	wetland	5	2008
32	Poltalloch	Lake Alexandrina	seasonal wetland	2	2008
33	Pt Sturt	Lake Alexandrina	seasonal wetland	2	2008
34	Teringie (existing SAMDBNRM Board community monitoring site)	Lake Alexandrina	seasonal wetland	4	pre-2008
35	Waltowa (existing SAMDBNRM Board community monitoring site)	Lake Albert	seasonal wetland	2	pre-2008
36	Narrung (existing SAMDBNRM Board community monitoring site)	Lake Alexandrina	seasonal wetland	4	pre-2008

Wetlands

At each survey site (Figure 1, Table 7), a transect running perpendicular to the shoreline and three, 1 x 3 m quadrats, separated by one metre, were established (Figure 4) at regular elevation intervals that represented the dominant plant communities (A. Rumbelow pers. comm.). In wetlands with an established monitoring program (Milang, Waltowa, Teringie and Narrung), existing sites were re-surveyed. For the remaining wetlands (Dunns Lagoon, Point Sturt, Hunters Creek, Goolwa Channel Drive, Bremer River Mouth, Angas River Mouth and Loveday Bay), a transect was established and quadrats placed in each plant community present during the spring 2008 survey. A minimum of one additional transect (but usually two or more in each wetland, except at the Angas and Bremer River mouths) was established, and quadrats were placed at the same elevations (determined using a laser level) as on the first transect. At sites where the elevation gradient was steep (e.g. Angas and Bremer River Mouth, Hunter's Creek) only edge and channel quadrats were surveyed. Cover and abundance of each species present in the quadrat were estimated using the method outlined in Heard and Channon (1997), except that N and T were replaced by 0.1 and 0.5 to enable statistical analyses (Table 8).

Table 8: Modified Braun-Blanquet (1932) scale estimating cover/abundance as per Heard and Channon (1997).

Score	Modified Score	Description
N	0.1	Not many, 1-10 individuals
T	0.5	Sparsely or very sparsely present; cover very small (less than 5%)
1	1	Plentiful but of small cover (less than 5%)
2	2	Any number of individuals covering 5-25% of the area
3	3	Any number of individuals covering 25-50% of the area
4	4	Any number of individuals covering 50-75% of the area
5	5	Covering more than 75% of the area

Lakeshores

With the exception of quadrat placement, lakeshores were surveyed using the same technique as wetlands. At each site, a transect running perpendicular to the shoreline was established and three, 1 x 3 m quadrats, separated by one metre, were established at elevation intervals of +0.8, +0.6, +0.4, +0.2, 0 and -0.5 m AHD (Figure 4) (*sensu* Marsland and Nicol 2009; Gehrig and Nicol 2010a; Gehrig *et al.* 2010).



Figure 4: Vegetation surveying protocol for lakeshore sites: plan view showing placement of quadrats relative to the shoreline.

Plant identification and Nomenclature

Plants were identified using keys in Sainty and Jacobs (1981), Jessop and Tolken (1986), Prescott (1988), Cunningham *et al.* (1992), Dashorst and Jessop (1998), Romanowski (1998), Sainty and Jacobs (2003) and Jessop *et al.* (2006). In some cases, due to immature individuals or lack of floral structures, plants were identified to genus only. Nomenclature follows the Centre for Australian National Biodiversity Research and Council of Heads of Australasian Herbaria (2021).

2.3. Data Analysis

Changes in floristic composition through time, at all elevations, in each of the five habitats (Lake Alexandrina, Lake Albert, Goolwa Channel, permanent wetlands and temporary wetlands) were assessed by non-metric multidimensional scaling (nMDS) ordination using the package PRIMER version 7.0.12 (Clarke and Gorley 2015). Bray-Curtis (1957) similarities were used to construct the similarity matrices for the nMDS ordinations on untransformed data.

Native species richness at +0.6 m AHD in Lakes Alexandrina, Lake Albert and Goolwa Channel was plotted through time to assess the benefit of the managed draw down in autumn 2018. This elevation was chosen because it represented the zone that was inundated in spring and exposed by the managed draw-down in autumn 2018, 2019 and 2020.

3. RESULTS

3.1. Change through time of the Lower Lakes plant community from spring 2008 to autumn 2021

Over the 12 years of condition monitoring (spring 2008 to autumn 2021), a total of 164 taxa (including 77 exotics, two weeds of national significance, five proclaimed pest plants in South Australia and one species listed as rare in South Australia) were recorded throughout the Lower Lakes (Appendix 1). Species list of each habitat (Lake Alexandrina, Lake Albert, Goolwa Channel, permanent wetlands and temporary wetlands) and the surveys they were recorded are presented in Appendices 3 to 7). Lake Alexandrina was the most species rich of the habitats with 120 taxa (including 56 exotics) recorded between spring 2008 and autumn 2021, followed by permanent wetlands (98 taxa, including 36 exotics), then temporary wetlands (95 taxa, including 43 exotics), Goolwa Channel (80 taxa, including 30 exotics) and Lake Albert the least species rich with 61 taxa (including 32 exotics).

Patterns of temporal change in the plant community for each habitat showed a shift in floristic composition over the condition monitoring program (nMDS ordination; Figures 5 to 9). Furthermore, for each habitat except the seasonal wetlands, there was greater change in the plant community between the early surveys that reduced through time with very little change in vegetation between the more recent surveys (Figures 5 to 9).

Lake Alexandrina

In spring 2008, water levels in Lake Alexandrina were at historical lows (Figure 2) and the plant community was dominated by terrestrial species (predominantly agricultural weeds). The plant community remained dominated by terrestrial taxa until spring 2010, when water levels were reinstated (Figure 2) and the terrestrial species were extirpated resulting in a large change in floristic composition (Figure 5). From spring 2010 to autumn 2013, there was an increase in the abundance of emergent, amphibious and submergent species (Frahm *et al.* 2014). However, there were also seasonal patterns over this period (Figure 5) with emergent taxa typically more abundant in spring and amphibious taxa in autumn (Frahm *et al.* 2014). From autumn 2013 to spring 2017 the change in the plant community has been small, in comparison to previous years, but seasonal patterns were similar (Figure 5) with emergent taxa more abundant in spring and amphibious and submergent taxa in autumn. There was a small shift in the plant community between spring 2017 and autumn 2018 due to an increase in species richness in the littoral zone and less change between autumn and spring 2018 (compared to previous years) (Figure 5) as most of these species persisted. Between spring 2018 and 2019 there was also a small change in the vegetation with the community becoming more similar to the communities present in autumn 2016 and 2017 (Figure 5). There was a larger change in the plant community between spring 2019 and 2020, compared to recent years, which was due to an increase in submergent species in the aquatic zone and amphibious species in the littoral zone (in particular at +0.8 m AHD) (Figure 5).




Figure 5: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in Lake Alexandrina (Sp denotes spring; Au denotes autumn).

Lake Albert

Similar to Lake Alexandrina (Figure 5), the plant community in Lake Albert was dominated by terrestrial species with large changes in floristic composition (Figure 6) whilst water levels were low during the Millennium Drought (Figure 2). After water levels were reinstated in spring 2010 (Figure 2), the change in plant community (Figure 6) was driven by an increase in emergent and amphibious species. Since spring 2010, there has been a gradual change in floristic composition (Figure 6) primarily driven by an increase in the abundance of *Typha domingensis*. In contrast to Lake Alexandrina, seasonal changes in floristic composition were not evident and there was little change between autumn 2017 and spring 2019 (Figure 6). However, between spring 2019 and spring 2020 there was a comparatively large change in the plant community driven by decreases in abundance of native amphibious species and *Paspalum distichum* and *Cenchrus clandestinus* (Figure 6).




Figure 6: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in Lake Albert (Sp denotes spring; Au denotes autumn).

Goolwa Channel

Similar to lakes Alexandrina (Figure 5) and Albert (Figure 6), the plant community in Goolwa Channel was dominated by terrestrial taxa whilst water levels were low prior to spring 2009 (Figure 2). Water levels rose to around +0.8 m AHD in spring 2009 (Figure 2) due to the completion of the Clayton Regulator and there was a large change in floristic composition (Figure 7). This change was driven by terrestrial species being extirpated with extensive beds of the submergent species *Potamogeton pectinatus* recruiting throughout Goolwa Channel, the lower Finniss River and Lower Currency Creek (Gehrig and Nicol 2010a). There was a significant change in the plant community between spring of 2009 and 2010 (Figure 7), which was a result of the Clayton Regulator being breached and a rapid reduction in surface water salinity (Figure 3). These changes in floristic composition were driven by a decrease in the abundance of *Potamogeton pectinatus* and increase in submergent species adapted to lower salinity environments (e.g. *Ceratophyllum demersum*, *Potamogeton crispus*, *Myriophyllum salsugineum* and *Vallisneria australis*) (Bailey *et al.* 2002). After spring 2010, water levels and salinities returned to historical levels (Figure 2 and Figure 3) but the plant community continued to change (Figure 7). The change between spring 2010 and spring 2011 (Figure 7) was driven primarily by an increase in the abundance of *Typha domingensis*. There were seasonal changes in vegetation between spring 2011 spring 2013 (Figure 7) driven by higher abundances of *Typha domingensis* and *Phragmites australis* in autumn. After spring 2013, there was very little change in floristic composition and until autumn 2018 when there was an increase in amphibious taxa in the littoral zone and submergents in the deep-water zone with little change since then (Figure 7).




Figure 7: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in Goolwa Channel (Sp denotes spring; Au denotes autumn).

Permanent Wetlands

All permanent wetlands surveyed in the condition monitoring program are hydrologically connected to Lake Alexandrina; therefore, water levels (and salinities to a lesser degree) in these habitats reflect conditions in Lake Alexandrina (Figures 2 and 3). Similar to the other habitats in the Lower Lakes, permanent wetlands were dominated by terrestrial taxa whilst water levels were low, most of which were extirpated when water levels were reinstated in spring 2010 (Figure 2). Since spring 2010, there has generally been an increase in the abundance of emergent, submergent and amphibious species in permanent wetlands, which has driven the change in floristic composition (Figure 8). Since autumn 2013, the change in the plant community was much smaller than observed in the earlier surveys of the condition monitoring program (Figure 7). Similar to lakes Alexandrina and Albert, there was little change in floristic composition between autumn 2018 and spring 2019 but a larger shift between spring 2019 and spring 2020 (Figure 7). The recent change was driven by an increase in submergent species in the aquatic zone and amphibious taxa in the littoral zone.




Figure 8: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in permanent wetlands (Sp denotes spring; Au denotes autumn).

Temporary Wetlands

Strong seasonal changes in vegetation are evident in the seasonal wetlands that were surveyed (Figure 9). Despite lack of hydrological connectivity to the Lower Lakes between spring 2008 and autumn 2010, all wetlands were partially inundated in spring 2008 and spring 2009 due to local rainfall and runoff hence the seasonal patterns in floristic composition during this period (Figure 7). The submergent species *Ruppia tuberosa* and *Lamprothamnium macropogon* were present in the inundated areas of several of the seasonal wetlands in spring 2008 and spring 2009 and absent in autumn 2009 and 2010 when the wetlands were dominated by terrestrial taxa. After water levels were reinstated in spring 2010 and the hydrological connection with the lakes restored, in contrast to the other habitats, the plant community was more similar to the community present in spring 2009 than in spring 2011 (Figure 9). There was; however, a change between spring 2010 and autumn 2011, after which there was very little change in floristic composition between autumn surveys (Figure 7). The change was driven by an increase in the abundance of *Typha domingensis*, *Bolboschoenus caldwellii* and *Schoenoplectus pungens* between spring 2010 and autumn 2011. The seasonal patterns observed between autumn 2011 and autumn 2021 (Figure 7) were due to the presence of submergent species (*Ruppia tuberosa*, *Ruppia polycarpa*, *Myriophyllum verrucosum*, *Myriophyllum salsugineum*, *Chara* sp. and *Lamprothamnium macropogon*) in spring.





Figure 9: nMDS ordination comparing the plant community between spring 2008 and autumn 2021 in seasonal wetlands (Sp denotes spring; Au denotes autumn).

Effect of variable water levels

Managed water level cycling for ecological outcomes has been undertaken in the Lower Lakes with spring surcharging (when water is available) and draw-down in late summer/autumn being undertaken between autumn 2018 and 2020 (Figure 2). This has resulted in quadrats at +0.6 m AHD being subjected to repeated patterns of inundation and exposure (Figure 2). Native amphibious species richness in Lake Alexandrina, Lake Albert and Goolwa Channel was plotted over the condition monitoring program to assess the benefit of the extended draw-down on the vegetation of the aforementioned habitats (Figure 10a). Native amphibious species richness at +0.6 m AHD declined during the drought but has generally increased since water levels were reinstated until the two latest surveys (Figure 10). The plant community at +0.6 m AHD in Lake Albert is depauperate compared to Lake Alexandrina and Goolwa Channel (Figure 10). Native amphibious species richness at +0.6 m AHD increased between spring 2017 and spring 2018 in all habitats except Lake Alexandrina, with the highest species richness over the condition monitoring program recorded in this habitat in autumn 2018 (Figure 10). Native species richness declined between spring 2018 and spring 2019 in Lake Albert and Goolwa Channel but remained similar in Lake Alexandrina (Figure 10a). Further declines across all habitats were observed between spring 2019 and spring 2020, with a small increase in autumn 2021 (Figure 10a).

a.



Native amphibious species richness at +0.8 m AHD decreased across all habitats during the drought but generally increased after water levels were reinstated except for the most recent survey in Lake Albert and Goolwa Channel (Figure 10b). In contrast to +0.6 m AHD, this elevation is exposed more often and for longer so there is a greater opportunity for species that require exposure to recruit but there may be periods of low soil moisture. There are generally less native amphibious species at +0.8 m AHD compared to +0.6 m AHD across all habitats with Lake Albert having the lowest native amphibious species richness (Figure 10b).

b.




Figure 10: Mean native amphibious species richness per quadrat for Lake Alexandrina, Lake Albert and Goolwa Channel at a. +0.6 m and b. +0.8 m AHD over the condition monitoring program (error bars = ±1 S.E.).

3.2. TLM targets

The following section graphically presents the progress of achievement for each of the targets for each habitat outlined in Tables 1-5 over the duration of the condition monitoring program (spring 2008 to autumn 2021). Target thresholds were defined by the proportion (percentage) of quadrats containing a species or functional group above a certain percentage cover (Tables 1–5). Target thresholds presented in red on the graphs denote targets that are achieved when the percentage of quadrats is lower than the threshold (undesirable taxa) and thresholds presented in blue are met when the percentage of quadrats is higher than the threshold (desirable taxa). In addition, the habitat condition score calculated from the targets achieved from each habitat and the WOISS (calculated from the habitat condition scores) are presented for the duration of the condition monitoring program.

Lake Alexandrina targets

Littoral Zone

Figure 11 shows the percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 75% in the littoral zone from spring 2008 to autumn 2021. Prior to spring 2018, there was a seasonal trend with a higher proportion of quadrats containing a combined cover of these species greater than 75% in autumn compared to spring (Figure 11). In addition, there was a general upward trend of the indicator since water levels were reinstated in spring 2010 to autumn 2015, followed by a sharp decline then another upward trend. The percentage of quadrats with a combined cover greater than 75% did not exceed 40% and the target was consistently achieved (Figure 11).




Figure 11: Percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 75% in the littoral zone of Lake Alexandrina from spring 2008 to autumn 2021.

Figure 12 shows the percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone from spring 2008 to autumn 2021. Between autumn 2009 and autumn 2010, more than 20% of quadrats contained a combined cover of these species greater than 50%; hence, the target was not met. However, when water levels were reinstated in spring 2010 the number of quadrats with a combined cover greater than 50% of these species fell below 20%, the target was achieved and there has been a general downward trend (Figure 12). The lowest proportion of quadrats with these species with a combined cover of greater than 50% was in spring 2018 (Figure 12).




Figure 12: Percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone of Lake Alexandrina from spring 2008 to autumn 2021.

Figure 13 shows the percentage of quadrats containing a cover of native amphibious species $\geq 5\%$ in the littoral zone from spring 2008 to autumn 2021. There has been an upward trend of the indicator since water levels were reinstated; however, this indicator did not exceed 50% of quadrats until autumn 2015 after which it has been consistently achieved, except in spring 2017 (Figure 13).




Figure 13: Percentage of quadrats containing a cover native amphibious species greater than 5% in the littoral zone of Lake Alexandrina from spring 2008 to autumn 2021.

Figure 14 shows the percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* $\geq 5\%$ in the littoral zone from spring 2008 to autumn 2021. The indicator has not exceeded 50% and peaked in spring 2015 (12% of quadrats). Therefore, this target has not been achieved during the condition monitoring program (Figure 14). The number of quadrats containing a cover of these species $\geq 5\%$ has generally increased since water levels were reinstated until autumn 2017, but declined in spring 2017 and has since remained low (Figure 14).




Figure 14: Percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* greater than 5% in the littoral zone of Lake Alexandrina from spring 2008 to autumn 2021.

Aquatic Zone

Figure 15 shows the percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% in the aquatic zone from spring 2008 to autumn 2021. This indicator has generally increased; however, it has not exceeded 40% and the target has been achieved since spring 2008 (Figure 15).




Figure 15: Percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% in the aquatic zone of Lake Alexandrina from spring 2008 to autumn 2021.

Figure 16 shows the percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* $\geq 5\%$ in the aquatic zone from spring 2008 to autumn 2021. This indicator generally increased since water levels were reinstated, peaking in spring 2015 and spring 2018 (18%). There was a seasonal pattern after spring 2013 with higher proportions of quadrats containing these species with a cover of greater than 5% in spring compared to autumn (Figure 16). The target of 20% of quadrats has not been achieved during the condition monitoring program (Figure 16).




Figure 16: Percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* greater than 5% in the aquatic zone of Lake Alexandrina from spring 2008 to autumn 2021.

Figure 17 shows the percentage of quadrats containing a cover of native submergent species $\geq 5\%$ in the aquatic zone from spring 2008 to autumn 2021. During the drought, the aquatic zone was dry; hence, no submergent species were present and it was not until spring 2011 before a significant number of quadrats contained native submergent species. There was an increasing trend for this indicator after spring 2011 until autumn 2015, followed by a general decrease until spring 2017 and peaking in the most recent survey (31% of quadrats) (Figure 17). There was also a seasonal pattern with more quadrats containing native submergents $\geq 5\%$ in autumn compared to spring, except in spring 2019 (Figure 17). The proportion of quadrats containing $\geq 5\%$ cover of these species has never exceeded 35%; therefore, the target has not been achieved to date despite a recent trend in improvement.




Figure 17: Percentage of quadrats containing a cover of native submergent species greater than 5% in the aquatic zone of Lake Alexandrina from spring 2008 to autumn 2021.

Whole of habitat condition

The whole of habitat condition score (the proportion of targets achieved) in Lake Alexandrina is shown in Figure 18. The increase between autumn 2010 and spring 2010 was due to water levels being reinstated and the target for the deep water zone being achieved and the number of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone falling below 20% (Figure 12). No additional targets were achieved until autumn 2015 when the number of quadrats containing cover of native amphibious species $\geq 5\%$ in the littoral zone exceeded 50% and the target was achieved every year since, except in spring 2017 (Figure 13).




Figure 18: Whole of habitat condition score for Lake Alexandrina from spring 2008 to autumn 2021.

Lake Albert

Littoral Zone

Figure 19 shows the percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 75% in the littoral zone from spring 2008 to autumn 2021. There were no quadrats containing a combined cover of these species greater than 75% in the littoral zone until spring 2013, after which, there was a general upward trend peaking at 23% in autumn 2018, after which there was a general decline (Figure 19). The number of quadrats has remained well below 40%; therefore, the target has consistently been achieved since spring 2008.




Figure 19: Percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 75% in the littoral zone of Lake Albert from spring 2008 to autumn 2021.

Figure 20 shows the percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone from spring 2008 to autumn 2021. Since autumn 2010 this indicator has generally decreased, with the number of quadrats falling below 20% in spring 2015 and the target achieved thereafter except in autumn 2018 (Figure 20). There were no quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone in spring 2018, spring 2020 and autumn 2021 (Figure 20).




Figure 20: Percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone of Lake Albert from spring 2008 to autumn 2021.

Figure 21 shows the percentage of quadrats containing a cover of native amphibious species $\geq 5\%$ in the littoral zone from spring 2008 to autumn 2021. The number of quadrats containing these species with a cover of $\geq 5\%$ has been variable since spring 2008 showing no seasonal patterns or general trends over the condition monitoring program (Figure 21). However, there was an increase from spring 2017 with the target being achieved for the first time in spring 2019 after which, there was a decline and the target not achieved for the two most recent surveys (Figure 21).




Figure 21: Percentage of quadrats containing a cover of native amphibious species greater than 5% in the littoral zone of Lake Albert from spring 2008 to autumn 2021.

The combined cover of native emergent species other than *Typha domingensis* and *Phragmites australis* has not exceeded 5% in any quadrats in the littoral zone of Lake Albert since spring 2008; therefore, this target has not been achieved (Table 2).

Aquatic Zone

Figure 22 shows the percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% in the aquatic zone from spring 2008 to autumn 2021. There was a general upward trend of the indicator after water levels were reinstated, except in autumn 2014 and spring 2020 (Figure 22). The proportion of quadrats with a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% has not exceeded 40% (the largest number of quadrats was 18% in spring 2019) in the aquatic zone of Lake Albert and the target has been achieved since spring 2008 (Figure 22).




Figure 22: Percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% in the aquatic zone of Lake Albert from spring 2008 to autumn 2021.

Figure 23 shows the percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* $\geq 5\%$ in the aquatic zone from spring 2008 to autumn 2021. These species are uncommon in Lake Albert and quadrats with a combined cover in the aquatic zone $\geq 5\%$ were recorded on five occasions; autumn 2013, autumn 2015, autumn 2016, autumn 2017 and spring 2019 (peaking at 7% in autumn 2015, autumn 2017 and spring 2019) (Figure 23). Therefore, the target has not been achieved over the survey period.




Figure 23: Percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* greater than 5% in the aquatic zone of Lake Albert from spring 2008 to autumn 2021.

Figure 24 shows the percentage of quadrats containing a cover of native submergent species $\geq 5\%$ in the aquatic zone from spring 2008 to autumn 2021. During the drought the aquatic zone was dry; hence, no submergent species were present. There was only one occasion (spring 2011, 6% of quadrats) when native submergent species were present $\geq 5\%$ cover (Figure 24) in any quadrats; hence, the target has not been achieved.




Figure 24: Percentage of quadrats containing a cover of native submergent species greater than 5% in the aquatic zone of Lake Albert from spring 2008 to autumn 2021.

Whole of habitat condition

The whole of habitat condition score for Lake Albert is shown in Figure 25. The increase between autumn and spring of 2010 was due to water levels being reinstated and the target for the deep water zone being achieved (Figure 25). No additional targets were achieved until spring 2015 when the number of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone fell below 20% (Figure 20). However, the proportion of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone increased to 20% in autumn 2018 (Figure 20) and target was not achieved for this survey; hence, the decrease in condition score in autumn 2018 (Figure 25). This target was achieved in spring 2018 (Figure 20), resulting in an increase in habitat condition score between autumn and spring 2018 (Figure 25). There was a further increase between spring 2018 and spring 2019 (Figure 25), which was due to the percentage of quadrats containing cover of native amphibious species $\geq 5\%$ in the littoral zone exceeding 35%, resulting in the target being achieved for the first time (Figure 21). However, this target was not achieved for the two most recent surveys (Figure 21) resulting in a decline in condition score (Figure 25).




Figure 25: Whole of habitat condition score for Lake Albert from spring 2008 to autumn 2021.

Goolwa Channel

Littoral Zone

Figure 26 shows the percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 75% in the littoral zone from spring 2008 to autumn 2021. The indicator exceeded 50% of quadrats on six occasions (Figure 26). In recent surveys it was below 50%, except in spring 2018, resulting in the target being achieved except in in spring 2018 (Figure 26).




Figure 26: Percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 75% in the littoral zone of Goolwa Channel from spring 2008 to autumn 2021.

Figure 27 shows the percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone from spring 2008 to autumn 2021. The only time this target was not achieved was in autumn 2010, after which there has been a general downward trend (except between autumn 2015 and spring 2017) of the indicator (Figure 27).




Figure 27: Percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone of Goolwa Channel from spring 2008 to autumn 2021.

Figure 28 shows the percentage of quadrats containing a cover of native amphibious species $\geq 5\%$ in the littoral zone from spring 2008 to autumn 2021. The indicator has trended upwards since spring 2008; however, there were also strong seasonal patterns from spring 2009 to spring 2013 with higher abundances of these species in spring (Figure 28). After spring 2013 there were no seasonal patterns and the target has been achieved each subsequent survey (Figure 28).




Figure 28: Percentage of quadrats containing a cover native amphibious species greater than 5% in the littoral zone of Goolwa Channel from spring 2008 to autumn 2021.

Figure 29 shows the percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* $\geq 5\%$ in the littoral zone from spring 2008 to autumn 2021. The indicator has not exceeded 50% of quadrats; therefore, this target has not been achieved during the condition monitoring program (Figure 29). However, there was an increasing trend from spring 2009 to spring 2013 (13% of quadrats) followed by a decrease in autumn 2014 (Figure 29). After autumn 2014 there was another upward trend peaking in spring 2018 at 23% of quadrats followed by a downward trend (Figure 29).




Figure 29: Percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* greater than 5% in the littoral zone of Goolwa Channel from spring 2008 to autumn 2021.

Aquatic Zone

Figure 30 shows the percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% in the aquatic zone from spring 2008 to autumn 2021. The indicator has exceeded 50% of quadrats on four occasions with the target achieved since autumn 2016 (Figure 30).




Figure 30: Percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% in the aquatic zone of Goolwa Channel from spring 2008 to autumn 2021.

Figure 31 shows the percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* $\geq 5\%$ in the aquatic zone from spring 2008 to autumn 2021. The indicator exceeded 50% of quadrats (and the target was achieved) for the first time in spring 2018 and again in spring 2020 with a general upwards trend since spring 2009 (Figure 31). However, there was a decrease between spring 2018 and spring 2019 and spring 2020 and autumn 2021 with the percentage of quadrats falling below 20% and the target not being achieved in these surveys (Figure 31).




Figure 31: Percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* greater than 5% in the aquatic zone of Goolwa Channel from spring 2008 to autumn 2021.

Figure 32 shows the percentage of quadrats containing a cover of native submergent species $\geq 5\%$ in the aquatic zone from spring 2008 to autumn 2021. Before spring 2009 the aquatic zone was dry; hence, no submergent species were present but after 2009 there was an increasing trend in the number of quadrats containing native submergent species with a cover of $\geq 5\%$ peaking in autumn 2016 at 27% of quadrats (Figure 32). Between autumn 2016 and spring 2017 there was a downwards trend followed by an upwards trend with the target being achieved for the first (and only) time in spring 2020 (Figure 32).




Figure 32: Percentage of quadrats containing a cover of native submergent species greater than 5% in the aquatic zone of Goolwa Channel from spring 2008 to autumn 2021.

Deep water zone

Figure 33 shows the percentage of quadrats containing a cover of native submergent species $\geq 5\%$ in the deep water zone from spring 2008 to spring 2019. Before spring 2009, much of the deep water zone was dry; hence, no submergent species were present (Figure 33). Between spring 2009 and spring 2010, there was a large increase in the number of quadrats with $\geq 5\%$ cover of native submergent species (79% in spring 2010) due to the dominance of *Potamogeton pectinatus* after the Clayton Regulator was constructed. After the Clayton Regulator was breached there was a decrease in the number of quadrats with $\geq 5\%$ cover of native submergents but the number above the target of 20% of quadrats until spring 2015 when it decreased and the target was not achieved again until autumn 2018 and again for the two most recent surveys (Figure 33).




Figure 33: Percentage of quadrats containing a cover of native submergent species greater than 5% in the deep water zone of Goolwa Channel from spring 2008 to autumn 2021.

Whole of habitat condition

The whole of habitat condition score for Goolwa Channel is shown in Figure 34. In contrast to lakes Alexandrina (Figure 18) and Albert (Figure 25) there has not been a sustained increase in habitat condition score over the condition monitoring program for Goolwa Channel (Figure 34). The generally increasing trend between spring 2008 and autumn 2015 was due to the deep water target being achieved over this period (Figure 33). The minor fluctuations over this period were due to the *Typha domingensis* and *Phragmites australis* targets in the littoral (Figure 26) and aquatic (Figure 30) zones and the native amphibious species target (Figure 28) in the littoral zone being achieved for some surveys and not others (seasonal patterns in abundance). The decrease in habitat condition score between autumn 2015 and autumn 2016 was because the deep water (Figure 33) and littoral *Typha domingensis* and *Phragmites australis* targets (Figure 26) were not achieved. The increase in condition score between autumn 2017 and autumn 2018 was due to the littoral *Typha domingensis* and *Phragmites australis* target being achieved in spring 2017 and autumn 2018 (Figure 26) and the deep water target being achieved in autumn 2018 (Figure 33). The recent decline in condition score was due to the littoral zone *Typha domingensis* and *Phragmites australis* (Figure 26) and deep water submergent vegetation (Figure 33) targets not being achieved in spring 2018, despite the emergent species other than *Typha domingensis* and *Phragmites australis* target being achieved in the aquatic zone (Figure 31). The further decline between spring 2018 and 2019 was due to the emergent species other than *Typha domingensis* and *Phragmites australis* target not being achieved in the aquatic zone in spring 2019 (Figure 31). The increase between spring 2019 and spring 2020 was due to the deep water (Figure 33) and the aquatic zone native submergent targets being met (Figure 32). The decline between spring 2020 and autumn 2021 was due to the native submergent target in the aquatic zone not being achieved (Figure 32).




Figure 34: Whole of habitat condition score for Goolwa Channel from spring 2008 to autumn 2021.

Permanent wetlands

Littoral zone

Figure 35 shows the percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 75% in the littoral zone from spring 2008 to autumn 2021. Quadrats in the littoral zone containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 75% were uncommon in permanent wetlands and has never exceeded the target of 35% of quadrats (Figure 35). Therefore, this target has been achieved throughout the condition monitoring program (Figure 35).




Figure 35: Percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% in the littoral zone of permanent wetlands from spring 2008 to autumn 2021.

Figure 36 shows the percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone from spring 2008 to autumn 2021. In contrast to lakes Alexandrina (Figure 12) and Albert (Figure 20) and Goolwa Channel (Figure 27), the reinstatement of water levels did not result in a decrease in the indicator. Between autumn 2010 and autumn 2016 there was a decreasing trend but a large increase between autumn 2016 and autumn 2018 (Figure 36). The highest proportion of quadrats (54%) containing more than 50% cover of these species was in autumn 2018, after which there was a general decline but not sufficient to achieve the target (Figure 36). The target was only achieved in spring 2008 and spring 2009 (Figure 36).




Figure 36: Percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% in the littoral zone of permanent wetlands from spring 2008 to autumn 2021.

Figure 37 shows the percentage of quadrats containing a cover of native amphibious species $\geq 5\%$ in the littoral zone from spring 2008 to autumn 2021. The was highly variable but has generally tended upwards for the duration of the monitoring (Figure 37). The target of 50% of quadrats having a cover of native amphibious species $\geq 5\%$ was achieved for the first time in spring 2020 and again in autumn 2021 (Figure 37).




Figure 37: Percentage of quadrats containing a cover native amphibious species greater than 5% in the littoral zone of permanent wetlands from spring 2008 to autumn 2021.

Figure 38 shows the percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* $\geq 5\%$ in the littoral zone from spring 2008 to autumn 2021. The indicator has not exceeded 50% of quadrats; therefore, this target has not been achieved during the condition monitoring program (Figure 38). There was a general increasing trend of the indicator until autumn 2016, after which it decreased sharply but there has been an increasing trend since autumn 2018 (Figure 38).




Figure 38: Percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* greater than 5% in the littoral zone of permanent wetlands from spring 2008 to autumn 2021.

Aquatic zone

Figure 39 shows the percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% in the aquatic zone from spring 2008 to autumn 2021. The indicator showed an increasing trend between autumn 2009 and autumn 2015 after which it levelled and decreased to zero in spring 2019 (Figure 39). The number of quadrats has not exceeded 40% and the target was consistently achieved over the monitoring program (Figure 39).




Figure 39: Percentage of quadrats containing a combined cover of *Typha domingensis* and *Phragmites australis* greater than 50% in the aquatic zone of permanent wetlands from spring 2008 to autumn 2021.

Figure 40 shows the percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* $\geq 5\%$ in the aquatic zone from spring 2008 to autumn 2021. There was an increase in the indicator after water levels were reinstated but that decreased to zero by autumn 2013 (Figure 40). The number of quadrats remained at 5% or lower until spring 2017 and peaked in autumn 2018 with 17%, decreased to 5% in spring 2018 and has since shown an upward trend (Figure 40). The target has not been achieved during the condition monitoring program (Figure 40).




Figure 40: Percentage of quadrats containing a cover of native emergent species other than *Typha domingensis* and *Phragmites australis* greater than 5% in the aquatic zone of permanent wetlands from spring 2008 to autumn 2021.

Figure 41 shows the percentage of quadrats containing a cover of native submergent species between 5 and 50% in the aquatic zone from spring 2008 to autumn 2021. Before spring 2010 the cover of native submergent species was low due to this zone largely being dry (although there were isolated puddles in spring 2008 and 2009 that supported submergent species) (Figure 41). After water levels were reinstated there has been a general increasing trend, with the target being achieved for the first time in spring 2020 and again in autumn 2021 (Figure 41).




Figure 41: Percentage of quadrats containing a cover of native submergent species between 5 and 50% in the aquatic zone of permanent wetlands from spring 2008 to autumn 2021.

Whole of habitat condition

The whole of habitat condition score for permanent wetlands is shown in Figure 42. There has been little change in the habitat condition score for permanent wetlands between spring 2008 and spring 2019 (Figure 42). The fall in condition score between spring 2008 and autumn 2009, rise between autumn 2009 and spring 2009 and fall between spring 2009 and autumn 2010 was due to the littoral zone *Paspalum distichum* and *Cenchrus clandestinus* being achieved in spring 2008 and spring 2009 (Figure 36). From autumn 2010 there was no change in condition score until spring 2020 with the only targets that were achieved consistently over this period being the *Typha domingensis* and *Phragmites australis* targets in the littoral (Figure 35) and aquatic (Figure 39) zones. In spring 2020 and autumn 2021 the littoral zone amphibious (Figure 46) and aquatic zone submergent (Figure 48) targets were achieved resulting in an increase in habitat condition score.




Figure 42: Whole of habitat condition score for permanent wetlands from spring 2008 to autumn 2021.

Seasonal wetlands

Wetland edge

Figure 43 shows the percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% around the edges of seasonal wetlands from spring 2008 to autumn 2021. There has been an increasing trend in the indicator around the edges of seasonal wetlands over the duration of the condition monitoring program (Figure 43). In addition, there was a seasonal pattern with higher abundances in autumn when water levels are low (Figure 43). The target of a maximum of 20% of quadrats was exceeded (target not achieved) in autumn 2012, autumn 2013 and from autumn 2014 onwards (Figure 43).




Figure 43: Percentage of quadrats containing a combined cover of *Paspalum distichum* and *Cenchrus clandestinus* greater than 50% around the edge of seasonal wetlands from spring 2008 to autumn 2021.

Figure 44 shows the percentage of quadrats containing a cover of native amphibious species $\geq 5\%$ in the around the edges of seasonal wetlands from spring 2008 to autumn 2021. Native amphibious species were common around the edges of seasonal wetlands and the number of quadrats with a cover of $\geq 5\%$ was higher than the 50% target throughout the survey period despite the downward trend from autumn 2016 (Figure 44). Therefore, this target was consistently achieved over the condition monitoring program (Figure 44).




Figure 44: Percentage of quadrats containing a cover native amphibious species greater than 5% around the edge of seasonal wetlands from spring 2008 to autumn 2021.

Figure 45 shows the percentage of quadrats containing a cover of native emergent species $\geq 5\%$ around the edges of seasonal wetlands from spring 2008 to autumn 2021. The percentage of quadrats with a cover of native emergent species $\geq 5\%$ has not exceeded 50% of quadrats; therefore, this target has not been achieved during the condition monitoring program (Figure 45). However, there was a general increasing trend from over the condition monitoring program (Figure 45).




Figure 45: Percentage of quadrats containing a cover of native emergent species greater than 5% around the edge of seasonal wetlands from spring 2008 to autumn 2021.

Wetland bed

Figure 46 shows the percentage of quadrats containing a cover of native amphibious species $\geq 5\%$ on the beds of seasonal wetlands from spring 2008 to autumn 2021. Native amphibious species were less common on the beds of seasonal wetlands compared to the edges (Figure 44). The number of quadrats with a combined cover of these species $\geq 5\%$ peaked in spring 2010 (59%), after which it fell to 26% and fluctuated between 15% and 32% until spring 2016 when it rose to 50% but fell to 18% in autumn 2017 (Figure 46). The target was achieved 13 times over the condition monitoring program but not in the four most recent surveys, with spring 2020 having the lowest percentage (13%) of quadrats containing native amphibious species with a cover of $\geq 5\%$ (Figure 46).




Figure 46: Percentage of quadrats containing a cover of native amphibious species greater than 5% on the bed of seasonal wetlands from spring 2008 to autumn 2021.

Figure 47 shows the percentage of quadrats containing a cover of native emergent species $\geq 5\%$ on the beds of seasonal wetlands from spring 2008 to autumn 2021. Between spring 2008 and spring 2009 there was a decrease of the indicator (Figure 47). However, the number of quadrats increased between spring 2009 and autumn 2011, after which there was a seasonal pattern with higher abundances in autumn compared to spring (Figure 47). The target of 20% of quadrats was first achieved in autumn 2011 and was achieved each subsequent survey, except in spring 2013 and spring 2018 (Figure 47).




Figure 47: Percentage of quadrats containing a cover of native emergent species greater than 5% on the beds of seasonal wetlands from spring 2008 to autumn 2021.

Figure 48 shows the percentage of quadrats containing a cover of native submergent species $\geq 25\%$ on the beds of seasonal wetlands in spring from 2008 to 2020. Before spring 2011 the cover of native emergent species was low due to seasonal wetlands beds largely being dry (although there were isolated puddles in spring 2008 and 2009 that supported submergent species) (Figure 41). After water levels were reinstated in spring 2010, there has been a generally increasing trend in the indicator, with the highest percentage of quadrats with submergent species greater than 25% cover (45%) in spring 2019 (Figure 41). However, the target of 50% of quadrats has not been achieved over the duration of the condition monitoring program (Figure 41).




Figure 48: Percentage of quadrats containing a cover of native submergent species greater than 25% in spring on the beds of seasonal wetlands from 2008 to 2020.

Whole of habitat condition

The whole of habitat condition score for seasonal wetlands is shown in Figure 49. There has not been a sustained increase in habitat condition score over the condition monitoring program for temporary wetlands (Figure 49). There is a seasonal pattern in wetland condition score with scores usually higher in autumn compared to spring, which is due to the higher abundance of native amphibious (Figure 44; Figure 46) and emergent (Figure 45; Figure 47) species and there being no submergent species target in autumn (which has never been achieved in spring) (Figure 48). There was no change in score between autumn 2017 and autumn 2018 after which there was a decrease resulting in the lowest habitat condition score over the condition monitoring program (Figure 49). The decrease between autumn and spring 2018 was due to the native amphibious (Figure 46) and emergent species (Figure 47) targets for the wetland beds not being achieved in spring 2018. However, there was an increase between spring 2018 and spring 2019 due to the percentage of quadrats containing a cover of native emergent species $\geq 5\%$ on wetlands beds of being greater than 20% and the target being achieved (Figure 47). There was no change between spring 2019 and spring 2020 and the increase in autumn 2021 again due to the absence of a submergent target (Figure 47).




Figure 49: Whole of habitat condition score for seasonal wetlands from spring 2008 to autumn 2021.

Whole of lakes condition

The whole of Icon Site score (WOISS) for aquatic and littoral vegetation represents the proportion of targets achieved throughout the five different habitats and using the Matter 8 condition rating. Based on this score, objective V3 has been achieved because a 'good' rating has been reached in the two most recent surveys (Table 9).

During the period of low water levels (surveys prior to spring 2010) the WOISS was low and fluctuated between 0.32 and 0.34 (poor condition rating (Table 9)). There was an increase between spring 2009 and spring 2011, after which it fluctuated until spring 2018 after which there was an increase and a 'good' rating was achieved in the two latest surveys (Figure 50). The increase in condition score between spring 2009 and spring 2010 (increase from poor to fair condition (Table 9) was due to increases in habitat condition scores in Lake Alexandrina (Figure 18), Lake Albert (Figure 25), Goolwa Channel (Figure 34) and seasonal wetlands (Figure 49). The decrease between spring 2011 and spring 2013 was due to decreases in habitat condition scores in Goolwa Channel (Figure 34) and seasonal wetlands (Figure 49) and the increase between spring 2013 and autumn 2015 was due to increases in habitat condition scores in Lake Alexandrina (Figure 18), Goolwa Channel (Figure 34) and seasonal wetlands (Figure 49). The decrease in condition score between autumn 2015 and spring 2015 was due to decreases in condition score in Goolwa Channel (Figure 34) and seasonal wetlands (Figure 49). There was no change in condition scores for all habitats between autumn 2016 and spring 2017; hence, the whole of lakes condition score remained the same during that period (Figure 50). The increase between spring 2017 and autumn 2018 was predominantly due to the increase in condition score in Goolwa Channel over the same period (Figure 34). The decrease between autumn and spring 2018 was due to decreases in habitat condition scores for Goolwa Channel (Figure 34) and seasonal wetlands (Figure 49), despite the increase in Lake Albert (Figure 25). The increase in condition score between spring 2018 (0.43) and spring 2019 (0.48) (Table 9) was due the increases in habitat condition scores in Lake Albert (Figure 25) and seasonal wetlands (Figure 49), despite the decrease in Goolwa Channel (Figure 34). The further increase in condition score (Figure 50) and improvement in vegetation condition from fair to good between spring 2019 and 2020 (Table 9) was due to increases in habitat condition scores in Goolwa Channel (Figure 34), permanent (Figure 42) and temporary wetlands (Figure 49) over this period.




Figure 50: Whole of lakes condition score from spring 2008 to autumn 2021.

Table 9: Whole of lakes condition score and Matter 8 condition rating (Department for Environment and Water 2019) from spring 2008 to autumn 2021.

Survey Date	Whole of lakes condition score	Condition rating
Spring 2008	0.32	Poor
Autumn 2009	0.34	Poor
Spring 2009	0.32	Poor
Autumn 2010	0.36	Poor
Spring 2010	0.50	Fair
Autumn 2011	0.56	Fair
Spring 2011	0.56	Fair
Autumn 2012	0.49	Fair
Spring 2012	0.49	Fair
Autumn 2013	0.50	Fair
Spring 2013	0.48	Fair
Autumn 2014	0.49	Fair
Autumn 2015	0.55	Fair
Spring 2015	0.45	Fair
Autumn 2016	0.46	Fair
Autumn 2017	0.46	Fair
Spring 2017	0.46	Fair
Autumn 2018	0.53	Fair
Spring 2018	0.43	Fair
Spring 2019	0.48	Fair
Spring 2020	0.63	Good
Autumn 2021	0.60	Good

4. DISCUSSION AND MANAGEMENT IMPLICATIONS

4.1. Impacts of water level and salinity

During the most recent survey period (spring 2020 to autumn 2021), salinity (Figure 3) in Lake Alexandrina and the Goolwa Channel remained similar to values recorded since spring 2010. Electrical conductivity also remained stable in Lake Albert and was higher than Lake Alexandrina, although the higher salinity in Lake Albert is probably not biologically significant for the plant species present. Water levels in summer/autumn 2020-21 were not drawn-down as far as in the previous three years with the minimum water level around +0.6 m AHD.

During the drought-induced draw-down period (2007 to 2010), plant assemblages had shifted towards terrestrial taxa. However, following restoration of water levels in the Lower Lakes in late August 2010 (and the subsequent reconnection of most wetlands) there has been a general increasing trend in the abundance and diversity of aquatic dependent taxa (e.g. submergent, amphibious and emergent), suggesting the vegetation of the system is still recovering.

During 2012/13, water level management in the Lower Lakes involved two draw-down and refilling cycles (between +0.4 and +0.8 m AHD) with the aim to reduce salinity in Lake Albert (Figure 2). There have been no deliberate lake level cycles since then; however, the typical seasonal cycle of high water levels in spring and early summer and low water levels in autumn has occurred each year (Figure 2). Stable water levels have been identified as detrimental to aquatic plant communities, with a greater diversity of aquatic plants generally in systems with fluctuating water levels (e.g. Nielsen and Chick 1997). Increases in water levels between autumn 2016 and autumn 2017 periodically inundated areas at higher elevations (above +0.9 m AHD) in spring 2016, which may have resulted in the increase in abundance of *Cenchrus clandestinus* and *Paspalum distichum* in permanent and temporary wetland habitats. The lower water levels in autumn 2017, 2018 and 2019 exposed the fringes of lakeshores and wetlands, which provided opportunities for species requiring exposure to germinate (e.g. *Persicaria lapathifolia*, *Berula erecta*, *Calystegia sepium*, *Ludwigia peploides*, *Juncus* spp., *Cyperus gymnocaulos*) (Nicol 2004). There may be limited opportunity for recruitment of species that require exposure to germinate due to fringing areas being densely vegetated with emergent species such as *Typha domingensis* or *Phragmites australis*. However, native amphibious plant species richness at +0.6 m AHD at the site scale was higher in the autumn 2017 and 2018 surveys compared to spring 2017 in all habitats indicating draw-down provided opportunities for species to recruit. Furthermore, native amphibious species richness in spring 2018 and 2019 (despite decreasing during this period

in Lake Albert and Goolwa Channel) was higher than spring 2017 across all habitats (Figure 10) indicating an increase that persisted for several years, most likely due to seasonal lake level cycling. However, there was a decrease in native amphibious species richness at +0.6 m AHD in the two most recent surveys across all habitats that may be due to the higher lake levels throughout the 2020-21 water year. Despite the decrease in native amphibious species richness at +0.6 m AHD, it was maintained at +0.8 m AHD indicating that these species have not been lost from the littoral zone and should recruit in the future at lower elevations when draw-down below +0.6 m AHD. Often shorelines that are not densely vegetated are subjected to wave action, which can prevent seedlings from establishing (e.g. Foote and Kadlec 1988). Nevertheless, seasonal water level fluctuations between $>+0.8$ and $+0.5$ m AHD are recommended because areas of submergent vegetation are maintained and the establishment of amphibious taxa in areas protected from wave action is facilitated. The main downside to these water level fluctuations is the increase in abundance of *Paspalum distichum* and (to a lesser extent) *Cenchrus clandestinus* in recent years.

4.2. Change in plant community, spring 2008 to autumn 2021

The change in floristic composition observed over the duration of the condition monitoring program (spring 2008 to autumn 2021) has provided information regarding the recovery of the aquatic and littoral plant community, after the Millennium Drought, which resulted in complete loss of the submergent plant community and a decrease in the abundance of amphibious and emergent species. Pooling data from each habitat, although at the cost of losing information regarding the response of individual wetlands, has enabled the change in floristic composition to be analysed at a broader spatial scale. There were similarities in the patterns of change among habitats, such as the expected large changes observed when water levels were reinstated in spring 2010, the decrease in change between surveys through time and the seasonal patterns evident in some habitats.

The smaller change in floristic composition in recent surveys for all habitats except seasonal wetlands may indicate that the current plant community may persist into the future with only minor changes, providing recent salinity and water level regimes are maintained. However, multiple, minor, non-seasonal changes through time can result in large (albeit gradual) changes in the plant community through time. There is evidence this may have occurred in recent years in all habitats (Figure 6). The points on this ordination from the latest surveys, whilst showing less change in floristic composition among surveys compared to earlier surveys (pre autumn 2011), exhibited a temporal directional change. Furthermore, many of the TLM targets have shown decreasing or increasing trends in the abundances of species

or functional groups in recent years that suggests there may be gradual changes in floristic composition that continue to occur through time.

The patterns observed in the temporary wetlands were expected due to the patterns in seasonal inundation and spring surveys being undertaken when submergent species were present. Whilst the spring plant community in seasonal wetlands was variable, the autumn plant community was similar among surveys. The variability among the spring surveys was due to differences in the abundances of submergent species over time. In comparison, the plant community in autumn was dominated by *Phragmites australis*, *Sarcocornia quinqueflora* and *Paspalum distichum*.

It is unknown whether the plant community present in recent years is comparable to the community prior to 2007 because direct quantitative comparisons between the condition monitoring data and the small amount of data collected prior to 2007 cannot be made. However, for sites where data does exist (Teringie, Narrung, Clayton Bay, Dunn's Lagoon, Milang, Loveday Bay, Point Sturt and Hunters Creek), the diversity and abundance of submergent species were higher before 2007 compared to recent surveys (Holt *et al.* 2005; Nicol *et al.* 2006). For example, Holt *et al.* (2005) reported extensive beds of *Vallisneria australis* and *Myriophyllum salsgineum* throughout Dunns Lagoon almost completely covering the permanently inundated areas in spring 2004. In addition, Nicol *et al.* (2006) reported a bed of dense *Ruppia polycarpa* covering the entire inundated area of Point Sturt wetland. In the most recent surveys, *Myriophyllum salsgineum* and *Vallisneria australis* were present in Dunns Lagoon and abundant in places, but overall vegetation cover across the lagoon was patchy. In addition, *Ruppia polycarpa* has not been recorded in Point Sturt Wetland during the condition monitoring program but in the most recent three spring surveys, the low elevations were dominated by *Ruppia tuberosa*.

4.3. The Living Murray targets and condition scores

The original vegetation target (now an objective) (V3) for the Lower Lakes: *maintain or improve aquatic and littoral vegetation in the Lower Lakes*, whilst an appropriate management aim and ecological objective for the system, could not properly be assessed because there is no quantitative baseline. Furthermore, baseline data would need to be collected over a minimum of 5–10 years to determine the natural (acceptable) variability of the system. Davis and Brock (2008) identified this as a problem when determining limits of acceptable change for wetlands of international importance under the Ramsar Convention. These authors proposed that conceptual models be developed to determine limits of acceptable change and to design a monitoring program to assess and refine the proposed

limits of acceptable change (Davis and Brock 2008). Nicol (2016) proposed limits of acceptable change (and management triggers) for aquatic and littoral vegetation in the Coorong and Lakes Alexandrina and Albert Ramsar Wetland using conceptual models (*sensu* Davis and Brock 2008), and TLM aquatic and littoral vegetation targets were based on proposed limits of acceptable change management triggers. In addition, the Whole Of Icon Site Scores (WOISS) have been used for the South Australian Basin Plan environmental outcome reporting (Matter 8) and the scale used to report on the achievement of Objective V3.

The refined targets (Department for Environment, Water and Natural Resources 2017) were based largely on expert opinion; however, data from the 2005 (Holt *et al.* 2005), 2006 (Nicol *et al.* 2006), habitat mapping (Seaman 2003), biological surveys of conservation reserves adjacent to the Murray Mouth (Brandle *et al.* 2002), a survey of the aquatic vegetation of Hindmarsh Island (Renfrey *et al.* 1989) and condition monitoring data were also used to develop the targets. The condition monitoring program was designed prior to the development of the refined targets but the data collected could be used to assess and refine the targets in addition to providing data to monitor change in floristic composition through time. Continued repeated surveys will allow the targets to be refined in the future.

The habitats with the highest proportion of targets achieved in the most recent surveys and; therefore, having the highest condition score were Lake Alexandrina and Goolwa Channel. The condition scores for lakes Alexandrina and Albert have generally been stable or increasing since water levels were reinstated. In contrast, the condition score for Goolwa Channel was highly variable and decreased between autumn 2018 and spring 2019 but increased in the two most recent surveys.

In Lake Alexandrina, there has generally been progress towards achieving the targets that require an increase in the abundance of desirable species (native amphibious and submergent species) that have not yet been achieved. However, emergent species other than *Typha domingensis* and *Phragmites australis* in the littoral and aquatic zones have generally declined in recent surveys. There was also a downward trend in the number of quadrats that were dominated by *Paspalum distichum* and *Cenchrus clandestinus* after spring 2010. However, there was an increase in the number of quadrats dominated by *Typha domingensis* and *Phragmites australis* but the proportion of those quadrats was not close to the target. These trends suggest that the condition score in Lake Alexandrina will continue to improve through time as more targets are achieved providing the abundances of *Typha domingensis* and *Phragmites australis* remain at current levels and there is not an increase

in *Paspalum distichum* and *Cenchrus clandestinus* abundance as observed in permanent and temporary wetlands.

The habitat condition score for Lake Albert has also been generally stable or increasing since water levels were reinstated. There has been a general downward trend in quadrats dominated by *Paspalum distichum* and *Cenchrus clandestinus* after spring 2010 except between spring 2017 and autumn 2018 and the target was not achieved for this survey; hence, the decline in habitat condition score over this period. Between autumn 2018 and autumn 2021 there was a general decrease in quadrats dominated by *Paspalum distichum* and *Cenchrus clandestinus* and the target was achieved resulting in an increase in habitat score. Similar to Lake Alexandrina, there has been an increase in the number of quadrats dominated by *Typha domingensis* and *Phragmites australis* but not to a level close to the target. The progress towards achievement of targets that require an increase in the abundance of desirable species observed in Lake Alexandrina has generally not occurred in Lake Albert, except for native amphibious species in the littoral zone. Quadrats that contained these species with $\geq 5\%$ cover exceeded 35% in spring 2019, resulting in the target being achieved for the first time and (Figure 21), which resulted in an increase in habitat score in spring 2019 but there was a decrease in the two most recent surveys resulting in a decrease in the habitat score (Figure 25).

In contrast to condition scores in lakes Alexandrina and Albert, the condition score in Goolwa Channel is highly variable. This is primarily due to an increase in the number of quadrats dominated by *Typha domingensis* and *Phragmites australis* in the littoral zone and a decrease in the abundance of submergent species in the deep water zone. However, since spring 2010 the number of quadrats dominated by *Typha domingensis* and *Phragmites australis* in the littoral and aquatic zones has been relatively stable and fluctuated around the target level (Figure 26 and Figure 30). The target was met for the three most recent survey in both zones (Figure 26 and Figure 30). The deep water zone target was achieved in autumn 2018 for the first time since autumn 2015 and again for the two most recent surveys. In Spring 2020 all targets except the emergent other than *Typha domingensis* and *Phragmites australis* in the littoral zone (Figure 29) were achieved resulting in the highest habitat condition score for the condition monitoring program (Figure 34). The submergent (Figure 32) and emergent species other than *Typha domingensis* and *Phragmites australis* in the aquatic zone (Figure 31) were not met in the most recent survey resulting in a decrease in habitat condition score but still the second highest score over monitoring program. Similar to Lake Alexandrina, there has generally been progress towards achieving the targets that require an increase in the

abundance of desirable species and if these trends continue, the condition score for Goolwa Channel will increase again and be maintained at a high level.

The condition habitat score for permanent wetlands has remained constant between spring 2009 and spring 2019 and was lower than that for lakes Alexandrina and Albert over the same period. There was a downward trend in the number of quadrats dominated by *Paspalum distichum* and *Cenchrus clandestinus* between spring 2010 and autumn 2016 but since then there has been a general increase and the target has not been achieved since spring 2009. There has been progress towards achieving the targets that require an increase in the abundance of desirable species since water levels were reinstated (except emergent species other than *Typha domingensis* and *Phragmites australis* in the littoral and aquatic zones) with the targets for amphibious taxa in the littoral zone (Figure 37) and submergents in the aquatic zone (Figure 41) being achieved in the two most recent surveys. This resulted in an increase in the habitat condition score similar to the values in Lakes Alexandrina (Figure 18) and Albert (Figure 25).

The condition score for seasonal wetlands over the duration of the condition monitoring program was variable until autumn 2015, after which it has trended downwards. The peaks in autumn were due to the absence of a submergent vegetation target for this season. The downward trend since autumn 2015 and autumn 2017 was due to an increase in the number of quadrats dominated by *Paspalum distichum* and *Cenchrus clandestinus* (which has shown an increasing trend since spring 2008). Furthermore, the percentage of quadrats containing native amphibious and emergent species with a cover of greater than 5% on the wetland bed have fluctuated around the target with both targets not being met in spring 2018 but the emergent target achieved in spring 2019 onwards. This resulted in an increase in the habitat score for this survey, which was maintained in spring 2020. The number of quadrats containing submergent species with a cover $\geq 25\%$ on the wetland bed in spring (the most recent survey had the highest percentage recorded (45%)) and those with a cover of native emergent species around the edge of $\geq 5\%$ have generally increased since spring 2010. These trends, if they continue, suggest that the condition score for seasonal wetlands will increase in the future.

The WOISS has fluctuated between 0.56 and 0.43 between autumn 2011 and spring 2019, which is classed as being in fair condition but not achieving objective V3. Many of the individual indicators have fluctuated around their target values, which has resulted in small to moderate variations in the habitat condition scores and hence the whole of lakes condition score. However, in the two most recent surveys, several of the increasing trends for targets that require an increase in the abundance of desirable taxa exceeded the threshold for

targets being achieved, resulting in increases in habitat scores and the whole of lakes condition score. In spring 2020 and autumn 2021 the whole of lakes condition score was higher than 0.6, which resulted in the condition rating improving from fair to good and meeting the ecological objective for the first time in the condition monitoring program. Furthermore, most of the targets not achieved in the two most recent surveys are trending towards being met, suggesting that under the current hydrological and salinity regime the plant community is improving through time. Therefore, it is important that the current salinity and water level regimes are maintained to provide conditions for the continual improvement of vegetation condition.

4.4. Further studies

Suggested further studies (in priority order) to improve the understanding of the vegetation dynamics of the Lower Lakes and the impacts of changes in water levels and salinity include:

1. continuation of the condition monitoring program (with both spring and autumn surveys and *Melaleuca halmaturorum* demographics) to continue to improve understanding of the medium to long-term vegetation dynamics of the system, monitor the recovery trajectory post hydrological restoration (e.g. do current trends persist or is there an equilibrium state?) and to refine indicators;
2. mapping of large-scale plant communities in the Goolwa Channel (*sensu* Gehrig *et al.* 2011a), expanding to key wetlands and lakeshore areas to complement the condition monitoring program and gain a better understanding of vegetation dynamics at the landscape scale;
3. integration of existing data sets for plant and other biotic groups such as fish, birds and invertebrates to better understand relationships among components of the wider aquatic ecosystem to inform (a) development of broader ecological indicators and (b) future research directions;
4. investigation of different control methods for *Paspalum distichum* and *Cenchrus clandestinus* such as controlled summer grazing, herbicides and mowing and monitor to determine effectiveness and native species recovery;
5. investigation of the salinity tolerances of potential local ecotypes of key species (e.g. *Typha domingensis*, *Phragmites australis*, *Schoenoplectus tabernaemontani*, *Vallisneria australis*, *Myriophyllum salsugineum*);
6. investigation of the effects of elevated but sub-lethal salinities on key species;
7. determine propagule longevity under different conditions (e.g. salinity, pH, soil moisture);

8. investigation of the the current submergent plant propagule bank in key wetlands and Goolwa Channel;
9. trial emergent vegetation control at *Melaleuca halmaturorum* stands and monitor to determine whether competition is restricting recruitment.

REFERENCES

- Abernethy, B. and Rutherford, I.D. (1998). Where along a river's length will vegetation most effectively stabilise stream banks? *Geomorphology* **23**: 55-75.
- Bailey, P., Boon, P. and Morris, K. (2002). Salt sensitivity database. Land and Water Australia.
- Bassett, I.E., Paynter, Q. and Beggs, J.R. (2012). Invertebrate community composition differs between invasive herb alligator weed and native sedges. *Acta Oecologica* **41**: 65-73.
- Bell, N., Riis, T., Suren, A.M. and Baattrup-Pedersen, A. (2013). Distribution of invertebrates within beds of two morphologically contrasting stream macrophyte species. *Fundamental and Applied Limnology* **183**: 309-321.
- Bice, C., Wilson, P. and Ye, Q. (2008). Threatened fish populations in the Lower Lakes of the River Murray in spring 2007 and summer 2008. South Australian Research and Development Institute (Aquatic Sciences), F2008/000801-1, Adelaide.
- Blanch, S.J., Ganf, G.G. and Walker, K.F. (1999a). Growth and resource allocation in response to flooding in the emergent sedge *Bolboschoenus medianus*. *Aquatic Botany* **63**: 145-160.
- Blanch, S.J., Ganf, G.G. and Walker, K.F. (1999b). Tolerance of riverine plants to flooding and exposure by water regime. *Regulated Rivers: Research and Management* **15**: 43-62.
- Blanch, S.J., Walker, K.F. and Ganf, G.G. (2000). Water regimes and littoral plants in four weir pools of the River Murray, Australia. *Regulated Rivers Research and Management* **16**: 445-456.
- Brandle, R., Hammer, M., Wedderburn, S., Seaman, R., Noye, R. and Queale, L. (2002). A biological survey of the Murray Mouth Reserves, South Australia March 2002. Department for Environment and Heritage, Adelaide.
- Braun-Blanquet, J. (1932). 'Plant Sociology.' (McGraw-Hill: New York).
- Bray, J.R. and Curtis, J.T. (1957). An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs* **27**: 325-349.
- Brock, M.A. and Casanova, M.T. (1997). Plant life at the edge of wetlands: ecological responses to wetting and drying patterns. In 'Frontiers in Ecology: Building the Links'. (Eds Klomp, N. and Lunt, I.) pp. 181-192. (Elsevier Science: Oxford).

Camargo, A., Pezzato, M., Henry-Silva, G. and Assumpcao, A. (2006). Primary production of *Utricularia foliosa* L., *Egeria densa* Planchon and *Cabomba furcata* Schult & Schult.f from rivers of the coastal plain of the State of Sao Paulo, Brazil. *Hydrobiologia* **570**: 35-39.

Casanova, M.T. (2011). Using water plant functional groups to investigate environmental water requirements. *Freshwater Biology* **56**: 2637-2652.

Centre for Australian National Biodiversity Research and Council of Heads of Australasian Herbaria (2021). Australian Plant Census, IBIS database, <http://www.chah.gov.au/apc/index.html>.

Clarke, K.R. and Gorley, R.N. (2015). PRIMER Version 7.0.12: User Manual/Tutorial. (PRIMER-E: Plymouth).

Cunningham, G.M., Mulham, W.E., Milthorpe, P.L. and Leigh, J.H. (1992). 'Plants of Western New South Wales.' (CSIRO Publishing: Collingwood).

Dashorst, G.R.M. and Jessop, J.P. (1998). 'Plants of the Adelaide Plains and Hills.' (The Botanic Gardens of Adelaide and State Herbarium: Adelaide).

Davis, J. and Brock, M.A. (2008). Detecting unacceptable change in the ecological character of Ramsar wetlands. *Ecological Management and Restoration* **9**: 26-32.

Department for Environment and Water (2019). South Australian River Murray Basin Plan Environmental Outcome Evaluation: Coorong, Lower Lakes and Murray Mouth (CLLMM) Priority Environmental Asset, DEW Technical report 2020/18, Government of South Australia, Department for Environment and Water, Adelaide.

Department for Environment and Water (2021a). Unpublished surface water electrical conductivity for Lake Alexandrina, Lake Albert and Goolwa Channel from August 2008 to May 2021.

Department for Environment and Water (2021b). Unpublished surface water levels for Lake Alexandrina, Lake Albert and Goolwa Channel from August 2008 to May 2021.

Department for Environment, Water and Natural Resources (2017). Condition Monitoring Plan (Revised) 2017. The Living Murray – Lower Lakes, Coorong and Murray Mouth Icon Site. DEWNR, Adelaide.

dos Santos, A.M. and Esteves, F.D. (2002). Primary production and mortality of *Eleocharis interstincta* in response to water level fluctuations. *Aquatic Botany* **74**: 189-199.

Fitzpatrick, R., Grealish, G., Shand, P., Marvanek, S., Thomas, B., Creeper, N., Merry, R. and Raven, M. (2009a). Preliminary assessment of acid sulfate soil materials in Currency Creek, Finniss River, Tookayerta Creek and Black Swamp region, South Australia. CSIRO Land and Water, Adelaide.

Fitzpatrick, R.W., Grealish, G., Chappell, A., Marvanek, S. and Shand, P. (2010). Spatial variability of subaqueous and terrestrial acid sulfate soils and their properties, for the Lower Lakes South Australia. CSIRO Sustainable Agriculture National Research Flagship R-00689-01-015, Adelaide.

Fitzpatrick, R.W., Grealish, G., Shand, P., Simpson, S.L., Merry, R.H. and Raven, M.D. (2009b). Acid sulfate soil assessment in Finniss River, Currency Creek, Black Swamp and Goolwa Channel, South Australia. CSIRO Land and Water, 26/09, Adelaide.

Fitzpatrick, R.W., Grealish, G.H., Shand, P. and N.L., C. (2011). Monitoring and assessment of reflooded Acid Sulfate Soil materials in Currency Creek and Finniss River region, South Australia. CSIRO: Sustainable Agriculture Research Flagship, R-325-8-6, Adelaide.

Foote, A.L. and Kadlec, J.A. (1988). Effects of wave energy on plant establishment in shallow lacustrine wetlands. *Journal of Freshwater Ecology* 4: 523-532.

Frahn, K.A., Gehrig, S.L., Nicol, J.M. and Marsland, K.B. (2013). Lower Lakes vegetation condition monitoring – 2012/2013. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2009/000370-5, Adelaide.

Frahn, K., Gehrig, S., Nicol, J. and Marsland, K. (2014). Lower Lakes Vegetation Condition Monitoring – 2013/2014. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2009/000370-6, Adelaide.

Gehrig, S.L. and Nicol, J.M. (2010a). Aquatic and littoral vegetation monitoring of Goolwa Channel 2009-10. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2010/000383-1, Adelaide.

Gehrig, S.L. and Nicol, J.M. (2010b). Aquatic and littoral vegetation of the Murray River downstream of Lock 1, the Lower Lakes, Murray Estuary and Coorong. A literature review. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2010/000297-1, Adelaide.

Gehrig, S.L., Nicol, J.M. and Bucater, L. (2011a). Aquatic and littoral vegetation monitoring of Goolwa Channel 2009-11. South Australian Research and Development Institute (Aquatic Sciences), F2010/000383-2, Adelaide.

Gehrig, S.L., Nicol, J.M., Frahn, K.A. and Marsland, K.B. (2012). Lower Lakes vegetation condition monitoring – 2011/2012. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2010/000370-4, Adelaide.

Gehrig, S.L., Nicol, J.M. and Marsland, K.B. (2010). Lower Lakes aquatic and littoral vegetation condition monitoring 2009-10. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2010/000370-2, Adelaide.

Gehrig, S.L., Nicol, J.M. and Marsland, K.B. (2011b). Lower Lakes vegetation condition monitoring – 2010/2011. South Australian Research and Development Institute (Aquatic Sciences), F2009/000370-3, Adelaide.

Hart, B.T., Bailey, P., Edwards, R., Hortle, K.G., James, K., McMahon, A., Meredith, C. and Swadling, K. (1991). A review of the salt sensitivity of the Australian freshwater biota. *Hydrobiologia* **210**: 150-144.

Heard, L. and Channon, B. (1997). Guide to a native vegetation survey using the biological survey of South Australia. South Australian Department of Environment and Natural Resources, Adelaide.

Holt, M., Swingler, K., O'Donnell, E., Shirley, M., Lake, M., Conallin, A., Meredith, S., Ho, S., Prider, J., Poulsen, D., Richardson, S. and Cooling, M. (2005). River Murray Wetlands Baseline Survey. River Murray Catchment Water Management Board, Berri.

James, W.F., Best, E.P. and Barko, J.W. (2004). Sediment resuspension and light attenuation in Peoria Lake: can macrophytes improve water quality in this shallow system? *Hydrobiologia* **515**: 193-201.

Jessop, J., Dashorst, G.R.M. and James, F.R. (2006). 'Grasses of South Australia. An illustrated guide to the native and naturalised species.' (Wakefield Press: Adelaide).

Jessop, J.P. and Tolken, H.R. (1986). 'The Flora of South Australia.' (Government of South Australia Printer: Adelaide).

Kingsford, R.T., Fairweather, P.G., Geddes, M.C., Lester, R.E., Sammut, J. and Walker, K.F. (2009). Engineering a crisis in a Ramsar Wetland: the Coorong, Lower Lakes and Murray Mouth, Australia. Australian Wetlands and Rivers Centre, The University of New South Wales, Sydney.

Kingsford, R.T., Walker, K.F., Lester, R.E., Young, W.J., Fairweather, P.G., Sammut, J. and Geddes, M.C. (2011). A Ramsar wetland in crisis – the Coorong, Lower Lakes and Murray Mouth, Australia. *Marine and Freshwater Research* **62**: 255-265.

Marsland, K. and Nicol, J. (2006). Current and proposed monitoring activities in relation to The Living Murray Lower Lakes, Coorong and Murray Mouth Icon Site Environmental Management Plan. SARDI Aquatic Sciences, SARDI Publication Number F2006/000081 P01, Adelaide.

Marsland, K.B. and Nicol, J.M. (2009). Lower Lakes vegetation condition monitoring-2008/09. South Australian Research and Development Institute (Aquatic Sciences), F2009/000370-1, Adelaide.

Matuszak, A., Mörtl, M., Quillfeldt, P. and Bauer, H.G. (2014). Macrophyte-associated macroinvertebrates as an important food source for wintering waterbirds at Lake Constance. *Limnology* **15**: 69-76.

Maunsell Australia Pty Ltd (2009). Lower Lakes, Coorong and Murray Mouth Icon Site condition monitoring plan. South Australian Murray-Darling Basin Natural Resources Management Board, Adelaide.

Merry, R.H., Fitzpatrick, R.W., Barnett, E.J., Davies, P.J., Fotheringham, D.G., Thomas, B.P. and Hicks, W.S. (2003). South Australian inventory of acid sulfate soil risk (atlas). CSIRO Land and Water, Adelaide.

Nicol, J.M. (2004). Vegetation Dynamics of the Menindee Lakes with Reference to the Seed Bank. PhD thesis, The University of Adelaide.

Nicol, J.M. (2016). An assessment of Ramsar criteria and Limits of Acceptable Change for aquatic and littoral vegetation in the Coorong and Lakes Alexandrina and Albert Ramsar Wetland. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2016/000262-1, Adelaide.

Nicol, J.M., Frahn, K.A., Gehrig, S.L. and Marsland, K.B. (2016a). Lower Lakes Vegetation Condition Monitoring – 2015-16. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2009/000370-7, Adelaide.

Nicol, J.M., Frahn, K.A., Gehrig, S.L. and Marsland, K.B. (2017). Lower Lakes Vegetation Condition Monitoring – 2016-17. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2009/000370-8, Adelaide.

Nicol, J.M., Frahn, K.A., Gehrig, S.L. and Marsland, K.B. (2019a). Lower Lakes Vegetation Condition Monitoring – 2017-18. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2009/000370-9, Adelaide.

Nicol, J.M., Frahn, K.A., Gehrig, S.L. and Marsland, K.B. (2019b). Lower Lakes Vegetation Condition Monitoring – 2018-19. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2009/000370-10, Adelaide.

Nicol, J. M., Frahn, K.A., Gehrig, S.L. and Marsland, K.B. (2020). Lower Lakes Vegetation Condition Monitoring – 2019-20. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2009/000370-11, Adelaide.

Nicol, J.M., Ganf, G.G. and Pelton, G.A. (2003). Seed banks of a southern Australian wetland: the influence of water regime on final species composition. *Plant Ecology* **168**: 191-205.

Nicol, J.M., Gehrig, S.L. and Frahn, K.A. (2016b). Establishment success and benefits to the aquatic plant community of planting *Schoenoplectus tabernaemontani* around the shorelines of lakes Alexandrina and Albert 2013 – 2016. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2013/000414-4, Adelaide.

Nicol, J.M., Weedon, J.T. and Doonan, A. (2006). Vegetation Surveys. In 'River Murray Wetlands Baseline Survey – 2005'. (Eds Simpson, D., Holt, M., Champion, T., Horan, A. and Shirley, M.). (South Australian Murray-Darling Basin Natural Resources Management Board: Berri).

Nielsen, D.L. and Brock, M.A. (2009). Modified water regime and salinity as a consequence of climate change: prospects for wetlands of Southern Australia. *Climate Change* **95**: 532-533.

Nielsen, D.L., Brock, M.A., Petrie, R. and Crossle, K. (2007). The impact of salinity pulses on the emergence of plant and zooplankton from wetland seed and egg banks. *Freshwater Biology* **52**: 784-795.

Nielsen, D.L., Brock, M.A., Rees, G.N. and Baldwin, D.S. (2003). Effects of increasing salinity on freshwater ecosystems in Australia. *Australian Journal of Botany* **51**: 655 - 665

Nielsen, D.L. and Chick, A.J. (1997). Flood-mediated changes in aquatic macrophyte community structure. *Marine and Freshwater Research* **48**: 153-157.

Noges, T., Luup, H. and Feldmann, T. (2010). Primary production of aquatic macrophytes and their epiphytes in two shallow lakes (Peipsi and Vortsjarv) in Estonia. *Aquatic Ecology* **44**: 83-92.

Papas, P. (2007). Effect of macrophytes on aquatic invertebrates – a literature review. Arthur Rylah Institute for Environmental Research, Department of Sustainability and Environment and Melbourne Water, Technical Report Series No. 158, Melbourne.

Phillips, B. and Muller, K. (2006). Ecological character of the Coorong, Lakes Alexandrina and Albert Wetland of International Importance. Department for Environment and Heritage, Adelaide.

PIRSA Spatial Information Services (2009). Erosion mapping of the shorelines of Lakes Alexandrina and Albert. Primary Industries and Resources South Australia (Spatial Information Services) and Goolwa to Wellington Local Action Planning Board Adelaide.

Prescott, A. (1988). 'It's Blue with Five Petals. Wild Flowers of the Adelaide Region.' (Ann Prescott: Prospect, South Australia).

Renfrey, A.P.C., Rea, N. and Ganf, G.G. (1989). The aquatic flora of Hindmarsh Island, South Australia. Department of Environment and Planning, Adelaide.

Robinson, W.A. (2015) The Living Murray Condition Monitoring Plan Refinement Project: Summary Report. Technical Report to the MDBA, March 2015. 95 pp.

Romanowski, N. (1998). 'Aquatic and Wetland Plants. A Field Guide for Non-tropical Australia.' (University of New South Wales Press: Sydney).

Sainty, G.R. and Jacobs, S.W.L. (1981). 'Water Plants of New South Wales.' (Water Resources Commission New South Wales: Sydney).

Sainty, G.R. and Jacobs, S.W.L. (2003). 'Waterplants in Australia.' (Sainty and Associates: Darlinghurst, N.S.W., Australia).

Seaman, R.L. (2003). Coorong and Lower Lakes Ramsar habitat mapping program. South Australian Department for Environment and Heritage, Adelaide.

van Dijk, A.I.J.M., Beck, H.E., Crosbie, R.S., de Jeu, R.A.M., Liu, Y.Y., Podger, G.M., Timbal, B. and Viney, N.R. (2013). The Millennium Drought in southeast Australia (2001–2009): natural and human causes and implications for water resources, ecosystems, economy, and society. *Water Resources Research* **49**: 1040-1057.

Walker, P.D., Wijnhoven, S. and van der Velde, G. (2013). Macrophyte presence and growth form influence macroinvertebrate community structure. *Aquatic Botany* **104**: 80-87.

Webster, I.T., Parslow, J.S., Grayson, R.B., Molloy, R.P., Andrewartha, J., Sakov, P., Tan, K.S., Walker, S.J. and Wallace, B.B. (2001). Gippsland Lakes environmental study: assessing options for improving water quality and ecological function. Gippsland Coastal Board, Melbourne.

Wedderburn, S.D., Walker, K.F. and Zampatti, B.P. (2007). Habitat separation of *Craterocephalus* (Atherinidae) species and populations in off-channel areas of the lower River Murray, Australia. *Ecology of Freshwater Fish* **16**: 442–449

Wright, J.F., Gunn, R.J.M., Winder, J.M., Wiggers, R., Vowles, K., Clarke, R.T. and Harris, I. (2002). A comparison of the macrophyte cover and macroinvertebrate fauna at three sites on the River Kennet in the mid 1970s and late 1990s. *The Science of The Total Environment* **282-283**: 121-142.

APPENDICES

Appendix 1: Species list, functional classification (Gehrig and Nicol 2010b), life history strategy and conservation status (state conservation status from listings in Barker *et al.* (2005) and regional conservation status from listings in Lang and Kaeheneuhl (2001) from all sites and survey dates (*denotes exotic taxon, **denotes proclaimed pest plant in South Australia, ***denotes weed of national significance # denotes listed as rare in South Australia).

Taxon	Functional Group	Life history strategy	Status and Comments
<i>Acacia myrtifolia</i>	Terrestrial dry	Perennial	Native
<i>Anagallis arvensis</i> *	Terrestrial damp	Annual	Exotic
<i>Apium graveolens</i> *	Terrestrial damp	Annual	Exotic
<i>Arctotheca calendula</i> *	Terrestrial dry	Annual	Exotic
<i>Asparagus asparagoides</i> ***	Terrestrial dry	Perennial	Exotic
<i>Asparagus officinalis</i> *	Terrestrial dry	Perennial	Exotic
<i>Atriplex prostrata</i> *	Terrestrial damp	Perennial	Exotic
<i>Atriplex</i> spp.	Terrestrial dry	Perennial	Native
<i>Atriplex suberecta</i>	Floodplain	Perennial	Native
<i>Avena</i> spp.*	Terrestrial dry	Annual	Exotic- <i>Avena</i> spp. is comprised of <i>Avena barbata</i> and <i>Avena fatua</i>
<i>Azolla filiculoides</i>	Floating	Perennial	Native
<i>Baumea juncea</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Berula erecta</i> *	Emergent	Perennial	Exotic
<i>Bolboschoenus caldwellii</i>	Emergent	Perennial	Native
<i>Brassica rapa</i> *	Terrestrial dry	Annual	Exotic
<i>Brassica tournifortii</i> *	Terrestrial dry	Annual	Exotic
<i>Briza minor</i> *	Terrestrial dry	Annual	Exotic
<i>Bromus catharticus</i> *	Terrestrial dry	Annual	Exotic
<i>Bromus diandrus</i> *	Terrestrial dry	Annual	Exotic
<i>Bromus hordeaceus</i> ssp. <i>hordeaceus</i> *	Terrestrial dry	Annual	Exotic
<i>Bromus rubens</i> *	Terrestrial dry	Annual	Exotic
<i>Calystegia sepium</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native-Listed as Uncommon in the Murray and Southern Lofty Regions
<i>Carex apressa</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Carex fascicularis</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Cenchrus clandestinus</i>	Terrestrial dry	Perennial	Exotic
<i>Centaurea calcitrapa</i> *	Terrestrial damp	Annual	Exotic
<i>Centaureum tenuiflorum</i> *	Terrestrial damp	Annual	Exotic
<i>Centella asiatica</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Ceratophyllum demersum</i> #	Submergent (k-selected)	Perennial	Native-Listed as Rare in South Australia
<i>Chara</i> spp.	Submergent (r-selected)	Annual	Native
<i>Chenopodium album</i> *	Terrestrial damp	Annual	Exotic
<i>Chenopodium glaucum</i> *	Terrestrial damp	Annual	Exotic
<i>Chenopodium nitrariaceum</i>	Terrestrial dry	Perennial	Native
<i>Conyza bonariensis</i> *	Terrestrial damp	Annual	Exotic
<i>Cotula coronopifolia</i> *	Amphibious fluctuation responder-plastic	Perennial	Exotic
<i>Crassula helmsii</i>	Amphibious fluctuation tolerator-low growing	Perennial	Native
<i>Cyanogeton procera</i>	Emergent	Perennial	Native-Listed as Uncommon in the Southern Lofty Region
<i>Cyperus exaltatus</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Cyperus gymnocaulos</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Dianella revoluta</i>	Terrestrial dry	Perennial	Native
<i>Disphyma crassifolium</i>	Terrestrial dry	Perennial	Native

Taxon	Functional Group	Life history strategy	Status and Comments
<i>Distichlis distichophylla</i>	Terrestrial damp	Perennial	Native-Listed as Uncommon in the Murray Region
<i>Duma florulenta</i>	Amphibious fluctuation tolerator-woody	Perennial	Native
<i>Echinochloa crus-galli</i> *	Terrestrial damp	Annual	Exotic
<i>Ehrharta longiflora</i> *	Terrestrial damp	Annual	Exotic
<i>Einadia nutans</i>	Terrestrial dry	Perennial	Native
<i>Eleocharis acuta</i>	Emergent	Perennial	Native
<i>Enchylaena tomentosa</i>	Terrestrial dry	Perennial	Native
<i>Epilobium pallidiflorum</i>	Terrestrial damp	Perennial	Native-Listed as Uncertain in the Murray Region and uncommon in the Southern Lofty Region
<i>Eragrostis australasica</i>	Floodplain	Perennial	Native
<i>Eragrostis curvula</i> **	Terrestrial damp	Annual	Exotic-Proclaimed pest plant in SA
<i>Eragrostis</i> sp.	Terrestrial damp	Annual	Native-could not identify to species
<i>Erodium cicutarium</i> *	Terrestrial dry	Annual	Exotic
<i>Euphorbia terracina</i> **	Terrestrial dry	Annual	Exotic-Proclaimed pest plant in SA
<i>Ficinia nodosa</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Foeniculum vulgare</i> *	Terrestrial damp	Annual	Exotic
<i>Frankenia pauciflora</i>	Terrestrial dry	Perennial	Native
<i>Fumaria bastardii</i> *	Terrestrial damp	Annual	Exotic
<i>Gahnia clarkii</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Gahnia filum</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native-Listed as Rare in the Murray and Southern Lofty Regions
<i>Galenia secunda</i> *	Terrestrial dry	Perennial	Exotic
<i>Glyceria australis</i>	Emergent	Perennial	Native
<i>Heliotropium europaeum</i> *	Floodplain	Annual	Exotic
<i>Holcus lanatus</i> *	Terrestrial damp	Annual	Exotic
<i>Hordeum vulgare</i> *	Terrestrial dry	Annual	Exotic
<i>Hypochoeris glabra</i> *	Terrestrial dry	Annual	Exotic
<i>Hypochoeris radicata</i> *	Terrestrial dry	Annual	Exotic
<i>Iris</i> spp.*	Terrestrial dry	Perennial	Exotic
<i>Isolepis producta</i>	Amphibious fluctuation tolerator-low growing	Perennial	Native
<i>Juncus acutus</i> **	Amphibious fluctuation tolerator-emergent	Perennial	Exotic
<i>Juncus holoschoenus</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Juncus kraussii</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Juncus subsecundus</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Juncus usitatus</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Lachnagrostis filiformis</i>	Floodplain	Annual	Native
<i>Lactuca saligna</i> *	Terrestrial dry	Annual	Exotic
<i>Lactuca serriola</i> *	Terrestrial dry	Annual	Exotic
<i>Lagurus ovatus</i> *	Terrestrial dry	Annual	Exotic
<i>Lamprothamnium macropogon</i>	Submergent r-selected	Annual	Native
<i>Lemna</i> spp.	Floating	Perennial	Native
<i>Limosella australis</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Lobelia anceps</i>	Terrestrial damp	Perennial	Native
<i>Lolium</i> spp.*	Terrestrial dry	Annual	Exotic- <i>Lolium</i> spp. comprises of <i>Lolium perenne</i> and <i>Lolium rigidum</i>
<i>Ludwigia peploides</i> ssp. <i>montevidensis</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Lupinus cosentinii</i> *	Terrestrial dry	Annual	Exotic

Taxon	Functional Group	Life history strategy	Status and Comments
<i>Lycium ferocissimum</i> ***	Terrestrial dry	Perennial	Exotic-Proclaimed pest plant in SA
<i>Lycopus australis</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native-Listed as Rare in the Murray Region
<i>Lythrum hyssopifolia</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Lythrum salicaria</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Malva parviflora</i> *	Terrestrial dry	Annual	Exotic
<i>Marrubium vulgare</i> **	Terrestrial dry	Annual	Exotic
<i>Medicago</i> spp.*	Terrestrial dry	Annual	Exotic- <i>Medicago</i> spp. comprises of <i>Medicago polymorpha</i> , <i>Medicago truncatula</i> and <i>Medicago minima</i>
<i>Melaleuca halmaturorum</i>	Amphibious fluctuation tolerator-woody	Perennial	Native
<i>Melilotus albus</i> *	Terrestrial dry	Annual	Exotic
<i>Melilotus indicus</i> *	Terrestrial dry	Annual	Exotic
<i>Mentha australis</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Mentha</i> spp.*	Amphibious fluctuation tolerator-emergent	Perennial	Exotic- <i>Mentha</i> spp. comprises of <i>Mentha piperita</i> , <i>Mentha pulegium</i> and <i>Mentha spicata</i>
<i>Myoporum insulare</i>	Terrestrial dry	Perennial	Native
<i>Myriophyllum caput-medusae</i>	Submergent k-selected	Perennial	Native
<i>Myriophyllum muelleri</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Myriophyllum salsugineum</i>	Submergent k-selected	Perennial	Native-Listed as Uncertain in the Southern Lofty Region
<i>Myriophyllum verrucosum</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Onopordum acanthium</i> *	Terrestrial damp	Annual	Exotic
<i>Paspalum distichum</i> *	Terrestrial damp	Perennial	Exotic
<i>Persicaria lapathifolia</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Phragmites australis</i>	Emergent	Perennial	Native
<i>Phyla canescens</i> *	Amphibious fluctuation tolerator-low growing	Perennial	Exotic
<i>Picris angustifolia</i> ssp. <i>angustifolia</i>	Terrestrial dry	Annual	Native
<i>Plantago coronopus</i> *	Terrestrial dry	Annual	Exotic
<i>Plantago lanceolata</i> *	Terrestrial dry	Annual	Exotic
<i>Polygonum aviculare</i> *	Terrestrial dry	Perennial	Exotic
<i>Polypogon monspeliensis</i> *	Amphibious fluctuation tolerator-emergent	Annual	Exotic
<i>Potamogeton crispus</i>	Submergent k-selected	Perennial	Native
<i>Potamogeton pectinatus</i>	Submergent k-selected	Perennial	Native
<i>Pseudognaphalium luteoalbum</i>	Floodplain	Annual	Native
<i>Puccinellia</i> sp.*	Terrestrial damp	Annual	Exotic-could not be identified to species but was not <i>Puccinellia stricta</i> or <i>Puccinellia perlaxa</i>
<i>Ranunculus trichophyllus</i> *	Submergent (r-selected)	Annual	Exotic
<i>Ranunculus trilobus</i> *	Amphibious fluctuation tolerator-emergent	Annual	Exotic
<i>Reichardia tingitana</i> *	Terrestrial dry	Annual	Exotic
<i>Rorippa nasturtium-aquaticum</i> *	Amphibious fluctuation responder-plastic	Annual	Exotic
<i>Rorippa palustris</i> *	Floodplain	Annual	Exotic
<i>Rumex bidens</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Ruppia megacarpa</i>	Submergent k-selected	Perennial	Native
<i>Ruppia polycarpa</i>	Submergent r-selected	Annual	Native
<i>Ruppia tuberosa</i>	Submergent r-selected	Annual	Native
<i>Salix babylonica</i> *	Emergent	Perennial	Exotic
<i>Salsola australis</i>	Terrestrial dry	Perennial	Native

Taxon	Functional Group	Life history strategy	Status and Comments
<i>Samolus repens</i>	Terrestrial damp	Perennial	Native- Listed as Rare in the Murray Region and Uncommon the Southern Lofty Region
<i>Sarcocornia quinqueflora</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Scabiosa atropurpurea</i> *	Terrestrial dry	Annual	Exotic
<i>Scaevola calendulacea</i>	Terrestrial dry	Perennial	Native
<i>Schoenoplectus pungens</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native-Listed as Rare in the Southern Lofty Region
<i>Schoenoplectus tabernaemontani</i>	Emergent	Perennial	Native
<i>Sclerolaena blackiana</i>	Terrestrial dry	Perennial	Native-Listed as Rare in SA
<i>Senecio cunninghamii</i>	Floodplain	Perennial	Native
<i>Senecio pterophorus</i> *	Terrestrial dry	Annual	Exotic
<i>Senecio runcinifolius</i>	Floodplain	Perennial	Native-Listed as Uncommon in the Murray Region
<i>Silybum marianum</i> **	Terrestrial damp	Annual	Exotic-Proclaimed pest plant in SA
<i>Solanum lycopersicum</i> *	Terrestrial dry	Annual	Exotic
<i>Solanum nigrum</i> *	Terrestrial damp	Annual	Exotic
<i>Sonchus asper</i> *	Terrestrial damp	Annual	Exotic
<i>Sonchus oleraceus</i> *	Terrestrial damp	Annual	Exotic
<i>Spergularia brevifolia</i> *	Terrestrial damp	Annual	Exotic
<i>Stenotaphrum secundatum</i> *	Terrestrial dry	Perennial	Exotic
<i>Suaeda australis</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Symphotrichum subulatum</i> *	Terrestrial damp	Annual	Exotic
<i>Thyridia repens</i>	Amphibious fluctuation tolerator-low growing	Perennial	Native
<i>Trifolium</i> spp.*	Terrestrial dry	Annual	Exotic- <i>Trifolium</i> spp. comprises of <i>Trifolium angustifolium</i> , <i>Trifolium arvense</i> , <i>Trifolium repens</i> and <i>Trifolium subterraneum</i>
<i>Triglochin striata</i>	Amphibious fluctuation tolerator-low growing	Perennial	Native
<i>Triticum</i> sp.*	Terrestrial dry	Annual	Exotic-could not be identified to species
<i>Typha domingensis</i>	Emergent	Perennial	Native
<i>Urtica urens</i> *	Terrestrial damp	Annual	Exotic
<i>Vallisneria australis</i>	Submergent k-selected	Perennial	Native-Listed as Uncommon in the Murray Region and Threatened in the Southern Lofty Region
<i>Vicia sativa</i> *	Terrestrial dry	Annual	Exotic
<i>Wilsonia rotundifolia</i>	Terrestrial damp	Perennial	Native

Appendix 2: GPS coordinates (UTM format, map datum WGS84) for lakeshore and wetland understory vegetation monitoring sites (site numbers correspond with site numbers in Figure 1).

Site #	Site	Easting	Northing	Site type
1	Bremer Mouth Lakeshore	323061	6081991	lakeshore
2	Brown Beach 1	350172	6052777	lakeshore
3	Brown Beach 2	350287	6053158	lakeshore
4	Clayton Bay	311301	6070626	lakeshore
5	Currency Creek 3	296772	6074222	lakeshore
6	Currency Creek 4	301013	6071800	lakeshore
7	Goolwa North	303330	6070156	lakeshore
8	Goolwa South	300490	6066366	lakeshore
9	Hindmarsh Island Bridge 01	299670	6068521	lakeshore
10	Hindmarsh Island Bridge 02	299695	6068616	lakeshore
11	Lake Reserve Rd	339298	6089987	lakeshore
12	Loveday Bay	329431	6058407	lakeshore
13	Loveday Bay Lakeshore	326621	6061647	lakeshore
14	Lower Finnis 02	305131	6076401	lakeshore
15	Milang	315964	6079870	lakeshore
16	Milang Lakeshore	316081	6079746	lakeshore
17	Pt Sturt Lakeshore	322811	6069643	lakeshore
18	Pt Sturt Water Reserve	317673	6070784	lakeshore
19	Terlingie Lakeshore	327461	6066887	lakeshore
20	Upstream of Clayton Regulator	312281	6069151	lakeshore
21	Wally's Landing	303066	6079631	lakeshore
22	Warrengie 1	347722	6049163	lakeshore
23	Lower Finnis 03	305131	6072406	lakeshore
24	Narrung Lakeshore	333762	6069807	lakeshore
25	Nurra Nurra	341786	6063837	lakeshore
26	Warrengie 2	348487	6049133	lakeshore
27	Angas Mouth	318391	6081206	wetland
28	Bremer Mouth	323056	6082019	wetland
29	Dunns Lagoon	312417	6070300	wetland
30	Goolwa Channel Drive	307024	6064437	wetland
31	Hunters Creek	308219	6065526	wetland
32	Poltalloch	343248	6071554	wetland
33	Pt Sturt	322778	6069794	wetland
34	Terlingie	327334	6065286	wetland
35	Waltowa	353908	6057756	wetland
36	Narrung	334542	6068744	wetland

Appendix 3: Taxa present (green shading) in Lake Alexandrina spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).

Taxon	Survey Date																					
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Apium graveolens</i> *		*	*				*	*	*		*		*	*	*		*	*				
<i>Arctotheca calendula</i> *	*		*																			
<i>Atriplex prostrata</i> *	*	*		*	*																	
<i>Atriplex</i> spp.				*																		
<i>Atriplex suberecta</i>		*	*														*					
<i>Avena</i> spp.*	*		*		*				*													
<i>Azolla filiculoides</i>						*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Berula erecta</i> *	*							*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Bolboschoenus caldwellii</i>		*		*	*	*	*		*	*	*	*	*	*		*	*	*			*	*
<i>Brassica rapa</i> *	*																					
<i>Brassica tournifortii</i> *				*								*										
<i>Briza minor</i> *			*																			
<i>Bromus diandrus</i> *	*		*																			
<i>Bromus hordeaceus</i> ssp. <i>hordeaceus</i> *			*						*													
<i>Bromus rubens</i> *									*													
<i>Calystegia sepium</i>	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Carex apressa</i>															*					*		
<i>Carex fascicularis</i>									*				*				*	*		*		
<i>Cenchrus clandestinus</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Centaurea calcitrapa</i> *	*	*	*	*					*				*	*								*
<i>Centaurium tenuiflorum</i> *	*		*																			
<i>Centella asiatica</i>	*							*	*		*	*	*	*	*	*	*	*	*	*	*	*
<i>Ceratophyllum demersum</i> #						*	*	*	*	*	*	*	*	*	*	*	*	*		*	*	
<i>Chara</i> spp.										*			*									
<i>Chenopodium glaucum</i> *		*		*																		
<i>Chenopodium nitriaceum</i>				*																		
<i>Conyza bonariensis</i> *	*	*	*	*									*	*								
<i>Cotula coronopifolia</i> *	*	*	*		*	*	*	*	*					*	*			*	*	*	*	
<i>Crassula helmsii</i>																*		*	*	*	*	*
<i>Cycnogeton procera</i>					*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Cyperus gymnocaulos</i>	*	*	*	*	*	*	*	*	*	*		*		*			*				*	
<i>Distichlis distichophylla</i>	*		*																			
<i>Duma florulenta</i>	*			*																		
<i>Ehrharta longiflora</i> *			*																			
<i>Einadia nutans</i>		*		*																		
<i>Eleocharis acuta</i>							*		*	*	*	*		*	*	*	*	*	*	*	*	*
<i>Enchylaena tomentosa</i>			*																			
<i>Epilobium pallidiflorum</i>												*		*		*		*			*	*
<i>Eragrostis curvula</i> **	*		*																			
<i>Eragrostis</i> sp.	*	*	*																			
<i>Ficinia nodosa</i>		*	*	*	*							*	*				*					
<i>Foeniculum vulgare</i> *	*	*		*																		
<i>Frankenia pauciflora</i>		*																				
<i>Fumaria bastardii</i> *			*																			
<i>Galenia secunda</i> *													*									
<i>Glyceria australis</i>							*															
<i>Holcus lanatus</i> *	*						*										*					
<i>Hordeum vulgare</i> *	*		*		*																	

Taxon	Survey Date																					
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Hypochoeris glabra</i> *	*		*		*																	
<i>Hypochoeris radicata</i> *	*	*	*	*																		
<i>Isolepis producta</i>	*	*	*						*													*
<i>Juncus acutus</i> **	*	*	*	*																		
<i>Juncus holoschoenus</i>												*		*	*	*						
<i>Juncus kraussii</i>	*	*	*	*	*	*	*	*	*	*	*				*	*						
<i>Juncus usitatus</i>		*	*	*	*	*			*					*			*	*	*	*	*	*
<i>Lachnagrostis filiformis</i>	*	*	*	*		*	*	*			*		*			*						
<i>Lactuca saligna</i> *			*																			
<i>Lactuca serriola</i> *		*	*	*	*						*	*	*									
<i>Lagurus ovatus</i> *			*																			
<i>Lemna</i> spp.						*	*				*	*		*	*	*	*	*				
<i>Limosella australis</i>								*		*				*	*	*	*	*				
<i>Lobelia anceps</i>			*										*					*				
<i>Lolium</i> spp.*	*		*		*		*		*		*			*			*				*	*
<i>Ludwigia peploides</i> ssp. <i>montevidensis</i>						*			*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Lycopus australis</i>		*	*	*			*			*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Lythrum hyssopifolia</i>											*											*
<i>Medicago</i> spp.*	*		*				*	*		*											*	*
<i>Melilotus indicus</i> *	*		*										*							*		
<i>Mentha australis</i>	*							*										*	*	*		
<i>Mentha</i> spp.*		*	*	*	*	*				*	*	*	*	*	*	*	*					
<i>Myriophyllum muelleri</i>																*		*				
<i>Myriophyllum salsugineum</i>							*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Onopordum acanthium</i> *	*																					
<i>Paspalum distichum</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Persicaria lapathifolia</i>	*		*	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Phragmites australis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Picris angustifolia</i> ssp. <i>angustifolia</i>	*	*	*	*																		
<i>Plantago coronopus</i> *	*	*	*	*	*																	
<i>Plantago lanceolata</i> *									*													
<i>Polygonum aviculare</i> *		*	*			*																
<i>Polypogon monspeliensis</i> *	*	*	*	*		*	*		*	*	*			*					*			
<i>Potamogeton crispus</i>											*			*			*		*	*		*
<i>Potamogeton pectinatus</i>									*		*	*	*		*					*	*	*
<i>Pseudognaphalium luteoalbum</i>	*		*	*							*											
<i>Puccinellia</i> sp.*			*																			
<i>Ranunculus trichophyllus</i> *									*			*	*	*	*	*	*	*	*	*	*	*
<i>Reichardia tingitana</i> *	*		*	*																		
<i>Rorippa nasturtium-aquaticum</i> *										*	*				*			*				
<i>Rorippa palustris</i> *	*																					
<i>Rumex bidens</i>	*						*	*	*	*	*		*	*	*	*	*		*	*	*	*
<i>Ruppia tuberosa</i>	*											*	*	*	*	*	*		*	*	*	*
<i>Salix babylonica</i> *	*																					
<i>Sarcocornia quinqueflora</i>	*	*	*	*																		
<i>Schoenoplectus pungens</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*		*		*
<i>Schoenoplectus tabernaemontani</i>	*		*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Sclerolaena blackiana</i>	*																					
<i>Senecio cunninghamii</i>			*																			

Taxon	Survey Date																					
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Senecio pterophorus</i> *	*	*	*	*				*	*	*	*	*		*	*		*	*				
<i>Senecio runcinifolius</i>		*											*									
<i>Silybum marianum</i> **			*																			
<i>Solanum lycopersicum</i> *															*							
<i>Solanum nigrum</i> *		*	*			*		*		*					*							
<i>Sonchus asper</i> *			*	*		*	*	*														
<i>Sonchus oleraceus</i> *	*	*	*	*	*		*		*	*			*		*		*					
<i>Spergularia brevifolia</i> *	*		*	*																		
<i>Suaeda australis</i>	*	*	*	*									*									
<i>Symphytotrichum subulatum</i> *	*	*	*	*		*	*	*	*	*		*	*		*	*		*	*	*		*
<i>Thyridia repens</i>		*										*										
<i>Trifolium spp.</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*		*	*	*	*	*	
<i>Triglochin striata</i>			*	*		*														*		
<i>Triticum sp.</i> *			*																			
<i>Typha domingensis</i>	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Urtica urens</i> *			*	*																		
<i>Vallisneria australis</i>											*	*		*			*			*	*	*
<i>Vicia sativa</i> *	*		*				*										*	*				
<i>Wilsonia rotundifolia</i>		*																				

Appendix 4: Taxa present (green shading) in Lake Albert spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).

Taxon	Survey Date																					
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Acacia myrtifolia</i>		*	*	*																		
<i>Anagallis arvensis*</i>										*												
<i>Arctotheca calendula*</i>			*																			
<i>Avena spp.*</i>	*		*						*													
<i>Bolboschoenus caldwellii</i>									*													
<i>Bromus catharticus*</i>									*													
<i>Bromus diandrus*</i>	*		*																			
<i>Bromus hordeaceus ssp. hordeaceus*</i>			*																			
<i>Calystegia sepium</i>									*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Cenchrus clandestinus*</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Centaurea calcitrapa*</i>	*		*						*													
<i>Chenopodium album*</i>				*									*									
<i>Chenopodium glaucum*</i>				*																		
<i>Conyza bonariensis*</i>	*							*														
<i>Cotula coronopifolia*</i>	*		*	*			*	*														
<i>Cyperus gymnocaulos</i>	*	*	*	*	*		*		*	*		*	*				*	*	*			
<i>Distichlis distichophylla</i>	*																					
<i>Duma florulenta</i>							*					*			*		*	*	*	*	*	*
<i>Ehrharta longiflora*</i>			*																			
<i>Enchylaena tomentosa</i>				*																		
<i>Eragrostis australasica</i>		*		*																		
<i>Eragrostis curvula**</i>			*																			
<i>Euphorbia terracina**</i>			*																			
<i>Ficinia nodosa</i>		*	*	*	*																	
<i>Hordeum vulgare*</i>	*		*																			
<i>Hypochoeris glabra*</i>			*																			
<i>Hypochoeris radicata*</i>			*	*																		
<i>Isolepis producta</i>	*			*									*									
<i>Lachnagrostis filiformis</i>	*	*										*	*									
<i>Lactuca serriola*</i>									*													
<i>Lagurus ovatus*</i>			*																			
<i>Lolium spp.*</i>	*		*						*													
<i>Lythrum hyssopifolia</i>									*			*			*							
<i>Lythrum salicaria</i>												*										
<i>Medicago spp.*</i>	*		*																			
<i>Melaleuca halmaturorum</i>			*	*									*									
<i>Melilotus indicus*</i>	*		*	*														*				
<i>Myriophyllum salsugineum</i>																		*				*
<i>Paspalum distichum*</i>		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Phragmites australis</i>		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Plantago coronopus*</i>			*	*					*		*											
<i>Polypogon monspeliensis*</i>	*	*	*																			

Taxon	Survey Date																						
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021	
<i>Potamogeton pectinatus</i>							*										*						
<i>Puccinellia</i> sp.*			*	*																			
<i>Reichardia tingitana</i> *	*	*	*	*																			
<i>Rumex bidens</i>									*							*						*	
<i>Sarcocornia quinqueflora</i>		*	*	*		*																	
<i>Scaevola calendulacea</i>				*																			
<i>Schoenoplectus pungens</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*		*	*	*			*
<i>Schoenoplectus tabernaemontani</i>	*				*	*			*	*			*		*	*		*	*	*			
<i>Senecio pterophorus</i> *			*																				
<i>Sonchus oleraceus</i> *	*		*	*				*	*			*											
<i>Spergularia brevifolia</i> *			*	*																			
<i>Suaeda australis</i>			*	*																			
<i>Symphyotrichum subulatum</i> *	*		*	*				*				*	*					*		*			*
<i>Thyridia repens</i>	*			*				*		*													*
<i>Trifolium</i> spp.*	*		*	*					*				*						*				
<i>Triglochin striata</i>			*	*																			
<i>Typha domingensis</i>										*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Vicia sativa</i> *	*		*						*														

Appendix 5: Taxa present (green shading) in Goolwa Channel spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).

Taxon	Survey Date																					
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Acacia myrtifolia</i>												*	*	*	*			*		*	*	*
<i>Asparagus asparagoides</i> ***																				*		
<i>Asparagus officinalis</i> *		*							*													
<i>Atriplex prostrata</i> *		*		*								*										
<i>Atriplex</i> spp.		*																				
<i>Azolla filiculoides</i>			*			*	*	*	*	*	*	*	*	*	*	*	*	*		*	*	*
<i>Berula erecta</i> *						*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Bolboschoenus caldwellii</i>			*	*	*	*	*	*		*	*							*	*	*	*	*
<i>Bouteloua dactyloides</i> *														*			*				*	*
<i>Brassica tournifortii</i> *			*																			
<i>Bromus diandrus</i> *			*																			
<i>Bromus hordeaceus</i> ssp. <i>hordeaceus</i> *			*				*															
<i>Calystegia sepium</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Cenchrus clandestinus</i> *			*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Centaurea calcitrapa</i> *	*																					*
<i>Centella asiatica</i>							*		*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Ceratophyllum demersum</i> #						*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Chenopodium glaucum</i> *	*	*		*																		
<i>Conyza bonariensis</i> *			*																			
<i>Cotula coronopifolia</i> *	*		*	*	*		*				*											
<i>Crassula helmsii</i>																				*	*	*
<i>Cynogeton procera</i>			*	*	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Cyperus exaltatus</i>			*																			
<i>Cyperus gymnocaulos</i>	*		*		*																	
<i>Duma florulenta</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Eleocharis acuta</i>				*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Enchylaena tomentosa</i>			*										*									
<i>Epilobium pallidiflorum</i>		*																				
<i>Eragrostis</i> sp.									*													
<i>Ficinia nodosa</i>			*		*		*				*				*		*	*	*	*		
<i>Gahnia clarkii</i>																						*
<i>Juncus kraussii</i>	*	*		*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Juncus usitatus</i>			*	*		*	*															
<i>Lachnagrostis filiformis</i>	*	*	*																			
<i>Lactuca saligna</i> *							*		*													
<i>Lemna</i> spp.				*		*					*	*		*	*		*					
<i>Lobelia anceps</i>													*									
<i>Lolium</i> spp.*			*			*																
<i>Lupinus cosentinii</i> *			*																			
<i>Lycopus australis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Lythrum hyssopifolia</i>									*													
<i>Lythrum salicaria</i>	*																*					
<i>Medicago</i> spp.*							*				*											
<i>Melilotus indicus</i> *			*				*															
<i>Mentha australis</i>							*		*									*	*	*	*	*
<i>Mentha</i> spp.*			*			*		*	*	*	*	*	*	*	*	*	*					
<i>Myriophyllum caput-medusae</i>									*						*							
<i>Myriophyllum muelleri</i>																		*				
<i>Myriophyllum salsugineum</i>				*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Paspalum distichum</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Persicaria lapathifolia</i>											*	*		*		*	*	*	*	*	*	*

Taxon	Survey Date																					
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Phragmites australis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Picris angustifolia</i> ssp. <i>angustifolia</i>		*		*		*		*				*										
<i>Plantago coronopus</i> *			*		*																	
<i>Plantago lanceolata</i> *			*		*			*	*	*												
<i>Polygonum aviculare</i> *				*																		
<i>Polypogon monspeliensis</i> *	*																					
<i>Potamogeton crispus</i>					*	*								*				*			*	*
<i>Potamogeton pectinatus</i>				*	*				*				*								*	
<i>Ranunculus trilobus</i> *							*		*		*	*	*	*	*							*
<i>Rumex bidens</i>				*	*		*	*	*		*	*		*	*		*		*	*		
<i>Salix babylonica</i> *				*	*	*		*	*	*												
<i>Samolus repens</i>									*	*	*			*			*				*	*
<i>Scabiosa atropurpurea</i> *			*																			
<i>Schoenoplectus pungens</i>			*	*	*		*		*		*	*										
<i>Schoenoplectus tabernaemontani</i>	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Silybum marianum</i> **			*																			
<i>Solanum nigrum</i> *			*																			
<i>Sonchus oleraceus</i> *	*		*				*	*	*		*											
<i>Suaeda australis</i>			*																			
<i>Symphotrichum subulatum</i> *	*	*	*	*				*	*	*	*		*	*		*	*	*				*
<i>Thyridia repens</i>				*																		
<i>Trifolium</i> spp.*												*										
<i>Triglochin striata</i>		*																				
<i>Typha domingensis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Vallisneria australis</i>							*	*		*	*	*					*	*	*		*	*

Appendix 6: Taxa present (green shading) in permanent wetlands spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).

Taxon	Survey Date																						
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021	
<i>Atriplex prostrata</i> *	*	*	*	*	*					*						*		*					
<i>Avena</i> spp.*			*		*												*						
<i>Azolla filiculoides</i>						*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Baumea juncea</i>																		*	*	*	*	*	
<i>Berula erecta</i> *	*													*	*	*	*	*	*	*	*	*	
<i>Bolboschoenus caldwellii</i>			*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
<i>Brassica rapa</i> *	*																						
<i>Brassica tournifortii</i> *							*																
<i>Bromus diandrus</i> *	*		*		*		*										*						
<i>Bromus hordeaceus</i> ssp. <i>hordeaceus</i> *	*		*														*						
<i>Bromus rubens</i> *																				*			
<i>Calystegia sepium</i>				*								*				*				*	*	*	
<i>Cenchrus clandestinus</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
<i>Centaurea calcitrapa</i> *	*	*			*	*														*			
<i>Centella asiatica</i>	*		*									*			*	*		*	*				
<i>Ceratophyllum demersum</i> #				*	*	*	*	*	*	*		*	*		*	*	*	*	*	*	*	*	
<i>Chenopodium album</i> *	*		*	*		*																	
<i>Chenopodium glaucum</i> *		*																					
<i>Conyza bonariensis</i> *	*				*																		
<i>Cotula coronopifolia</i> *	*		*		*	*	*	*	*		*	*											
<i>Crassula helmsii</i>											*		*	*				*	*	*		*	
<i>Cynogeton procera</i>	*		*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
<i>Cyperus exaltatus</i>				*																			
<i>Cyperus gymnocaulos</i>				*	*		*		*			*											
<i>Dianella revoluta</i>									*														
<i>Disphyma crassifolium</i>		*														*	*	*					
<i>Distichlis distichophylla</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
<i>Duma florulenta</i>					*		*																
<i>Echinochloa crus-galli</i> *									*														
<i>Eleocharis acuta</i>					*	*	*	*	*	*	*	*		*				*	*	*	*	*	
<i>Enchylaena tomentosa</i>			*																		*	*	
<i>Eragrostis curvula</i> **	*		*		*		*		*								*						
<i>Eragrostis</i> sp.	*		*				*		*														
<i>Ficinia nodosa</i>		*		*		*																	
<i>Gahnia filum</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
<i>Hordeum vulgare</i> *	*		*		*	*	*	*	*								*						
<i>Hypochoeris glabra</i> *			*			*		*															
<i>Hypochoeris radicata</i> *			*	*																			
<i>Iris</i> spp.*					*																		
<i>Isolepis producta</i>			*			*																	
<i>Juncus acutus</i> **			*						*														
<i>Juncus kraussii</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
<i>Juncus subsecundus</i>							*																
<i>Lachnagrostis filiformis</i>	*	*	*	*		*	*	*				*											

Taxon	Survey Date																					
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Lactuca saligna</i> *	*																					
<i>Lactuca serriola</i> *	*				*	*	*	*	*			*		*		*	*	*				
<i>Lemna</i> spp.					*		*			*	*	*	*	*	*		*		*	*	*	*
<i>Lobelia anceps</i>																						*
<i>Lolium</i> spp.*	*		*		*	*	*		*								*		*	*	*	
<i>Ludwigia peploides</i> ssp. <i>montevidensis</i>						*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Lycium ferocissimum</i> ***																		*				
<i>Lycopus australis</i>										*					*		*					
<i>Medicago</i> spp.*			*		*										*					*		
<i>Melaleuca halmaturorum</i>	*	*	*	*		*	*				*		*	*	*	*	*	*	*	*	*	*
<i>Meililotus indicus</i> *	*		*														*		*			
<i>Mentha australis</i>																	*				*	*
<i>Mentha</i> spp.*	*											*	*	*		*	*					
<i>Myoporum insulare</i>																	*			*	*	*
<i>Myriophyllum caput-medusae</i>									*	*		*	*									
<i>Myriophyllum muelleri</i>																		*				
<i>Myriophyllum salsugineum</i>					*	*	*	*		*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Paspalum distichum</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
<i>Persicaria lapathifolia</i>	*	*	*	*	*			*	*		*		*	*		*	*					
<i>Phragmites australis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Picris angustifolia</i> ssp. <i>angustifolia</i>				*	*																	
<i>Plantago coronopus</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Polygomon montpelienis</i> *	*				*		*		*		*			*								
<i>Potamogeton crispus</i>					*		*		*		*		*	*					*		*	*
<i>Potamogeton pectinatus</i>					*	*	*	*	*	*	*	*	*	*	*	*	*		*	*		
<i>Pseudognaphalium luteoalbum</i>			*	*			*															
<i>Reichardia tingitana</i> *	*		*	*		*	*	*														
<i>Rorippa nasturtium-aquaticum</i> *	*																		*			
<i>Rumex bidens</i>	*	*			*		*	*	*					*	*	*	*	*	*	*	*	*
<i>Ruppia megacarpa</i>					*	*	*															
<i>Ruppia polycarpa</i>					*																	
<i>Ruppia tuberosa</i>	*		*																			
<i>Samolus repens</i>	*		*		*					*	*	*	*	*	*	*	*		*	*	*	*
<i>Sarcocornia quinqueflora</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Scabiosa atropurpurea</i> *						*																
<i>Schoenoplectus pungens</i>	*		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Schoenoplectus tabernaemontani</i>						*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Senecio pterophorus</i> *	*	*	*	*		*			*	*	*	*										
<i>Senecio runcinifolius</i>			*																			
<i>Sonchus asper</i> *			*	*	*																	
<i>Sonchus oleraceus</i> *	*	*	*	*	*		*		*		*	*		*	*	*	*	*	*	*	*	*

Taxon	Survey Date																					
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Spergularia brevifolia</i> *	*	*	*	*																		
<i>Suaeda australis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Symphotrichum subulatum</i> *	*	*	*	*	*	*	*	*	*	*		*		*		*		*	*	*		*
<i>Thyridia repens</i>	*		*			*	*				*											
<i>Trifolium spp.</i> *			*														*					
<i>Triglochin striata</i>	*	*	*		*	*			*		*											*
<i>Typha domingensis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Urtica urens</i> *	*																					
<i>Vallisneria australis</i>	*		*			*	*	*	*	*	*	*	*	*	*	*	*	*	*		*	*
<i>Wilsonia rotundifolia</i>	*	*			*	*							*	*	*		*	*				

Appendix 7: Taxa present (green shading) in temporary wetlands spring 2008 to autumn 2021 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).

Taxon	Survey Date																						
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Spring 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Atriplex prostrata</i> *			*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Atriplex</i> spp.	*	*		*		*	*				*												
<i>Atriplex suberecta</i>		*	*	*	*								*		*			*					
<i>Avena</i> spp.*	*		*				*							*			*		*	*	*	*	*
<i>Azolla filiculoides</i>							*		*					*	*		*			*	*	*	*
<i>Berula erecta</i> *																	*						
<i>Bolboschoenus caldwellii</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Brassica tournfortii</i> *	*																						
<i>Bromus diandrus</i> *	*		*		*		*		*														
<i>Bromus hordeaceus</i> ssp. <i>hordeaceus</i> *			*				*		*														
<i>Bromus rubens</i> *																					*		
<i>Cenchrus clandestinus</i> *													*										
<i>Centaurea calcitrapa</i> *			*		*	*	*	*	*		*										*	*	
<i>Centaureum tenuiflorum</i> *																					*		
<i>Chara</i> spp.						*	*	*	*					*				*					
<i>Chenopodium album</i> *						*																	
<i>Chenopodium glaucum</i> *	*				*			*		*			*		*		*		*				*
<i>Conyza bonariensis</i> *	*				*								*		*		*		*				
<i>Cotula coronopifolia</i> *	*		*		*	*	*	*	*	*	*			*		*		*				*	
<i>Cynogeton procera</i>																						*	
<i>Cyperus gymnocaulos</i>	*	*		*		*		*	*		*												
<i>Disphyma crassifolium</i>					*	*	*	*	*	*	*		*	*		*	*	*		*	*	*	*
<i>Distichlis distichophylla</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Duma florulenta</i>						*	*			*				*	*	*	*	*		*			
<i>Einadia nutans</i>		*																					
<i>Eleocharis acuta</i>				*		*		*		*				*						*	*	*	*
<i>Epilobium paladiflorum</i>																							*
<i>Enchylaena tomentosa</i>	*	*	*	*	*	*		*	*			*	*		*			*					
<i>Eragrostis curvula</i> **	*		*		*		*		*					*				*			*		
<i>Eragrostis</i> sp.	*		*		*		*																
<i>Erodium cicutarium</i> *								*							*								
<i>Ficinia nodosa</i>	*		*									*	*				*						
<i>Frankenia pauciflora</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Glyceria australis</i>					*		*																
<i>Heliotropium europaeum</i> *				*		*																	
<i>Hordeum vulgare</i> *	*		*		*	*	*	*	*		*						*	*		*	*		
<i>Hypochoeris glabra</i> *						*														*	*		
<i>Isolepis producta</i>									*			*		*		*			*	*			*
<i>Juncus acutus</i> **													*										
<i>Juncus kraussii</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Juncus subsecundus</i>								*															*
<i>Lachnagrostis filiformis</i>	*		*		*			*	*	*	*				*			*	*			*	
<i>Lactuca saligna</i> *	*		*				*																
<i>Lactuca serriola</i> *	*		*		*	*	*	*	*		*						*	*		*			*
<i>Lagurus ovatus</i> *	*		*		*																		
<i>Lamprothamnium macropogon</i>	*		*		*	*	*		*		*			*		*		*		*	*	*	*
<i>Lemna</i> spp.						*			*							*		*		*	*	*	
<i>Limosella australis</i>																						*	
<i>Lobelia anceps</i>					*																		
<i>Lolium</i> spp.*	*		*	*	*	*	*	*	*		*		*	*		*		*		*	*	*	*
<i>Ludwigia peploides</i> ssp. <i>montevidensis</i>													*		*				*				

Taxon	Survey Date																						
	Spring 2008	Autumn 2009	Spring 2009	Autumn 2010	Spring 2010	Autumn 2011	Spring 2011	Autumn 2012	Spring 2012	Autumn 2013	Spring 2013	Autumn 2014	Autumn 2015	Spring 2015	Autumn 2016	Spring 2016	Autumn 2017	Spring 2017	Autumn 2018	Spring 2018	Spring 2019	Spring 2020	Autumn 2021
<i>Lycium ferocissimum</i> ***	*		*	*	*	*		*												*			
<i>Lythrum hyssopifolia</i>																						*	
<i>Lythrum salicaria</i>																		*					
<i>Malva parviflora</i> *						*																	
<i>Marrubium vulgare</i> **					*																		
<i>Medicago</i> spp.*	*		*		*	*	*	*		*					*								
<i>Melaleuca halmaturorum</i>							*			*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Melilotus albus</i> *									*														
<i>Melilotus indicus</i> *	*		*		*		*				*												
<i>Myoporum insulare</i>													*					*	*	*		*	*
<i>Myriophyllum salsugineum</i>					*									*			*						
<i>Myriophyllum verrucosum</i>													*										
<i>Paspalum distichum</i> *		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Phragmites australis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Phyla canescens</i> *	*																						
<i>Picris angustifolia</i> ssp. <i>angustifolia</i>					*	*																	
<i>Plantago coronopus</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Polygonum aviculare</i> *	*	*																					
<i>Polypogon monspeliensis</i> *	*		*		*		*		*		*			*		*		*	*	*	*	*	*
<i>Potamogeton pectinatus</i>						*	*	*	*	*													
<i>Pseudognaphalium luteoalbum</i>					*		*																
<i>Puccinellia</i> sp.*			*				*																
<i>Ranunculus trichophyllus</i>					*		*		*	*	*	*									*	*	*
<i>Reichardia tingitana</i> *	*		*			*	*		*														
<i>Rorippa nasturtium-aquaticum</i> *																				*			
<i>Rorippa palustris</i> *			*																				
<i>Rumex bidens</i>	*		*			*					*	*		*		*		*	*	*	*	*	*
<i>Ruppia polycarpa</i>					*	*	*		*													*	
<i>Ruppia tuberosa</i>	*		*		*	*		*	*		*			*		*		*		*	*	*	*
<i>Salsola australis</i>				*		*								*	*			*					
<i>Samolus repens</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Sarcocornia quinqueflora</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Schoenoplectus pungens</i>		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Schoenoplectus tabernaemontani</i>					*																		
<i>Senecio pterophorus</i> *	*		*	*						*								*	*	*			*
<i>Silybum marianum</i> **					*																		
<i>Sonchus asper</i> *															*						*	*	
<i>Sonchus oleraceus</i> *	*		*	*	*	*	*	*	*	*	*	*	*	*		*				*	*	*	
<i>Spergularia brevifolia</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Suaeda australis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Aster subulatus</i> *	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Thyridia repens</i>	*		*		*	*	*	*	*	*	*	*	*	*			*		*			*	
<i>Trifolium</i> spp.*	*		*		*	*	*		*		*	*		*		*		*		*	*	*	*
<i>Triglochin striata</i>	*				*	*	*	*	*	*	*		*	*		*	*	*	*	*	*	*	*
<i>Typha domingensis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Wilsonia rotundifolia</i>	*				*	*	*	*	*			*	*		*	*			*		*		*