

Algal bloom impact on key fish stocks in South Australia's gulfs

Updated: 11 Sept. 2026

Source: SARDI Aquatic and Livestock Sciences

Overview

Since March 2025, a large algal bloom caused by *Karenia* species has affected the waters of Gulf St Vincent, Spencer Gulf, Kangaroo Island and nearby areas. The bloom has resulted in widespread marine life deaths, impacting fish, shellfish and other sea life.

As part of the extensive SARDI Algal Bloom Response, fisheries and fish stocks are being assessed to understand the impact of the algal bloom on commercial fish species. This Fact Sheet outlines how the algal bloom is affecting key quota-managed species — King George Whiting, Southern Calamari, Southern Garfish, Western King Prawn, Blue Swimmer Crab, Southern Rock Lobster and Abalone — in Gulf St Vincent/Kangaroo Island (GSV/KI) and Spencer Gulf (SG), where the bloom has been most widespread (Figure 1). It also provides a summary report on 17 non-quota species in both gulfs.

Methods

This assessment uses both commercial fishing and independent survey data. Specifically, the assessment combines:

1. Analysis of commercial fishing data.
2. Scientific (fishery-independent) surveys.
3. Targeted fishing surveys with commercial operators.
4. Bycatch and species sampling.
5. Routine fishery stock assessment surveys for prawns, crabs, lobsters and abalone.

For all species, the assessment relies heavily on spatial and temporal analyses of commercial logbook data across all fishing sectors. To quantify fishery impacts, monthly catch and catch rate (expressed as catch per unit effort; CPUE) during the algal bloom period (March 2025 onwards), were compared against the 3-year monthly average (baseline) from February 2022 – February 2025 for each species. This allows quantification of a “before” and “after” algal bloom impact.

The GSV/KI zone was closed to commercial fishing on 01 November 2025 while SG was closed to Southern Calamari commercial fishing on 01 May 2026. Consequently, all catch data from these dates onwards for these fisheries are based on targeted fishery-dependent surveys. Indices are categorised as having changed by <50% of 3-year monthly average; 50–80% of 3-year monthly average, or >80% of 3-year monthly average. Where available, logbook data were supported by fishery-independent surveys, targeted fishing, bycatch sampling and stock assessment surveys.

Results

King George Whiting

Data sources: Commercial logbook, fishery-dependent research and post-settlement surveys.

Monthly catches and catch rates from GSV/KI during the algal bloom period from March–August 2025 remained broadly within recent ranges. Total catch in September and October 2025 declined sharply, with catch rates from targeted fishing also decreasing substantially between November 2025 and August 2026 (Figure 2). Both metrics to August 2026 were generally >80% below baseline levels at both zonal (Table 1; Table 2) and regional (Table 3; Table 4) scales. SG catch and catch rate have remained consistent with historical averages (Figure 2; Table 1; Table 2).

Post settlement surveys of King George Whiting were undertaken across GSV and SG in 2025 and 2026. Across the 18 locations sampled, post-settled King George Whiting densities were at their lowest levels on record, with declines of 37–97% since the most recent survey in 2017. The greatest reductions occurred in Lower Eyre Peninsula, Northern GSV and Southern GSV while Southern SG was the only region to show an increase.

Southern Calamari

Data sources: Commercial logbook, fishery-dependent research and fishery-independent dive surveys.

Commercial logbook and targeted fishing data from GSV/KI show marked declines, with catch and CPUE from July to October 2025 negligible relative to the 3-year average (Figure 3). Since the closure in November 2025, targeted CPUE has remained >80% below baseline to August 2026 at both zonal (Table 1; Table 2) and regional spatial scales (Table 3; Table 4). In SG, catch and CPUE remained broadly consistent with previous years until September 2025 (Figure 3). From October 2025 to April 2026, catch declined to >80% below baseline levels, while CPUE progressively decreased to >80% below baseline by June 2026 (Table 1; Table 2). Although CPUE remains low, there was some preliminary evidence of improvement in July and August, with CPUE increasing to 53% below baseline levels in August 2026. Overall declines are widespread across both northern and southern SG regions (Table 5; Table 6).

Seven fishery-independent diver surveys of Southern Calamari egg masses have been completed to date: six in GSV and one in SG. In GSV, no eggs were observed during October to December 2025 and June 2026 surveys, compared with the ~7000–10,500 eggs recorded at various sites in 2005–06. In SG, first-time surveys across four exploratory sites recorded four small egg masses (154 strands) in clear water with normal fauna, suggesting low Southern Calamari abundance rather than localised effects of the algal bloom.

Southern Garfish

Data sources: [Commercial logbook, fishery-dependent research and fishery-independent surveys.](#)

For Southern Garfish, logbook data from GSV/KI indicate strong negative impacts of the algal bloom, with catches and CPUE declining by >80% from May to October 2025 (Figure 4). Following the closure in November 2025, targeted CPUE generally remained >80% below baseline levels at both the zonal (Table 1, Table 2) and regional (Table 3; Table 4) scales. The exception was July and August 2026, when CPUE was 50–80% below historical averages. For SG, however, catches and CPUE were largely consistent with historical baseline data (Figure 4; Table 1; Table 2).

Fishery independent abundance surveys of Southern Garfish abundance were conducted in November 2025 at Glenelg and North Haven and February 2026 at Middle Beach and Edithburgh. Compared to similar surveys undertaken in 2016 and 2028, total catch declined by 61% at Glenelg, 88% at North Haven, 84% at Middle Beach and 25% at Edithburgh. Most of the observed Southern Garfish were small and few other species were observed. These surveys were the first in the series to be performed monthly during the new moon, and data must be considered with caution.

Western King Prawn

Data sources: [Commercial logbook and fishery-independent surveys](#)

Monthly logbook catch and catch rate data for Western King Prawn in SG indicated no negative impact from the algal bloom on Western King Prawn biomass (noting that fishing is not undertaken in SG from July to September; Figure 5). However, in GSV, both catch and CPUE were >80% below baseline from October to December 2025, except for CPUE in November.

In GSV, five algal bloom fishery-independent surveys were carried out from August to December 2025. CPUE showed a consistent decline of 72% during this period. Importantly, no elasmobranchs or Southern Calamari were observed as bycatch during the GSV surveys.

Eight fishery-independent surveys were carried out in SG between August 2025 and June 2026. There was no evidence to suggest a negative impact from the algal bloom on Western King Prawn in SG, with CPUE above the previous 10-year annual mean. Bycatch/byproduct that included elasmobranchs were recorded but Southern Calamari catches declined progressively from October 2025 onwards.

Blue Swimmer Crab

Data sources: [Commercial logbook and fishery-independent surveys](#)

Monthly logbook catch and catch rate data for GSV show clear negative impacts of the algal bloom on Blue Swimmer Crab. Catches and CPUE have declined rapidly since March 2025. From August to October 2025, both were >80% below baseline. Catch and catch rates have since risen above baseline averages

OFFICIAL

as of August 2026, following a fishery closure that lasted from 1 November 2025 until May 2026. There was no evidence of negative impacts of the algal bloom on Blue Swimmer Crabs in SG.

Three fishery-independent surveys for the algal bloom response were undertaken in GSV from 31 August to 3 September 2025, 25 September to 1 October 2025 and 27 January–5 February 2026, sampling consistent sites along Yorke Peninsula and the Metropolitan and Northern coastlines. Across the three trips, total catch from 1,243 pots was 2,847 kg (8,936 legal-size crabs), though catches varied widely by time and location. Trip 1 catch rates were very low (0.1–0.2 kg or 0.5–0.6 crabs/pot), while Trip 2 had higher catch rates (0.7–0.9 kg or 2.7–3.3 crabs/pot). However, in Trip 3 (27 January–5 February 2026), catch rates were 6.4 kg/pot, well above the long-term January average (3.1 kg/pot). Research-pot undersize catch rates during Trips 2 and 3 (0.2–1.2 kg/pot) were also within the expected March–April survey range (0.1–3.5 kg pot).

Annual stock assessment fishery-independent surveys were undertaken in both SG and GSV/KI during March 2026. Surveys were conducted in SG from 2–7 March and in GSV/KI from 17–22 March. In GSV/KI, research pot CPUE for legal-size crabs was 5.6 ± 0.4 (SE) kg potlift⁻¹, representing a 75% increase relative to the three-year average of 3.2 ± 0.73 (SE) kg potlift⁻¹. Commercial pot CPUE for legal-size crabs was 5.8 ± 0.1 (SE) kg potlift⁻¹, a 61% increase above the three-year average of 3.6 ± 0.9 (SE) kg potlift⁻¹.

In SG, research pot CPUE for legal-size crabs was 8.1 ± 0.4 (SE) kg potlift⁻¹, reflecting a 40% increase relative to the three-year average of 5.8 ± 1.1 (SE) kg potlift⁻¹. Commercial pot CPUE for legal-size crabs was 11.7 ± 0.2 (SE) kg potlift⁻¹, representing a 109% increase above the three-year average of 5.6 ± 0.8 (SE) kg potlift⁻¹. The March 2026 surveys indicate that the biomass in GSV/KI and SG are above the target reference level, consistent with a sustainable stock status classification under the harvest strategy.

Southern Rock Lobster

[Data sources: Commercial logbook and fishery-independent surveys](#)

Based on monthly logbook catch and catch rates in both the Northern and Southern Zones, there was no evidence of a negative impact from the algal bloom on Southern Rock Lobster (Figure 6). The latest fishery-independent survey was undertaken in the Southern Zone in January/February 2026. In the 2025/26 season, the catch rates of legal and undersized Southern Rock Lobsters were 79% and 76% higher than 2024/25 estimates, respectively.

Abalone

[Data sources: fishery-independent surveys](#)

Fishery-independent dive surveys were undertaken to the West (Corny Point, Hardwicke Bay, Port Victoria and Tiparra Reef) and East (Stansbury and Edithburgh) of Yorke Peninsula between August and November 2025.

OFFICIAL

The surveys at Corny Point and Tiparra Reef had the highest number of legal sized Greenlip Abalone since surveys began in 2015 and 1990, respectively. For Hardwicke Bay, counts of both legal- and sub-legal-sized Greenlip Abalone were higher in 2025 than the most recent survey in 2023, and similar to counts from surveys in 2019 and 2021. At Port Victoria, counts of Greenlip Abalone were similar to those recorded in the previous 2023 survey. From the 11 transects completed at Tiparra Reef, counts were the highest on record. Consequently, there was no evidence of algal bloom impacts on Greenlip Abalone in SG.

In contrast, at all sites surveyed by divers at Stansbury and Edithburgh, there was evidence that the algal bloom has impacted abalone in GSV. At Stansbury, divers observed unwell, dying or dead abalone of all three species (*Haliotis laevis*, *H. scalaris*, *H. cyclobates*). Samples of these three species (n = 10) were collected and tested for Abalone Viral Ganglioneuritis (AVG), with all tests returning a negative result. Some live abalone were observed at this location, and abundance varied among sites (Mean 1.075 Greenlip Abalone/min). There are no previous surveys for temporal comparison but estimates from all four dive sites suggested 70% to 90% of Greenlip Abalone were unwell or dead, and almost all other abalone species were dead. At Edithburgh, there was also evidence of impact from the algal bloom, but these appeared to be less severe than those observed at Stansbury.

Additional surveys in mid-May 2026 reinforced earlier observations, revealing substantial declines relative to 2025 estimates, including 100% in Hardwicke Bay, 68% in Port Victoria, 49% at Corny Point, and 20% at Tiparra Reef. Together, these results indicate a clear and significant impact of the algal bloom on Greenlip Abalone populations in GSV, particularly in southern sites.

Non-quota species

Data sources: [Commercial logbook](#).

A total of 17 species or species groups from the Marine Scalefish Fishery were assessed. The low monthly catches for many species limited the ability to detect potential impacts of the algal bloom; consequently, the assessment presented was based on qualitative interpretation rather than quantitative analysis.

Among the 17 species assessed in 2026, eleven (Yellowfin Whiting, Western Australian Salmon, Australian Herring, Whaler Sharks, Snook, Yellow-eye Mullet, Bluethroat Wrasse, Leatherjackets, Cuttlefish, Gummy Shark, School Shark) were considered to have experienced negative impacts from the algal bloom in Gulf St Vincent/Kangaroo Island, three (Silver Trevally, Rays and Skates, Black Bream) showed no apparent impact, and data were insufficient for three (Sand Crab, Mulloway, Ocean Leatherjacket) species. In Spencer Gulf, six (Cuttlefish, Western Australian Salmon, Whaler Sharks, Sand Crab, Bluethroat Wrasse, Gummy Shark,) species were similarly assessed as negatively affected, nine (Australian Herring, Yellow-eye Mullet, Leatherjackets, Yellowfin Whiting, Snook, Ocean Leatherjacket, Rays and Skates, Silver Trevally and School Shark) showed no impact, and two (Mulloway, Black Bream) lacked sufficient data.

OFFICIAL

Main findings

Gulf St Vincent/Kangaroo Island

- **King George Whiting:** Evidence of declines with widespread indication of post-settlement failure.
- **Southern Calamari and Southern Garfish:** Severely affected with additional evidence of egg mass/juvenile declines.
- **Western King Prawn:** Strong evidence from scientific surveys of low catches and very little bycatch.
- **Blue Swimmer Crab:** Catches have been impacted but estimates from recent targeted surveys and commercial logbooks indicate that abundances have improved.
- **Abalone:** High mortality observed in parts of the gulf.

Spencer Gulf

- **King George Whiting and Southern Garfish:** Generally stable
- **Southern Calamari:** Evidence of catch declines across broad spatial scales but some preliminary evidence of improved catch rates in July and August 2026.
- **Western King Prawn and Blue Swimmer Crab:** Catches remain within normal ranges.
- **Southern Rock Lobster:** Found mostly outside the affected areas – no sign of impact.

What this means

- The algal bloom has had the **strongest negative effects in Gulf St Vincent**, where several key fish species have declined sharply.
- **Spencer Gulf fisheries appear to be less affected**, except for Southern Calamari.
- Results will continue to be updated as more fishing and survey data become available.

Next steps

- SARDI continues to collect and analyse data to track impacts and provide further advice to PIRSA and industry.
- Updates will be shared as new information becomes available.

Figures




Figure 1. Management zones and marine fishing areas (blocks) in the Marine Scalefish Fishery. Abbreviations for management zones: WC, West Coast; NSG, Northern Spencer Gulf; SSG, Southern Spencer Gulf; NGSV, Northern Gulf St Vincent; SGSV, Southern Gulf St Vincent; SE, South East.

King George Whiting




Figure 2. Commercial Fishery statistics for King George Whiting in the Gulf St Vincent/Kangaroo Island and Spencer Gulf fishing zones. A) Total catch (t) and B) Targeted CPUE (kg/fisher-day) during the algal bloom period (circle: March 2025 to February 2026; square: March 2026 onwards) compared to the 3-year monthly average (February 2022 – February 2025; grey bars). Error bars indicate standard error. Green indicates change in catch/CPUE <50% of 3-year monthly average; orange indicates change in catch/CPUE of 50-80% of 3-year monthly average, red indicates change in catch/CPUE >80% of 3-year monthly average. Dashed line indicates closure of commercial fishery in Gulf St Vincent/Kangaroo Island fishing zone from 01 November 2025. Data from November 2025 onwards (GSV/KI) from targeted fishing program only.

Southern Calamari




Figure 3. Commercial Fishery statistics for Southern Calamari in the Gulf St Vincent/Kangaroo Island and Spencer Gulf fishing zones. A) Total catch (t) and B) Targeted CPUE (kg/fisher-day) during the algal bloom period (circle: March 2025 to February 2026; square: March 2026 onwards) compared to the 3-year monthly average (February 2022 – February 2025; grey bars). Error bars indicate standard error. Green indicates change in catch/CPUE <50% of 3-year monthly average; orange indicates change in catch/CPUE of 50-80% of 3-year monthly average, red indicates change in catch/CPUE >80% of 3-year monthly average. Dashed line indicates closure of commercial fishery in Gulf St Vincent/Kangaroo Island fishing zone from 01 November 2025 and Spencer Gulf from 01 May 2026. Data from November 2025 (GSV/KI) and May 2026 (SG) onwards from targeted fishing program only.

Southern Garfish




Figure 4. Commercial Fishery statistics for Southern Garfish in the Gulf St Vincent/Kangaroo Island and Spencer Gulf fishing zones. A) Total catch (t) and B) Targeted CPUE (kg/fisher-day) during the algal bloom period (circle: March 2025 to February 2026; square: March 2026 onwards) compared to the 3-year monthly average (February 2022 – February 2025; grey bars). Error bars indicate standard error. Green indicates change in catch/CPUE <50% of 3-year monthly average; orange indicates change in catch/CPUE of 50-80% of 3-year monthly average, red indicates change in catch/CPUE >80% of 3-year monthly average. Dashed line indicates closure of commercial fishery in Gulf St Vincent/Kangaroo Island fishing zone from 01 November 2025. Data from November 2025 onwards (GSV/KI) from targeted fishing program only. Grey bars indicate months where data were removed due to confidentiality (< 5 licences).

Western King Prawn




Figure 5. Commercial Fishery statistics for Western King Prawn in the Gulf St Vincent/Kangaroo Island and Spencer Gulf fishing zones. A) Total catch (t) and B) Targeted CPUE (kg/rawl-hour) during the algal bloom period (circle: March 2025 to February 2026; square: March 2026 onwards) compared to the 3-year monthly average (February 2022 – February 2025; grey bars). Error bars indicate standard error. Green indicates change in catch/CPUE <50% of 3-year monthly average; orange indicates change in catch/CPUE of 50-80% of 3-year monthly average, red indicates change in catch/CPUE >80% of 3-year monthly average. Grey bars indicate months where data were removed due to confidentiality (< 5 licences).

Southern Rock Lobster




Figure 6. Commercial Fishery statistics for Southern Rock Lobster in the Southern and Northern fishing zones. A) Total catch (t) and B) Targeted CPUE (kg/rawl-hour) during the algal bloom period (circle: March 2025 to February 2026; square: March 2026 onwards) compared to the 3-year monthly average (February 2022 – February 2025; grey bars). Error bars indicate standard error. Green indicates change in catch/CPUE <50% of 3-year monthly average; orange indicates change in catch/CPUE of 50-80% of 3-year monthly average, red indicates change in catch/CPUE >80% of 3-year monthly average. Low Southern Rock Lobster catches in March–April (Southern Zone) and July–August (Northern Zone) are a consequence of the TACC being close to fully harvested.

OFFICIAL

Table 1. Commercial fishery statistics for King George Whiting, Southern Calamari, Southern Garfish, Blue Crab and Western King Prawn in Gulf St Vincent/Kangaroo Island and Spencer Gulf fishing zones. Data is total catch (t) during the algal bloom period (March 2025 onwards) compared to the 3-year monthly average (March 2022 – February 2025). **Green** reflects change in catch by <50% of baseline (3-year average). **Orange** reflects a change in catch of 50-80% of baseline (3-year average). **Red** reflects change in catch of >80% of baseline (3-year average).

TOTAL CATCH																			
Gulf St Vincent/Kangaroo Island		2025										2026							
	King George Whiting - GSVKI	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	Total catch (t) (3 year average)	2.02	1.42	2.10	2.61	2.70	2.94	2.82	2.68	2.06	1.82	1.99	2.02	2.02	1.42	2.10	2.61	2.70	2.94
	Total catch (t) (Algal bloom period)	1.560	1.410	1.390	3.390	2.970	2.790	1.530	1.630	0.020	0.030	conf	0.060	0.110	conf	0.040	0.020	0.070	0.050
	Percentage of baseline	-22.86	-1.03	-33.92	30.04	9.81	-5.25	-45.77	-39.15	-99.03	-98.35		-97.03	-94.56		-98.10	-99.23	-97.41	-98.30
	Southern Calamari - GSVKI	6.86	12.10	13.78	7.44	6.63	9.95	11.25	13.16	15.81	9.66	9.32	7.52	6.86	12.10	13.78	7.44	6.63	9.95
	Total catch (t) (3 year average)	8.25	21.53	21.83	4.90	0.02	0.10	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.01	0.02	0.02	0.04	0.05
	Total catch (t) (Algal bloom period)	20.20	77.88	58.47	-34.11	-99.70	-98.99	-100.00	-99.85	-100.00	-100.00	-100.00	-100.00	-100.00	-99.92	-99.85	-99.73	-99.40	-99.50
	Percentage of baseline																		
	Southern Garfish - GSVKI	7.62	8.48	7.62	4.66	6.93	5.64	3.25	2.23	4.06	4.50	6.57	9.20	7.62	8.48	7.62	4.66	6.93	5.64
	Total catch (t) (3 year average)	7.19	7.99	3.98	2.46	2.71	0.78	conf	conf	0.00	conf	conf	0.02	0.05	conf	conf	0.15	conf	conf
	Total catch (t) (Algal bloom period)	-5.60	-5.77	-47.76	-47.16	-60.90	-86.16			-100.00			-99.78	-99.34			-96.78		
	Percentage of baseline																		
	Blue Crab - GSV	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf
	Total catch (t) (3 year average)	conf	conf	conf	conf	conf	conf	conf	conf	NA	NA	NA	NA	NA	NA	conf	conf	conf	conf
	Total catch (t) (Algal bloom period)																		
	Percentage of baseline																		
	Prawn - GSV	19.01	25.60	28.33	24.69	3.63	conf	NA	conf	22.33	18.05	NA	NA	19.01	25.60	28.33	24.69	3.63	conf
	Total catch (t) (3 year average)	conf	conf	conf	30.82	conf	11.17	conf	conf	conf	conf	NA	NA	NA	NA	NA	NA	NA	NA
	Total catch (t) (Algal bloom period)				24.83														
	Percentage of baseline																		
Spencer Gulf		2025										2026							
	King George Whiting - SG	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	Total catch (t) (3 year average)	4.01	2.54	3.83	9.57	14.12	11.26	7.17	4.96	5.25	4.09	4.93	5.28	4.01	2.54	3.83	9.57	14.12	11.26
	Total catch (t) (Algal bloom period)	2.61	2.28	4.35	11.82	14.45	12.59	8.58	5.59	4.87	5.98	3.62	4.15	5.25	1.90	4.14	10.50	13.59	7.11
	Percentage of baseline	-34.96	-10.24	13.60	23.54	2.34	11.78	19.61	12.64	-7.29	46.08	-26.64	-21.41	30.84	-25.20	8.11	9.74	-3.75	-36.88
	Southern Calamari - SG	7.77	13.05	16.50	10.00	11.17	10.77	16.15	19.63	15.81	14.81	13.02	9.12	7.77	13.05	16.50	10.00	11.17	10.77
	Total catch (t) (3 year average)	6.97	18.48	18.33	17.42	12.24	14.47	9.01	8.57	8.64	3.81	1.64	0.47	0.47	1.78	0.05	0.01	0.09	0.64
	Total catch (t) (Algal bloom period)	-10.30	41.63	11.09	74.25	9.55	34.35	-44.21	-56.33	-45.34	-74.28	-87.40	-94.85	-93.95	-86.36	-99.70	-99.90	-99.19	-94.06
	Percentage of baseline																		
	Southern Garfish - SG	6.51	5.96	8.88	5.74	5.06	7.72	3.92	4.12	5.76	4.78	10.38	10.03	6.51	5.96	8.88	5.74	5.06	7.72
	Total catch (t) (3 year average)	7.57	6.06	6.71	5.24	3.78	6.53	12.16	4.57	6.35	6.77	9.90	10.64	14.25	9.33	9.50	6.58	8.57	5.19
	Total catch (t) (Algal bloom period)	16.28	1.71	-24.43	-8.69	-25.30	-15.40	209.95	10.94	10.28	41.49	-4.59	6.03	118.88	56.59	7.00	14.66	69.37	-32.76
	Percentage of baseline																		
	Blue Crab - SG	conf	conf	conf	conf	conf	conf	conf	conf	conf	17.93	conf	39.50	conf	conf	conf	conf	conf	conf
	Total catch (t) (3 year average)	36.70	conf	conf	conf	conf	63.59	70.81	52.11	34.06	25.31	conf	41.28	46.64	conf	conf	conf	conf	conf
	Total catch (t) (Algal bloom period)										41.18		4.50						
	Percentage of baseline																		
	Prawn - SG	221.95	495.82	337.98	99.72	NA	NA	NA	72.61	215.80	186.41	NA	NA	221.95	495.82	337.98	99.72	NA	NA
	Total catch (t) (3 year average)	563.96	553.72	385.74	66.17	NA	NA	NA	262.25	225.52	NA	NA	NA	200.01	643.79	483.52	235.77	NA	NA
	Total catch (t) (Algal bloom period)	154.09	11.68	14.13	-33.64				261.20	4.50				-9.88	29.84	43.06	136.44		
	Percentage of baseline																		

OFFICIAL

Table 2. Commercial fishery statistics for King George Whiting, Southern Calamari, Southern Garfish, Blue Crab and Western King Prawn in Gulf St Vincent/Kangaroo Island and Spencer Gulf fishing zones. Data is targeted CPUE (kg/fisher-day) during the algal bloom period (March 2025 onwards) compared to the 3-year monthly average (February 2022 – February 2025). **Green** reflects change in CPUE by <50% of baseline (3-year average). **Orange** reflects a change in CPUE of 50-80% of baseline (3-year average). **Red** reflects change in CPUE of >80% of baseline (3-year average).

		Target CPUE																		
Gulf St Vincent/Kangaroo Island		2025										2026								
	King George Whiting - GSVKI	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	
	CPUE (kg/fisher-days) (3 year average)	15.74	11.44	14.88	20.21	20.05	20.72	22.46	22.93	19.64	19.83	17.75	22.28	15.74	11.44	14.88	20.21	20.05	20.72	
	CPUE (kg/fisher-days) (Algal bloom period)	27.01	18.14	14.76	21.00	30.03	25.68	23.48	15.52	2.00	2.62	conf	2.55	3.44	conf	1.52	1.21	7.01	2.50	
	Percentage of baseline	71.60	58.58	-0.83	3.92	49.76	23.96	4.53	-32.33	-89.82	-86.78		-88.56	-78.15		-89.79	-94.01	-65.04	-87.93	
	Southern Calamari - GSVKI	CPUE (kg/fisher-days) (3 year average)	22.87	27.25	24.87	23.28	20.25	27.33	32.74	40.44	47.33	39.76	39.09	32.72	22.87	27.25	24.87	23.28	20.25	27.33
	CPUE (kg/fisher-days) (Algal bloom period)	31.34	49.65	50.85	34.64	0.03	0.00	0.13	0.18	0.00	0.00	0.00	0.06	0.16	0.73	2.40	1.06	2.10	4.73	
	Percentage of baseline	37.04	82.21	104.46	48.81	-99.85	-100.00	-99.60	-99.55	-100.00	-100.00	-100.00	-99.82	-99.30	-97.32	-90.35	-95.45	-89.63	-82.70	
	Southern Garfish - GSVKI	CPUE (kg/fisher-days) (3 year average)	55.14	76.38	81.07	97.04	113.30	81.14	60.44	36.95	45.85	56.85	71.97	68.81	55.14	76.38	81.07	97.04	113.30	81.14
	CPUE (kg/fisher-days) (Algal bloom period)	75.61	106.25	99.57	conf	55.69	conf	conf	conf	0.25	conf	conf	1.93	2.79	conf	conf	7.88	conf	conf	
	Percentage of baseline	37.13	39.11	22.82		-50.85				-99.45			-97.20	-94.94			-91.88			
	Blue Crab - GSVKI	CPUE (kg/pot) (3 year average)	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	
	CPUE (kg/pot) (Algal bloom period)	conf	conf	conf	conf	conf	conf	conf	conf	NA	NA	NA	NA	NA	NA	conf	conf	conf	conf	
	Percentage of baseline																			
	Prawn - GSV	CPUE (kg/haul-hour) (3 year average)	41.30	43.64	41.21	39.65	24.59	conf	NA	conf	75.10	71.90	NA	NA	41.30	43.64	41.21	39.65	24.59	conf
	CPUE (kg/haul-hour) (Algal bloom period)	conf	conf	conf	66.77	conf	49.87	conf	conf	conf	conf	conf	NA	NA	NA	NA	NA	NA	NA	
	Percentage of baseline				68.41															
Spencer Gulf		2025										2026								
	King George Whiting - SG	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	
	CPUE (kg/fisher-days) (3 year average)	17.86	13.84	18.25	29.18	29.22	26.93	23.45	24.28	26.77	23.39	21.57	23.23	17.86	13.84	18.25	29.18	29.22	26.93	
	CPUE (kg/fisher-days) (Algal bloom period)	18.08	11.89	18.26	30.13	37.91	31.31	25.48	18.13	23.75	23.10	21.31	20.48	20.33	9.24	12.59	24.18	22.27	20.39	
	Percentage of baseline	1.20	-14.08	0.04	3.26	29.75	16.28	8.66	-25.32	-11.29	-1.25	-1.21	-11.85	13.80	-33.23	-31.02	-17.13	-23.78	-24.28	
	Southern Calamari - SG	CPUE (kg/fisher-days) (3 year average)	23.33	27.14	30.56	32.33	29.38	26.55	36.13	43.08	46.16	46.10	36.76	29.63	23.33	27.14	30.56	32.33	29.38	26.55
	CPUE (kg/fisher-days) (Algal bloom period)	25.33	35.05	34.67	47.93	40.27	35.43	30.12	24.80	31.74	20.00	13.29	7.10	3.90	10.63	8.50	0.38	10.16	12.36	
	Percentage of baseline	8.56	29.13	13.45	48.24	37.05	33.45	-16.63	-42.43	-31.23	-56.62	-63.85	-76.04	-83.29	-60.84	-72.19	-98.82	-65.42	-53.44	
	Southern Garfish - SG	CPUE (kg/fisher-days) (3 year average)	52.81	50.39	64.20	60.04	61.39	74.81	52.97	43.64	54.57	46.27	66.72	64.26	52.81	50.39	64.20	60.04	61.39	74.81
	CPUE (kg/fisher-days) (Algal bloom period)	63.35	58.75	59.85	83.32	43.55	60.87	150.74	46.15	42.44	40.93	54.77	52.89	48.91	47.37	59.29	57.37	96.06	49.79	
	Percentage of baseline	19.97	16.58	-6.77	38.77	-29.06	-18.63	184.60	5.75	-22.23	-11.53	-17.91	-17.69	-7.38	-6.00	-7.64	-4.45	56.48	-33.44	
	Blue Crab - SG	CPUE (kg/pot) (3 year average)	conf	conf	conf	conf	conf	conf	conf	conf	2.19	conf	4.53	conf	conf	conf	conf	conf	conf	
	CPUE (kg/pot) (Algal bloom period)	3.13	conf	conf	conf	conf	4.11	4.38	3.43	2.60	2.33	conf	3.97	4.24	conf	conf	conf	conf	conf	
	Percentage of baseline										6.34		-12.34							
	Prawn - SG	CPUE (kg/haul-hour) (3 year average)	78.38	119.14	84.67	59.29	NA	NA	NA	56.74	76.98	83.11	NA	NA	78.38	119.14	84.67	59.29	NA	NA
	CPUE (kg/haul-hour) (Algal bloom period)	148.14	135.84	110.91	55.94	NA	NA	NA	121.41	73.96	NA	NA	NA	98.09	182.38	131.98	85.85	NA	NA	
	Percentage of baseline	89.00	14.02	30.99	-5.66				113.96	-3.92				25.14	53.08	55.87	44.79			

OFFICIAL

Table 3. Regional commercial fishery statistics for King George Whiting, Southern Calamari and Southern Garfish in the Gulf St Vincent/Kangaroo Island fishing zone. Data is total catch (t) during the algal bloom period (March 2025 onwards) compared to the 3-year monthly average (February 2022 – February 2025). **Green** reflects change in catch by <50% of baseline (3-year average). **Orange** reflects a change in catch of 50-80% of baseline (3-year average). **Red** reflects change in catch of >80% of baseline (3-year average).

TOTAL CATCH																			
Northern Gulf St Vincent (34, 35, 36)		2025										2026							
	King George Whiting - NGSV	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	Total catch (t) (3 year average)	0.51	0.32	0.53	0.64	0.76	0.74	0.75	0.67	0.65	0.43	0.46	0.42	0.51	0.32	0.53	0.64	0.76	0.74
	Total catch (t) (Algal bloom period)	0.20	0.22	1.33	3.19	2.30	1.98	conf	0.12	0.01	0.01	0.02	0.01	0.01	0.01	0.00	0.02	0.01	0.00
	Percentage of baseline	-61.01	-31.32	150.35	401.15	202.90	168.47		-82.02	-98.45	-97.67	-95.64	-97.64	-98.05	-96.88	-100.00	-96.86	-98.68	-100.00
	Southern Calamari - NGSV																		
	Total catch (t) (3 year average)	4.95	10.79	11.67	6.43	5.25	7.57	5.37	2.44	1.98	0.91	0.85	1.30	4.95	10.79	11.67	6.43	5.25	7.57
	Total catch (t) (Algal bloom period)	4.84	19.61	21.72	4.90	0.02	0.10	NA	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.01	0.01
	Percentage of baseline	-2.31	81.68	86.15	-23.76	-99.62	-98.68		-100.00	-100.00	-100.00	-100.00	-100.00	-100.00	-99.91	-99.91	-99.84	-99.81	-99.87
	Southern Garfish - NGSV																		
	Total catch (t) (3 year average)	7.24	8.07	7.36	4.60	6.88	5.53	3.10	2.00	3.80	4.27	6.50	8.98	7.24	8.07	7.36	4.60	6.88	5.53
	Total catch (t) (Algal bloom period)	7.02	7.97	3.98	2.46	2.71	0.78	conf	conf	0.00	0.00	0.00	0.01	0.04	0.04	0.01	0.13	NA	0.07
	Percentage of baseline	-3.00	-1.28	-45.91	-46.58	-60.61	-85.89			-100.00	-100.00	-100.00	-99.89	-99.45	-99.50	-99.86	-97.18		-98.73
Southern Gulf St Vincent (40, 43, 44A)		2025										2026							
	King George Whiting - SGSV	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	Total catch (t) (3 year average)	0.34	0.19	0.52	0.85	0.94	0.90	0.96	0.50	conf	conf	conf	0.11	0.34	0.19	0.52	0.85	0.94	0.90
	Total catch (t) (Algal bloom period)	conf	conf	conf	conf	0.60	0.49	1.35	conf	0.01	0.00	0.00	0.02	0.04	0.00	0.01	0.00	0.05	0.04
	Percentage of baseline					-36.40	-45.52	40.45					-82.27	-88.20	-100.00	-98.06	-100.00	-94.70	-95.55
	Southern Calamari - SGSV																		
	Total catch (t) (3 year average)	1.66	0.78	1.36	0.75	1.22	2.02	5.56	10.56	13.76	8.73	8.39	6.11	1.66	0.78	1.36	0.75	1.22	2.02
	Total catch (t) (Algal bloom period)	2.96	1.66	conf	conf	0.00	0.00	conf	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.01	0.01	0.03
	Percentage of baseline	78.63	112.76			-100.00	-100.00		-100.00	-100.00	-100.00	-100.00	-100.00	-100.00	-100.00	-98.52	-98.67	-99.18	-98.51
	Southern Garfish - SGSV																		
	Total catch (t) (3 year average)	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	0.16	0.20	conf	conf	conf	conf
	Total catch (t) (Algal bloom period)	conf	NA	NA	NA	NA	NA	NA	NA	0.00	NA	0.00	0.01	0.00	NA	NA	0.00	0.05	0.06
	Percentage of baseline													-100.00					
Kangaroo Island (39B, 41, 42)		2025										2026							
	King George Whiting - KI	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	Total catch (t) (3 year average)	1.17	0.91	1.06	1.12	1.00	1.31	1.11	1.51	1.04	1.06	1.41	1.48	1.17	0.91	1.06	1.12	1.00	1.31
	Total catch (t) (Algal bloom period)	1.12	1.16	conf	conf	conf	conf	conf	0.35	0.00	0.02	NA	0.03	0.06	0.09	0.03	0.01	0.01	0.01
	Percentage of baseline	-4.33	27.10						-76.78	-100.00	-98.11		-97.98	-94.87	-90.14	-97.16	-99.11	-99.00	-99.23
	Southern Calamari - KI																		
	Total catch (t) (3 year average)	0.25	0.53	0.75	0.26	0.16	0.36	0.33	0.16	0.07	conf	conf	0.11	0.25	0.53	0.75	0.26	0.16	0.36
	Total catch (t) (Algal bloom period)	conf	conf	NA	conf	conf	conf	conf	conf	NA	0.00	NA	0.00	0.00	0.00	0.00	NA	0.01	0.01
	Percentage of baseline												-100.00	-100.00	-100.00	-100.00		-93.57	-97.19
	Southern Garfish - KI																		
	Total catch (t) (3 year average)	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	0.27	0.27	conf	conf	conf	conf
	Total catch (t) (Algal bloom period)	conf	conf	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.01	NA	NA	0.02	NA	NA
	Percentage of baseline													-96.29					

OFFICIAL

Table 4. Regional commercial fishery statistics for King George Whiting, Southern Calamari and Southern Garfish in the Gulf St Vincent/Kangaroo Island fishing zone. Data is targeted CPUE (kg/fisher-day) during the algal bloom period (March 2025 onwards) compared to the 3-year monthly average (February 2022 – February 2025). **Green** reflects change in CPUE by <50% of baseline (3-year average). **Orange** reflects a change in CPUE of 50-80% of baseline (3-year average). **Red** reflects change in CPUE of >80% of baseline (3-year average).

Target CPUE																			
Northern Gulf St Vincent (34, 35, 36)		2025										2026							
King George Whiting - NGSV		Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	CPUE (kg/fisher-days) (3 year average)	conf	8.21	8.45	10.88	13.46	11.68	13.43	14.47	16.46	17.45	conf	conf	14.16	8.21	8.45	10.88	13.46	11.68
	CPUE (kg/fisher-days) (Algal bloom period)	conf	conf	conf	29.01	conf	33.16	conf	3.30	1.29	1.55	2.19	0.61	0.46	3.38	0.31	1.07	2.68	0.20
	Percentage of baseline				166.67		183.98		-77.20	-92.16	-91.12			-96.75	-58.84	-96.33	-90.16	-80.10	-98.29
Southern Calamari - NGSV																			
	CPUE (kg/fisher-days) (3 year average)	23.81	30.14	26.84	26.42	22.29	30.00	27.93	21.12	20.97	14.63	20.15	17.42	23.81	30.14	26.84	26.42	22.29	30.00
	CPUE (kg/fisher-days) (Algal bloom period)	30.82	51.03	51.08	37.12	conf	conf	NA	0.00	0.00	0.00	0.00	0.00	0.05	1.27	1.80	0.44	1.40	1.30
	Percentage of baseline	29.47	69.29	90.32	40.52				-100.00	-100.00	-100.00	-100.00	-100.00	-99.79	-95.79	-93.29	-98.33	-93.72	-95.67
Southern Garfish - NGSV																			
	CPUE (kg/fisher-days) (3 year average)	55.43	75.40	81.26	98.67	114.69	80.72	59.09	36.54	43.78	58.01	73.76	69.54	55.43	75.40	81.26	98.67	114.69	80.72
	CPUE (kg/fisher-days) (Algal bloom period)	74.74	106.25	99.57	conf	55.69	conf	conf	conf	0.23	0.00	0.10	0.77	3.20	3.97	0.90	7.99	NA	17.75
	Percentage of baseline	34.83	40.92	22.53		-51.44				-99.47	-100.00	-99.86	-98.89	-94.23	-94.73	-98.89	-91.90		-78.01
Southern Gulf St Vincent (40, 43, 44A)		2025										2026							
King George Whiting - SGSV		Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	CPUE (kg/fisher-days) (3 year average)	25.87	24.75	27.41	36.13	29.01	28.10	41.85	56.01	conf	conf	conf	26.57	25.87	24.75	27.41	36.13	29.01	28.10
	CPUE (kg/fisher-days) (Algal bloom period)	conf	conf	conf	conf	conf	28.45	36.49	conf	6.80	2.90	0.05	6.67	21.00	0.00	2.50	0.23	16.30	8.18
	Percentage of baseline						1.24	-12.80					-74.90	-18.83	-100.00	-90.88	-99.36	-43.82	-70.89
Southern Calamari - SGSV																			
	CPUE (kg/fisher-days) (3 year average)	23.40	18.01	20.09	16.78	17.62	24.59	40.78	51.40	58.58	48.10	46.17	41.47	23.40	18.01	20.09	16.78	17.62	24.59
	CPUE (kg/fisher-days) (Algal bloom period)	37.03	36.11	conf	conf	conf	0.00	conf	0.00	0.00	0.00	0.00	0.10	0.16	0.25	5.00	2.90	1.97	6.80
	Percentage of baseline	58.25	100.50				-100.00		-100.00	-100.00	-100.00	-100.00	-99.76	-99.32	-98.61	-75.11	-82.72	-88.82	-72.35
Southern Garfish - SGSV																			
	CPUE (kg/fisher-days) (3 year average)	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	38.75	131.00	conf	conf	conf	conf
	CPUE (kg/fisher-days) (Algal bloom period)	conf	NA	NA	NA	NA	NA	NA	NA	0.35	NA	0.00	10.00	1.03	NA	NA	1.00	50.00	60.00
	Percentage of baseline													-97.34					
Kangaroo Island (39B, 41, 42)		2025										2026							
King George Whiting - KI		Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	CPUE (kg/fisher-days) (3 year average)	14.69	10.82	13.22	16.79	17.56	19.66	17.43	23.31	17.33	17.36	16.85	24.89	14.69	10.82	13.22	16.79	17.56	19.66
	CPUE (kg/fisher-days) (Algal bloom period)	33.27	20.50	conf	conf	conf	conf	conf	7.37	2.45	3.12	NA	4.67	4.12	28.67	2.48	2.90	3.90	2.28
	Percentage of baseline	126.46	89.48						-68.39	-85.86	-82.03		-81.24	-71.96	165.00	-81.24	-82.73	-77.79	-88.40
Southern Calamari - KI																			
	CPUE (kg/fisher-days) (3 year average)	10.53	11.70	15.26	8.25	conf	11.66	15.84	18.75	5.72	conf	conf	6.79	10.53	11.70	15.26	8.25	conf	11.66
	CPUE (kg/fisher-days) (Algal bloom period)	conf	conf	NA	conf	conf	conf	conf	conf	NA	0.00	NA	0.00	0.33	0.25	0.00	NA	10.00	11.50
	Percentage of baseline												-100.00	-96.87	-97.86	-100.00			-1.35
Southern Garfish - KI																			
	CPUE (kg/fisher-days) (3 year average)	conf	conf	conf	NA	conf	conf	conf	conf	conf	conf	conf	conf	86.61	95.17	conf	NA	conf	conf
	CPUE (kg/fisher-days) (Algal bloom period)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	3.20	NA	NA	20.00	NA	NA
	Percentage of baseline													-96.31					

OFFICIAL

Table 5. Regional commercial fishery statistics for King George Whiting, Southern Calamari and Southern Garfish in the Spencer Gulf fishing zone. Data is total catch (t) during the algal bloom period (March 2025 onwards) compared to the 3-year monthly average (February 2022 – February 2025). **Green** reflects change in catch by <50% of baseline (3-year average). **Orange** reflects a change in catch of 50-80% of baseline (3-year average). **Red** reflects change in catch of >80% of baseline (3-year average).

TOTAL CATCH																			
Northern Spencer Gulf (11, 19, 20, 21, 22, 23)		2025										2026							
 King George Whiting - NSG		Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	Total catch (t) (3 year average)	0.43	0.91	1.09	4.02	6.35	3.68	1.37	0.62	0.38	0.22	0.19	0.33	0.43	0.91	1.09	4.02	6.35	3.68
	Total catch (t) (Algal bloom period)	0.20	0.75	1.36	4.18	5.12	5.00	1.75	0.75	0.15	0.19	0.19	0.17	0.43	0.36	1.50	5.02	3.42	1.21
	Percentage of baseline	-53.00	-17.61	24.33	4.07	-19.37	35.74	27.96	20.65	-60.90	-14.63	2.54	-49.03	1.05	-60.45	37.13	24.98	-46.14	-67.15
 Southern Calamari - NSG																			
	Total catch (t) (3 year average)	2.70	9.24	12.23	8.18	8.78	7.63	6.05	2.32	1.52	1.98	1.04	1.59	2.70	9.24	12.23	8.18	8.78	7.63
	Total catch (t) (Algal bloom period)	2.10	11.99	13.13	16.12	11.60	13.79	7.23	2.04	0.65	0.22	0.05	0.04	0.21	0.76	0.02	NA	0.04	0.49
	Percentage of baseline	-22.22	29.72	7.33	97.06	32.08	80.69	19.60	-11.90	-57.31	-88.91	-95.21	-97.48	-92.22	-91.78	-99.84		-99.54	-93.58
 Southern Garfish - NSG																			
	Total catch (t) (3 year average)	5.91	5.29	7.86	4.99	4.78	7.49	3.85	3.91	5.27	4.15	8.92	9.17	5.91	5.29	7.86	4.99	4.78	7.49
	Total catch (t) (Algal bloom period)	7.17	5.09	6.59	4.67	3.63	6.45	12.04	4.29	5.77	5.12	8.77	9.48	12.79	9.02	9.08	6.18	8.30	5.08
	Percentage of baseline	21.25	-3.75	-16.20	-6.37	-24.12	-13.88	212.74	9.83	9.59	23.28	-1.68	3.33	116.29	70.56	15.46	23.90	73.50	-32.17
Southern Spencer Gulf (29, 30, 31, 32, 33, 39A)		2025										2026							
 King George Whiting - SSG		Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	Total catch (t) (3 year average)	3.36	1.48	1.85	4.59	7.28	6.96	5.13	3.75	4.49	3.57	4.63	4.78	3.36	1.48	1.85	4.59	7.28	6.96
	Total catch (t) (Algal bloom period)	2.25	0.89	1.39	6.94	8.75	7.00	6.16	4.09	3.87	5.57	3.37	3.81	4.32	1.06	1.58	4.30	8.76	4.93
	Percentage of baseline	-33.12	-39.72	-24.87	51.28	20.13	0.56	20.20	8.93	-13.79	55.85	-27.24	-20.31	28.42	-28.21	-14.60	-6.27	20.26	-29.18
 Southern Calamari - SSG																			
	Total catch (t) (3 year average)	5.07	3.80	4.26	1.80	2.38	3.14	10.09	17.28	14.28	12.83	11.98	7.52	5.07	3.80	4.26	1.80	2.38	3.14
	Total catch (t) (Algal bloom period)	4.87	6.50	5.16	1.29	0.63	0.67	1.76	6.50	7.95	3.58	1.59	0.44	0.25	1.00	0.04	0.01	0.05	0.15
	Percentage of baseline	-3.85	71.22	21.12	-28.50	-73.54	-78.65	-82.56	-62.39	-44.31	-72.09	-86.72	-94.15	-95.06	-73.66	-99.06	-99.45	-97.90	-95.22
 Southern Garfish - SSG																			
	Total catch (t) (3 year average)	0.45	0.62	1.01	0.75	0.27	conf	0.07	0.21	0.49	0.63	1.46	0.86	0.45	0.62	1.01	0.75	0.27	conf
	Total catch (t) (Algal bloom period)	0.41	0.97	conf	conf	conf	conf	conf	0.25	conf	1.64	1.12	1.14	1.45	0.31	conf	conf	conf	conf
	Percentage of baseline	-8.54	55.76							17.19		159.56	-23.09	32.55	223.46	-50.22			
Lower Eyre Peninsula (26, 27, 28, 37, 38, 47)		2025										2026							
 King George Whiting - LEP		Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
	Total catch (t) (3 year average)	0.22	0.15	0.89	0.96	0.49	0.62	0.68	0.59	0.38	0.30	0.12	0.17	0.22	0.15	0.89	0.96	0.49	0.62
	Total catch (t) (Algal bloom period)	0.17	0.64	1.60	conf	0.58	conf	conf	conf	0.85	conf	0.06	conf	0.50	0.47	1.06	1.19	1.41	0.97
	Percentage of baseline	-23.79	316.85	80.70		19.55				123.45		-49.12		124.15	206.12	19.72	23.51	190.62	56.75
 Southern Calamari - LEP																			
	Total catch (t) (3 year average)	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf
	Total catch (t) (Algal bloom period)	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	NA	conf	conf	NA	NA	NA	NA
	Percentage of baseline																		
 Southern Garfish - LEP																			
	Total catch (t) (3 year average)	conf	conf	conf	conf	conf	NA	NA	NA	conf	NA	NA	NA	conf	conf	conf	conf	conf	NA
	Total catch (t) (Algal bloom period)	NA	conf	conf	NA	NA	conf	NA	conf	NA	conf	NA	conf	NA	NA	conf	conf	conf	NA
	Percentage of baseline																		

OFFICIAL

Table 6. Regional commercial fishery statistics for King George Whiting, Southern Calamari and Southern Garfish in the Spencer Gulf fishing zone. Data is targeted CPUE (kg/fisher-day) during the algal bloom period (March 2025 onwards) compared to the 3-year monthly average (February 2022 – February 2025). **Green** reflects change in CPUE by <50% of baseline (3-year average). **Orange** reflects a change in CPUE of 50-80% of baseline (3-year average). **Red** reflects change in CPUE of >80% of baseline (3-year average).

Target CPUE																		
Northern Spencer Gulf (11, 19, 20, 21, 22, 23)			2025									2026						
King George Whiting - NSG			Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun
	CPUE (kg/fisher-days) (3 year average)		17.17	12.34	14.04	42.16	37.97	28.15	15.31	15.80	18.54	conf	14.92	17.32	17.17	12.34	14.04	42.16
	CPUE (kg/fisher-days) (Algal bloom period)		conf	12.05	12.51	29.68	48.07	39.04	13.12	7.70	5.82	4.85	conf	7.21	7.36	5.74	10.52	26.13
	Percentage of baseline			-2.33	-10.88	-29.61	26.61	38.70	-14.29	-51.26	-68.60			-58.38	-57.14	-53.47	-25.05	-38.03
Southern Calamari - NSG			2025									2026						
	CPUE (kg/fisher-days) (3 year average)		22.47	32.81	38.71	41.52	34.46	29.24	35.96	28.36	25.42	28.91	17.48	19.43	22.47	32.81	38.71	41.52
	CPUE (kg/fisher-days) (Algal bloom period)		16.70	35.50	36.89	57.95	47.59	41.98	39.02	18.99	15.62	5.73	conf	conf	1.81	8.96	8.00	NA
	Percentage of baseline		-25.69	8.19	-4.70	39.58	38.09	43.57	8.51	-33.03	-38.56	-80.18			-91.95	-72.69	-79.33	-78.01
Southern Garfish - NSG			2025									2026						
	CPUE (kg/fisher-days) (3 year average)		55.72	51.85	64.52	53.66	60.23	75.74	55.61	43.97	54.41	48.02	69.27	68.82	55.72	51.85	64.52	53.66
	CPUE (kg/fisher-days) (Algal bloom period)		66.11	56.89	62.43	81.51	44.38	62.67	155.54	48.40	43.78	38.02	56.75	53.12	48.56	50.06	62.81	60.52
	Percentage of baseline		18.64	9.72	-3.24	51.90	-26.32	-17.25	179.71	10.09	-19.54	-20.83	-18.08	-22.81	-12.85	-3.46	-2.65	12.78
Southern Spencer Gulf (29, 30, 31, 32, 33, 39A)			2025									2026						
King George Whiting - SSG			Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun
	CPUE (kg/fisher-days) (3 year average)		18.26	14.52	17.48	24.67	26.63	26.54	24.89	26.41	28.94	23.91	22.48	23.98	18.26	14.52	17.48	24.67
	CPUE (kg/fisher-days) (Algal bloom period)		19.69	9.83	13.77	30.16	35.68	29.12	27.50	21.41	26.86	24.59	23.02	22.04	22.15	9.09	12.06	22.26
	Percentage of baseline		7.85	-32.30	-21.23	22.27	33.98	9.72	10.50	-18.93	-7.18	2.85	2.42	-8.08	21.32	-37.40	-31.01	-9.76
Southern Calamari - SSG			2025									2026						
	CPUE (kg/fisher-days) (3 year average)		23.44	20.34	21.13	18.94	20.82	23.12	36.32	46.35	49.80	48.53	39.32	31.47	23.44	20.34	21.13	18.94
	CPUE (kg/fisher-days) (Algal bloom period)		28.85	34.62	33.35	17.89	14.07	10.88	17.93	28.75	37.10	22.02	14.92	8.18	4.62	12.81	8.75	0.38
	Percentage of baseline		23.09	70.22	57.84	-5.55	-32.43	-52.94	-50.63	-37.97	-25.50	-54.63	-62.06	-74.01	-80.29	-37.02	-58.59	-97.99
Southern Garfish - SSG			2025									2026						
	CPUE (kg/fisher-days) (3 year average)		31.59	52.63	69.95	84.56	71.19	conf	28.17	39.81	63.54	39.06	55.80	41.89	31.59	52.63	69.95	84.56
	CPUE (kg/fisher-days) (Algal bloom period)		40.44	64.44	conf	conf	conf	conf	conf	conf	conf	53.91	43.17	52.56	51.93	19.29	conf	conf
	Percentage of baseline		28.03	22.45								38.02	-22.63	25.46	64.40	-63.35		
Lower Eyre Peninsula (26, 27, 28, 37, 38, 47)			2025									2026						
King George Whiting - LEP			Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun
	CPUE (kg/fisher-days) (3 year average)		20.20	10.73	22.06	23.27	21.10	25.27	22.87	23.68	15.88	13.43	10.52	14.37	20.20	10.73	22.06	23.27
	CPUE (kg/fisher-days) (Algal bloom period)		10.61	16.49	29.71	conf	32.24	conf	conf	conf	20.35	conf	6.96	conf	20.08	13.55	17.66	25.31
	Percentage of baseline		-47.49	53.71	34.68		52.79				28.19		-33.86		-0.61	26.30	-19.94	8.77
Southern Calamari - LEP			2025									2026						
	CPUE (kg/fisher-days) (3 year average)		conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf	conf
	CPUE (kg/fisher-days) (Algal bloom period)		conf	conf	conf	conf	conf	conf	conf	conf	conf	NA	conf	NA	conf	conf	NA	NA
	Percentage of baseline																	
Southern Garfish - LEP			2025									2026						
	CPUE (kg/fisher-days) (3 year average)		conf	conf	conf	NA	conf	NA	NA	NA	NA	NA	NA	NA	conf	conf	conf	NA
	CPUE (kg/fisher-days) (Algal bloom period)		NA	NA	NA	NA	NA	NA	NA	conf	NA	conf	NA	conf	NA	NA	NA	conf
	Percentage of baseline																	